National Summary for Article 17 - Finland

1 General information

1.1 Number of SCIs and SACs

The table below provides the total number and total area of sites proposed and designated under the Habitats Directive (Sites of Community Importance, SCIs & Special Areas of Conservation, SACs), terrestrial area of sites and number and area of marine sites (i.e. any site with a marine component).

Empty cells in tables mean that the component requested is not applicable.

| | All | | Terrestrial | Marine | | | |
|-----------------------------------|------|------------|-------------|--------|------------|--|--|
| | No. | Area (km²) | Area (km²) | No. | Area (km²) | | |
| SCIs & SACs | 1721 | 48439 | 42290 | 172 | 6149 | | |
| SACs only | 0 | | | 0 | | | |
| Date of database used: 28-09-2012 | | | | | | | |

1.2 Number of sites with comprehensive management plans (Art. 6(1))

Number of sites for which comprehensive management plans have been adopted: 314

Percentage of network area covered by comprehensive management plans: 78%

Number of sites for which management plans are under preparation (optional): not reported

2. Number of habitats and species/subspecies

The table in this section gives the number of habitat types and species/subspecies in each Annex of the Habitats Directive by biogeographical and marine regions in Finland. The species and habitats with the following presence status are included in the table: 'present', species of which taxonomy is not clear (SR TAX), species where the link to the corresponding name in the Habitats Directive is not clear (LR), species extinct after the Directive came into force (EX) and optional reports (OP).

| Region | HABI | TATS | SPECIES | | | | | | | |
|----------------------|--------------|----------|--------------|----------|-----------------------------------|-----------------------------------|-----------------------------------|-----------------------------------|--|--|
| Region | Ann | ex I | Anne | Annex II | | Annex IV | | Annex V | | |
| | Non-priority | Priority | Non-priority | Priority | Including those in Annex II | Excluding those in Annex II | Including those in Annex II | Excluding those in Annex II | | |
| Number of habitats & | 48 | 21 | 78 | 7 | 68 | 20 | 21 | 18 | | |
| species in the MS | 69 | | 85 | | 68 | | 21 | | | |
| Alpine | 18 | 7 | 19 | 2 | 13 | 3 | 11 | 10 | | |
| Boreal | 42 | 20 | 68 | 6 | 64 | 20 | 18 | 17 | | |
| Marine Baltic | 5 | | 2 | | | | 2 | | | |

Additional information:

Number of assessments of marginal habitat types: 6

Number of assessments of marginal & occasional species: 14

Number of assessments of newly arriving species: 3

Number of species regionally extinct prior the Habitats Directive came into force: **none** Number of species regionally extinct after the Habitats Directive came into force: **none** Number of species globally extinct after the Habitats Directive came into force: **none** Number of assessments of species/habitat types for which no reports received: **none**

3. Information on Conservation status


Please note that the figures shown for 2001-2006 and 2007-2012 are not necessarily directly comparable because there can be differences in number of assessments between the reporting rounds, changes in how some features were allocated in biogeographical regions etc.


The following have been excluded from all statistics under section 3:

- Habitats reported as marginal (MAR) or with scientific reserve (SR)
- Species reported as marginal (MAR), occasional (OCC), newly arriving (ARR), regionally extinct before the Habitats Directive came into force (PEX) and introduced species (INT). In addition reports that give only an information about species without evaluation of the conservation status
- Redundant reports provided for both marine and terrestrial regions for habitats and species and species for which only one, either terrestrial or marine report was expected (IRM).

3.1 a) Overall assessment of conservation status of habitats and species (%)

These figures show the percentage of biogeographical assessments in each category of conservation status for habitats and species, respectively. The information on which these figures are based are presented in the table below the figures (real values).





Conservation status of habitats

Conservation status of species

FV - Favourable NA - Not reported XX - Unknown U1 - Unfavourable inadequate U2 - Unfavourable bad

| Year of HABITATS | | | | | | SPECIES | | | | |
|------------------|----|----|----|----|----|---------|----|----|----|----|
| assessment | FV | NA | xx | U1 | U2 | FV | NA | xx | U1 | U2 |
| 2007 | 31 | 1 | | 36 | 24 | 65 | | 16 | 49 | 14 |
| 2013 | 31 | | 1 | 36 | 24 | 69 | | 11 | 53 | 17 |


3.1 b) Percentage of assessments where the conservation status has changed between the reporting periods


This table shows the percentage of assessments where the Member State has indicated a change between two reporting rounds (2001-2006 and 2007-2012) and the percentages of all reported changes where the change has been reported as a genuine change. Change can be either a change from one conservation status category to another or a change within the same category (within the qualifiers '-', '+'. '=', 'x'). Data have been taken from the 'audit trail table' where the Member State indicates the nature of change. The Member State's results on this audit trail are shown under section 7.

| | SPECIES | HABITAT TYPES |
|---------------------------------------|---------|---------------|
| % of assessments that changed | 31% | 10% |
| % of total changes considered genuine | 10% | 2% |


3.2 Improving/deteriorating trends of habitats and species with an unfavourable conservation status (%)

These figures show the proportion of unfavourable assessments (U1 & U2) which are improving, deteriorating, stable or unknown.





Habitats - overall trend in Conservation Status



U (+) = unfavourable (inadequate and bad) improving, U (=) = unfavourable stable, U (-) = unfavourable declining, U (x) = unfavourable unknown trend

This table shows trends in conservation status of habitats & species separately for those cases where the overall conclusion is unfavourable inadequate (U1) and unfavourable bad (U2).


| Qualifiers of CS | U1+ | U1= | U1- | U1x | U2+ | U2= | U2- | U2x |
|------------------|-----|-----|-----|-----|-----|-----|-----|-----|
| Habitats | 4 | 13 | 19 | | 4 | 4 | 16 | |
| Species | 4 | 23 | 16 | 10 | 2 | 6 | 6 | 3 |

Note: U1+ = unfavourable-inadequate improving, U1= = unfavourable-inadequate stable, U1- = unfavourable-inadequate declining, U1x = unfavourable-inadequate trend unknown, U2+ = unfavourable-bad improving, U2= = unfavourable-bad stable, U2- = unfavourable-bad declining, U2x = unfavourable-bad trend unknown


3.3 Overall assessment of conservation status of habitats and species by biogeographical/marine region (%)

These figures show the percentage of assessments in each of conservation status category by biogeographical and marine region, for habitats and species, respectively.

Please note that some habitats reported as terrestrial in 2001-2006 have been reported as marine in 2007-2012 (e.g. estuaries). Some species (e.g. seals, marine turtles) which in some cases were reported for both marine and terrestrial regions were only reported for one region in 2007-2012 (this statement only applies to Member States with marine regions).



Conservation status of habitats in biogeographical and marine regions



Conservation status of **species** in biogeographical and marine regions


Note: wide bar corresponds to the 2007-2012 reporting period, and the narrow bar to the 2001-2006 reporting period. The number in brackets corresponds to the number of biogeographical assessments in the category.

3.4 Overall assessment of conservation status by habitat category/species group (%)

These figures show the percentage of biogeographical and marine assessments in each conservation status category by habitat category and by taxonomic group, for habitats and species, respectively.

The figures show the proportion of assessments in each conservation status class for 2007-2012 (upper bar) and 2001-2006 (lower bar). The information (number of assessments) on which these figures are based are presented in the tables below each figure (real values).

Habitats




Conservation status of habitats in biogeographical and marine regions

Note: wide bar corresponds to the 2007-2012 reporting period, and the narrow bar to the 2001-2006 reporting period. The number in brackets corresponds to the number of biogeographical assessments in the category.

| Grave | Year of | | | HABITATS | | |
|---------------------|------------|----|----|----------|----|----|
| Group | assessment | FV | NA | XX | U1 | U2 |
| Forests | 2007 | 4 | | | 9 | 4 |
| | 2013 | 4 | | | 9 | 4 |
| Rocky habitats | 2007 | 6 | | | 1 | |
| | 2013 | 6 | | | 1 | |
| Bogs, mires & fens | 2007 | 6 | 1 | | 6 | 3 |
| | 2013 | 6 | | 1 | 6 | 3 |
| Grasslands | 2007 | 3 | | | 1 | 9 |
| | 2013 | 3 | | | 1 | 9 |
| Heath & scrub | 2007 | 2 | | | 2 | 1 |
| | 2013 | 2 | | | 2 | 1 |
| Freshwater habitats | 2007 | 7 | | | 5 | 2 |
| | 2013 | 7 | | | 5 | 2 |
| Dunes habitats | 2007 | | | | 5 | 2 |
| | 2013 | | | | 5 | 2 |
| Coastal habitats | 2007 | 3 | | | 7 | 3 |
| | 2013 | 3 | | | 7 | 3 |

NB: Coastal habitats cover coastal and halophytic habitats (code 1xxx) and Dunes habitat types cover coastal sand dunes and inland dunes (code 2xxx) as listed in the Habitats Directive

Species



Conservation status of species in biogeographical and marine regions

Note: wide bar corresponds to the 2007-2012 reporting period, and the narrow bar to the 2001-2006 reporting period. The number in brackets corresponds to the number of biogeographical assessments in the category.

| Crew | Year of | | | SPECIES | | |
|---------------------|------------|----|----|---------|----|----|
| Group | assessment | FV | NA | XX | U1 | U2 |
| Other invertebrates | 2007 | | | 1 | | |
| | 2013 | | | 1 | | |
| Mammals | 2007 | 19 | | 6 | 3 | 2 |
| | 2013 | 15 | | 5 | 9 | 2 |
| Reptiles | 2007 | | | 1 | | |
| | 2013 | | | | 1 | |
| Amphibians | 2007 | 3 | | | 1 | |
| | 2013 | 3 | | | 1 | |
| Fish | 2007 | 9 | | 1 | | |
| | 2013 | 10 | | | 2 | |
| Arthropods | 2007 | 11 | | 1 | 20 | 3 |
| | 2013 | 15 | | 3 | 15 | 3 |
| Molluscs | 2007 | 2 | | 3 | 1 | |
| | 2013 | 3 | | 2 | | 2 |
| Vascular plants | 2007 | 17 | | | 12 | 8 |
| | 2013 | 17 | | | 14 | 7 |
| Non-vascular plants | 2007 | 4 | | 3 | 12 | 1 |
| | 2013 | 6 | | | 11 | 3 |

3.5 Reasons for change in reported values of parameters (%)

This table provides information on reasons for changes of values reported for the parameters 'Range', 'Area (habitat)', 'Population' and 'Habitat for the species' between reporting periods 2001-2006 and 2007-2012. The table gives the percentage of habitats/species assessments for which a particular reason for change in values was reported. The reporting format lists three principal reasons for change: genuine change, better knowledge/data and use of different method.

| Deepen for shores | Hab | itats | Species/subspecies | | | |
|-------------------------|-----------------------|-------------------------|-----------------------|-----------------|---------------------------------|--|
| Reason for change | Surface area of range | Surface area of habitat | Surface area of range | Population size | Area of habitat for the species | |
| Genuine change | 1 | 9 | 13 | 17 | 9 | |
| Better knowledge/data | 65 | 49 | 49 | 45 | 37 | |
| Use of different method | 97 | 28 | 87 | 59 | 43 | |

Note: More than one reason for change can be reported for each habitat and species.

4 Frequency of main pressures and threats (%)¹


This section provides information on the relative importance of pressures and threats (aggregated to level 1) reported for habitats and species. The figures show the percentage of biogeographical assessments reported as being affected by one or more pressures or threats categorised as of 'high importance'. The information for the number of pressures and threats on which these figures are based are presented in the tables below the figures.


¹ The following have been excluded:

[•] Habitats reported as marginal or with scientific reserve.

[•] Species reported as marginal, occasional, newly arriving, regionally extinct before the Habitats Directive came into force and introduced species. In addition reports that give only an information about species without evaluation of the conservation status.

Redundant reports provided for both marine and terrestrial regions for habitats and species and species for which only
one, either terrestrial or marine report was expected.





pressure threat


Note: Threats and pressures categories not reported are omitted.


Total number of assessments considered in the calculation: 92

Number of assessments with no high ranking threats (or no threats at all reported): 48

Number of assessment with no high ranking pressures (or no pressures at all): 47

| Pressures and threats | HABI | TATS |
|---|-------------------|------------------------|
| Pressures and threats | Number of threats | Number of pressures |
| A - Agriculture | 14 | 14 |
| B - Sylviculture, forestry | 10 | 11 |
| C - Mining, extraction of materials and energy production | 1 | |
| D - Transportation and service corridors | 2 | 1 |
| E - Urbanisation, residential and commercial development | 1 | 1 |
| G - Human intrusions and disturbances | 5 | 5 |
| H - Pollution | 9 | 10 |
| I - Invasive, other problematic species and genes | 2 | 2 |
| J - Natural System modifications | 9 | 11 |
| K - Natural biotic and abiotic processes (without catastrophes) | 9 | 7 |
| M - Climate change | 2 | |





pressure threat

Note: Threats and pressures categories not reported are omitted.

Total number of assessments considered in the calculation: 150

Number of assessments with no high ranking threats (or no threats at all reported): 117

Number of assessment with no high ranking pressures (or no pressures at all): 113

| Pressures and threats | SPE | CIES |
|---|-------------------|---------------------|
| Pressures and threats | Number of threats | Number of pressures |
| A - Agriculture | 5 | 5 |
| B - Sylviculture, forestry | 7 | 7 |
| C - Mining, extraction of materials and energy production | 1 | 3 |
| D - Transportation and service corridors | 3 | |
| E - Urbanisation, residential and commercial development | 1 | 1 |
| F - Biological resource use other than agriculture & forestry | 5 | 6 |
| G - Human intrusions and disturbances | | 2 |
| H - Pollution | | 3 |
| I - Invasive, other problematic species and genes | 4 | 3 |
| J - Natural System modifications | 10 | 10 |
| K - Natural biotic and abiotic processes (without catastrophes) | 9 | 10 |
| M - Climate change | 3 | 1 |
| U - Unknown threat or pressure | 1 | 1 |
| X - No threats or pressures | 1 | 1 |

5 Natura 2000 coverage and conservation measures ²

Note: The figures under section 5 cover only Annex I habitat types and Annex II species.

5.1 Natura 2000 coverage (%)


This section presents statistics on the coverage of Annex I habitats and Annex II species in Natura 2000 sites by habitat category/species group. These figures show the percentage of habitats/species assessments in three classes based on coverage by Natura 2000 sites, for habitats and species, respectively. The geometric mean is used if Member States have reported minimum and maximum values. The information for the number of assessments per coverage by Natura 2000 on which these figures are based are presented in the tables below the figures (real values). Please note that these statistics are based on Article 17 data and are independent from the results of the Biogeographical Seminars.

² The following have been excluded:

[•] Habitats reported as marginal or with scientific reserve.

[•] Species reported as marginal, occasional, newly arriving, regionally extinct before the Habitats Directive came into force and introduced species. In addition reports that give only an information about species without evaluation of the conservation status.

Redundant reports provided for both marine and terrestrial regions for habitats and species and species for which only
one, either terrestrial or marine report was expected.




% of habitat assessments in 3 classes of coverage by Natura 2000 sites

coverage by Natura 2000 sites : ■ 0-24% ■ 25-74% ■ 75-100%

Note: The number in brackets corresponds to the number of biogeographical assessments in the habitat category.

| Group | | HABITATS | | | | | | | |
|---------------------|-------|----------|---------|---------|--|--|--|--|--|
| Group | 0-24% | 25-74% | 75-100% | unknown | | | | | |
| Forests | 3 | 8 | 4 | 2 | | | | | |
| Rocky habitats | 1 | 2 | 4 | | | | | | |
| Bogs, mires & fens | 3 | 6 | 7 | | | | | | |
| Grasslands | 4 | 4 | 5 | | | | | | |
| Heath & scrub | | 1 | 4 | | | | | | |
| Freshwater habitats | 4 | 3 | 5 | 2 | | | | | |
| Dunes habitats | 1 | 4 | 2 | | | | | | |
| Coastal habitats | 5 | 8 | | | | | | | |



% of species assessments in 3 classes of coverage by Natura 2000 sites


coverage by Natura 2000 sites : • 0-24% • 25-74% • 75-100%

Note: The number in brackets corresponds to the number of biogeographical assessments in the species category.

| Group | | SPECIES | | | | | | | |
|---------------------|-------|---------|---------|---------|--|--|--|--|--|
| Group | 0-24% | 25-74% | 75-100% | unknown | | | | | |
| Mammals | 3 | 3 | 4 | | | | | | |
| Amphibians | | 1 | | | | | | | |
| Arthropods | 9 | 8 | 10 | 1 | | | | | |
| Molluscs | 1 | 1 | 5 | | | | | | |
| Vascular plants | 5 | 8 | 23 | | | | | | |
| Non-vascular plants | | 5 | 10 | | | | | | |

5.2 Main conservation measures (%)

This section provides information on the relative importance of conservation measures at level 1 implemented during the reporting period 2007-2012 for Annex I habitats and Annex II species. The figures show the percentage of biogeographical assessments for which one or more 'high importance' conservation measures was implemented. Measures not reported are omitted.




% of **habitat assessments** for which one or more 'high' importance measures were reported

Note: Numbers in brackets correspond to the number of assessments where measure 1, 2, etc. is noted as being of high importance. Occasional and extinct habitat types have been included in calculations.

Total number of assessments considered in the calculation: 92

Number of assessments with no high ranking conservation measures or no conservation measures at all reported: **18**



% of **species assessments** for which one or more 'high' importance measures were reported


Note: Numbers in brackets correspond to the number of assessments where measure 1, 2, etc. is noted as being of high importance. Occasional and extinct species have been included in calculations.

Total number of assessments considered in the calculation: 97

Number of assessments with no high ranking conservation measures or no conservation measures at all reported: **33**

5.3 Impact of conservation measures (%)

This section provides information on the effects of implemented conservation measures for each level 1 measure category. The figures show, for each level 1 measure category, the frequency of reported effects. The information for the number of assessments per measure category on which these figures are based are presented in the tables below the figures (full names of the measures are shown in the tables).




% of **habitat assessments** for which a particular effect of a measure was reported

■ maintain ■ enhance ■ longterm ■ no effect ■ unknown or not evaluated

Note: The numbers in brackets correspond to the numbers of biogeographical assessments for which one or more 'high' importance measure was reported.

| Measure | | HABITATS | | | | | | |
|---|----|----------|----------|--|--------------------------------|--|--|--|
| | | enhance | longterm | | unknown or not evaluated | | | |
| 2 - Measures related to agriculture and open habitats | | 18 | | | | | | |
| 3 - Measures related to forests and wooded habitats | | 1 | | | | | | |
| 4 - Measures related to wetland, freshwater and coastal habitats | | 6 | | | | | | |
| 6 - Measures related to spatial planning | 49 | 74 | 9 | | | | | |
| 7 - Measures related to hunting, taking and fishing and species management | 1 | 3 | | | | | | |
| 9 - Measures related to special resource use | | 1 | | | | | | |



% of **species assessments** for which a particular effect of a measure was reported

■ maintain ■ enhance ■ longterm ■ no effect ■ unknown or not evaluated

Note: The numbers in brackets correspond to the numbers of biogeographical assessments for which one or more 'high' importance measure was reported.

| | | | SPECIES | ; | |
|--|----|---------|----------|---|--------------------------------|
| Measure | | enhance | longterm | 1 | unknown or not evaluated |
| 1 - No measures | | | | | 1 |
| 2 - Measures related to agriculture and open habitats | | 3 | | | |
| 3 - Measures related to forests and wooded habitats | | 6 | 1 | | |
| 4 - Measures related to wetland, freshwater and coastal habitats | | 2 | | | |
| 6 - Measures related to spatial planning | 25 | 58 | 6 | | 1 |
| 7 - Measures related to hunting, taking and fishing and species management | 4 | 9 | 4 | | |
| 9 - Measures related to special resource use | | 1 | | | |

6 Data quality and completeness ³

The aim of this section is to provide an overview of the data gaps in the report; most of these gaps are due to insufficient knowledge. This section does not refer to potential errors or technical problems in the Member State's report and concentrates on what is relevant for evaluating data completeness.

The tables give percentages of habitats/species assessments with unknown or missing information for components of conservation status and conclusions.

³ The statistics on missing information take into account that for the plant species listed in Annex V at the genus level only 'Overall assessment of conservation status' and 'Overall trend' are mandatory. The same approach was used for the species extinct after the Habitats Directive came into force.

Habitats

| | Area | 0 |
|-----------------------|-----------------|-----|
| Liebitet ronge | Trend | 0 |
| Habitat range | Reference value | 0 |
| | Conclusion | 0 |
| | Area | 0 |
| | Trend | 0 |
| Habitat area | Reference value | 0 |
| | Conclusion | 0 |
| Structure & functions | Conclusion | 0 |
| Future prospects | Conclusion | 0 |
| Pressures | s & threats | 0 |
| Natura 2000 | Coverage | 0 |
| Natura 2000 | Measures | 1.1 |
| | Conclusion | 0 |
| Overall | Trend | 0 |
| | Maps | 0 |

Species

| | | 1 |
|----------------------|---------------------------|---|
| | Area | 0 |
| Species range | Trend | 0 |
| Species range | Reference value | 0 |
| | Conclusion | 0 |
| | Size | 0 |
| Species population | Trend | 0 |
| Species population | Reference value | 0 |
| | Conclusion | 0 |
| | Area | 0 |
| Lichitat far anagiag | Trend | 0 |
| Habitat for species | Area of suitable habitat* | 0 |
| | Conclusion | 0 |
| Future prospects | Conclusion | 0 |
| Pressures | s & threats | 0 |
| Natura 2000 | Coverage | 0 |
| Natura 2000 | Measures | 0 |
| | Conclusion | 0 |
| Overall | Trend | 0 |
| | Maps | 0 |

*This field is a mandatory field in the reporting format, however there is an inconsistency between the reporting format and the evaluation matrix as raised in the FAQ dated 14.2.2013

Habitats

| | Area | 0 |
|-----------------------|-----------------|-----|
| | Trend | 0 |
| Habitat range | Reference value | 1.1 |
| | Conclusion | 2 |
| | Area | 4 |
| | Trend | 2 |
| Habitat area | Reference value | 1.1 |
| | Conclusion | 3 |
| Structure & functions | Conclusion | 1.1 |
| Future prospects | Conclusion | 1.1 |
| Pressures | s & threats | 2 |
| Natura 2000 | Coverage | 0 |
| Natura 2000 | Measures | 0 |
| | Conclusion | 1.1 |
| Overall | Trend | 0 |
| | Maps | 0 |

Species

| | Area | 0 |
|----------------------|---------------------------|----|
| | Trend | 12 |
| Species range | Reference value | 6 |
| | Conclusion | 4 |
| | Size | 0 |
| Charica non-detion | Trend | 37 |
| Species population | Reference value | 13 |
| | Conclusion | 12 |
| | Area | 0 |
| Lichitat for anapian | Trend | 18 |
| Habitat for species | Area of suitable habitat* | 99 |
| | Conclusion | 6 |
| Future prospects | Conclusion | 11 |
| Pressures | s & threats | 5 |
| Natura 2000 | Coverage | 0 |
| Natura 2000 | Measures | 0 |
| | Conclusion | 7 |
| Overall | Trend | 19 |
| | Maps | 0 |

*This field is a mandatory field in the reporting format, however there remained an inconsistency between the reporting format and the evaluation matrix as raised in the FAQ dated 14.2.2013

6.2 Methods used to estimate values or trends in Member State reports (%)

This section presents information about the quality of estimated values and trends in habitat and species biogeographical reports. For some parameters and trends, the reporting format requires an indication of which of three methods (complete survey or a statistically robust estimate, partial data with some extrapolation and/or modelling, expert opinion with no or minimal sampling) have been used to estimate the values or trends. The tables in this section present percentage of habitats/species assessments for which values were estimated by each of the three methods mentioned above.

Habitats

| | Мар | Range | Area | Area trend | Str.&Funct. | N2000 | Average |
|---------------------|-----|-------|------|------------|-------------|-------|---------|
| Expert opinion (%) | 7 | 7 | 8 | 43 | 41 | 5 | 18 |
| Extrapolation (%) | 71 | 71 | 67 | 53 | 53 | 73 | 65 |
| Complete survey (%) | 22 | 22 | 21 | 3 | 7 | 22 | 16 |
| Absent data (%) | 0 | 0 | 4 | 1 | 0 | 0 | 1 |

Species

| | Мар | Range | Population | Pop. trend | Habitat | N2000* | Average |
|---------------------|-----|-------|------------|------------|---------|--------|---------|
| Expert opinion (%) | 2 | 2 | 31 | 24 | 47 | 30 | 23 |
| Extrapolation (%) | 15 | 16 | 52 | 37 | 47 | 55 | 37 |
| Complete survey (%) | 83 | 81 | 17 | 12 | 7 | 15 | 36 |
| Absent data (%) | 0 | 1 | 0 | 27 | 0 | 0 | 5 |

*This column covers only Annex II species

Source of information:

Link to the national general report on CDR

Link to the national report for habitats on CDR

Link to the national report for species on CDR

Other links (national links to be provided by the Member State)

7. List of habitats and species reported and their conservation status

This section lists habitats and species reported by the Member State and the overall conclusions on their conservation status for the reporting period 2001-2006 (indicated as 2007) and 2007-2012 (indicated as 2013). Information from the audit trail has been used for this list and its focus is on what was reported in 2013.

There are two tables for habitats and species if relevant for the Member State. The second table includes only habitats or species with a status OCC, SR, MAR etc. Please note that occurrences e.g. OCC if only reported in 2007, are included only in the second table.

In addition the list includes information provided by the Member State on the nature of change in the overall conservation status between the reporting periods.

The codes are the following :

- a = there is a genuine change: the overall conservation status improved (or deteriorated) due to natural or non-natural reasons (management, intervention, etc.)
- b1 = the change observed is due to more accurate data (e.g. better mapping of distribution) or improved knowledge (e.g. on ecology of species or habitat)
- b2 = the change observed is due to a taxonomic review: one taxon becoming several taxa, or vice versa
- c1 = the change observed is due to use of different methods to measure or evaluate individual parameters or the overall conservation status
- c2 = the change observed is mainly due to the use of different thresholds e.g. to fix Favourable reference values
- d = no information about the nature of change
- e = the change observed is due to less accurate or absent data than the one used in the previous reporting period
- nc = no change (e.g. overall trend in conservation status only evaluated in 2013 but assumed to be the same in 2007 or not known)

| Group | Name | Code | Year | ALP | BOR | MBAL |
|----------------|--|------|--------------|------------|------------------|------|
| Forests | Alluvial forests with Alnus glutinosa and Fraxinus excelsior (Alno-Padion, Alnion incanae, Salicion albae) | 91E0 | 2013 2007 | FV FV | U2- U2- | |
| | Bog woodland | 91D0 | 2013 2007 | FV FV | U1- U1 nc | |
| | Coniferous forests on, or connected to, glaciofluvial eskers | 9060 | 2013 2007 | | U2- U2- | |
| | Fennoscandian deciduous swamp woods | 9080 | 2013 2007 | | U2- U2 nc | |
| | Fennoscandian hemiboreal natural old broad-leaved deciduous forests (Quercus, Tilia, Acer, Fraxinus or | 9020 | 2013 2007 | | U1= U1+ c1 | |
| | Fennoscandian herb-rich forests with Picea abies | 9050 | 2013 2007 | FV FV | U1= U1 nc | |
| | Fennoscandian wooded pastures | 9070 | 2013 2007 | | U2- U2 b1 | |
| | Natural forests of primary succession stages of landupheaval coast | 9030 | 2013 2007 | | U1- U1 nc | |
| | Nordic subalpine/subarctic forests with Betula pubescens ssp. czerepanovii | 9040 | 2013 2007 | U1- U1- | U1= U1- c1 | |
| | Old acidophilous oak woods with Quercus robur on sandy plains | 9190 | 2013 2007 | | U1- U1 nc | |
| | Tilio-Acerion forests of slopes, screes and ravines | 9180 | 2013 2007 | | U1= U1 nc | |
| | Western Taïga | 9010 | 2013 2007 | FV FV | U1= U1 nc | |
| Rocky habitats | Calcareous rocky slopes with chasmophytic vegetation | 8210 | 2013 2007 | FV FV | U1- U1 nc | |

Habitats reported by Finland

| Group | Name | Code | Year | ALP | BOR | MBAL |
|--------------------|---|------|--------------|------------|------------------|------|
| | Siliceous rock with pioneer vegetation of the Sedo-Scleranthion or of the Sedo albi-Veronicion dillenii | 8230 | 2013 2007 | | FV FV | |
| | Siliceous rocky slopes with chasmophytic vegetation | 8220 | 2013 2007 | FV FV | FV FV | |
| | Siliceous scree of the montane to snow levels (Androsacetalia alpinae and Galeopsietalia ladani) | 8110 | 2013 2007 | FV FV | FV FV | |
| Bogs, mires & fens | Aapa mires | 7310 | 2013 2007 | FV FV | U1- U1 nc | |
| | Active raised bogs | 7110 | 2013 2007 | | U2- U2 nc | |
| | Alkaline fens | 7230 | 2013 2007 | FV FV | U1- U1- | |
| | Alpine pioneer formations of the Caricion bicoloris-atrofuscae | 7240 | 2013 2007 | FV FV | | |
| | Calcareous fens with Cladium mariscus and species of the Caricion davallianae | 7210 | 2013 2007 | | U1- U1 a | |
| | Fennoscandian mineral-rich springs and springfens | 7160 | 2013 2007 | FV FV | U2+ U2+ | |
| | Palsa mires | 7320 | 2013 2007 | U1- U1- | U2- U2- | |
| | Petrifying springs with tufa formation (Cratoneurion) | 7220 | 2013 2007 | FV FV | U1= U1 nc | |
| | Transition mires and quaking bogs | 7140 | 2013 2007 | FV FV | U1- U1 nc | |
| Grasslands | Fennoscandian lowland species-rich dry to mesic grasslands | 6270 | 2013 2007 | | U2- U2- | |
| | Fennoscandian wooded meadows | 6530 | 2013 2007 | | U2= U2 b1 | |
| | Hydrophilous tall herb fringe communities of plains and of the montane to alpine levels | 6430 | 2013 2007 | FV FV | U1= U1 b1 | |
| | Lowland hay meadows (Alopecurus pratensis, Sanguisorba officinalis) | 6510 | 2013 2007 | | U2- U2- | |
| | Molinia meadows on calcareous, peaty or clayey-silt-laden soils (Molinion caeruleae) | 6410 | 2013 2007 | | U2= U2- b1 | |
| | Mountain hay meadows | 6520 | 2013 2007 | | U2- U2- | |
| | Nordic alvar and precambrian calcareous flatrocks | 6280 | 2013 2007 | | U2- U2- | |
| | Northern boreal alluvial meadows | 6450 | 2013 2007 | | U2- U2- | |
| | Semi-natural dry grasslands and scrubland facies on calcareous substrates (Festuco-Brometalia) (* | 6210 | 2013 2007 | | U2- U2- | |
| | Siliceous alpine and boreal grasslands | 6150 | 2013 2007 | FV FV | FV FV | |
| | Species-rich Nardus grasslands, on silicious substrates in mountain areas (and submountain areas in | 6230 | 2013 2007 | | U2- U2- | |
| Heath & scrub | Alpine and Boreal heaths | 4060 | 2013 2007 | U1- U1- | U1= U1- c1 | |
| | European dry heaths | 4030 | 2013 2007 | | U2- U2- | |

| Group | Name | Code | Year | ALP | BOR | MBAL |
|---------------------|--|------|--------------|----------|-----------------|------|
| | Sub-Arctic Salix spp. scrub | 4080 | 2013 2007 | FV FV | FV FV | |
| Freshwater habitats | Alpine rivers and the herbaceous vegetation along their banks | 3220 | 2013 2007 | FV FV | FV FV | 1 |
| | Alpine rivers and their ligneous vegetation with Myricaria germanica | 3230 | 2013 2007 | FV FV | | |
| | Fennoscandian natural rivers | 3210 | 2013 2007 | FV FV | U1= U1 nc | |
| | Hard oligo-mesotrophic waters with benthic vegetation of Chara spp. | 3140 | 2013 2007 | | U1+ U1 nc | |
| | Natural dystrophic lakes and ponds | 3160 | 2013 2007 | FV FV | U1+ U1 nc | |
| | Natural eutrophic lakes with Magnopotamion or Hydrocharition — type vegetation | 3150 | 2013 2007 | | U2+ U2 nc | |
| | Oligotrophic to mesotrophic standing waters with vegetation of the Littorelletea uniflorae and/or of the | 3130 | 2013 2007 | | U1+ U1 nc | |
| | Oligotrophic waters containing very few minerals of sandy plains (Littorelletalia uniflorae) | 3110 | 2013 2007 | FV FV | U1+ U1 nc | |
| | Water courses of plain to montane levels with the Ranunculion fluitantis and Callitricho-Batrachion vegetation | 3260 | 2013 2007 | FV FV | U2+ U2+ | |
| Dunes habitats | Decalcified fixed dunes with Empetrum nigrum | 2140 | 2013 2007 | | U1= U1 nc | |
| | Dry sand heaths with Calluna and Empetrum nigrum | 2320 | 2013 2007 | | U1= U1 nc | |
| | Embryonic shifting dunes | 2110 | 2013 2007 | | U1= U1 nc | |
| | Fixed coastal dunes with herbaceous vegetation ("grey dunes') | 2130 | 2013 2007 | | U2= U2 nc | |
| | Humid dune slacks | 2190 | 2013 2007 | | U1- U1 nc | |
| | Shifting dunes along the shoreline with Ammophila arenaria ('white dunes') | 2120 | 2013 2007 | | U1- U1 nc | |
| | Wooded dunes of the Atlantic, Continental and Boreal region | 2180 | 2013 2007 | | U2- U2 nc | |
| Coastal habitats | Annual vegetation of drift lines | 1210 | 2013 2007 | | U1= U1 nc | |
| | Baltic esker islands with sandy, rocky and shingle beach vegetation and sublittoral vegetation | 1610 | 2013 2007 | | U1- U1 nc | |
| | Boreal Baltic coastal meadows | 1630 | 2013 2007 | | U2+ U2 a | |
| | Boreal Baltic islets and small islands | 1620 | 2013 2007 | | FV FV | |
| | Boreal Baltic narrow inlets | 1650 | 2013 2007 | | | U1- |
| | Boreal Baltic sandy beaches with perennial vegetation | 1640 | 2013 2007 | | U2- U2- | |
| | Coastal lagoons | 1150 | 2013 2007 | | U1- U1- | |
| | Estuaries | 1130 | 2013 2007 | | | U2= |

| Group | Name | Code | Year | ALP | BOR | MBAL |
|-------|--|------|--------------|-----|----------|-----------------|
| | Large shallow inlets and bays | 1160 | 2013 2007 | | | U1- U1- |
| | Perennial vegetation of stony banks | 1220 | 2013 2007 | | FV FV | |
| | Reefs | 1170 | 2013 2007 | | | U1- U1- |
| | Sandbanks which are slightly covered by sea water all the time | 1110 | 2013 2007 | | | U1- U1 nc |
| | Vegetated sea cliffs of the Atlantic and Baltic Coasts | 1230 | 2013 2007 | | FV FV | |

Habitat types reported as scientific reserve (SR), marginal (MAR), invalid report in marine region (IRM) etc. (only listed when a scientific reserve etc has been reported)

| Group | Name | Code | Year | ALP |
|--------------------|---|------|--------------|-----|
| Forests | Coniferous forests on, or connected to, glaciofluvial eskers | 9060 | 2013 2007 | MAR |
| | Fennoscandian deciduous swamp woods | 9080 | 2013 2007 | MAR |
| Bogs, mires & fens | Active raised bogs | 7110 | 2013 2007 | MAR |
| Grasslands | Fennoscandian lowland species-rich dry to mesic grasslands | 6270 | 2013 2007 | MAR |
| | Northern boreal alluvial meadows | 6450 | 2013 2007 | MAR |
| Heath & scrub | European dry heaths | 4030 | 2013 2007 | MAR |

Species reported by Finland

| Group | Name | Code | Year | ALP | BOR | MBAL |
|---------------------|-------------------------------------|------|--------------|-----------|-----------|------|
| | | | | | | |
| Non-vascular plants | Buxbaumia viridis | 1386 | 2013 2007 | | U2= U1 | |
| | | | 2007 | | b1 | |
| | Cephalozia macounii | 1980 | 2013 | | U2= | |
| | | | 2007 | | U1 | |
| | Cladonia spp. (subgenus Cladina) | 1378 | 2013 | U1= | a FV | |
| | Cladolila Spp. (Subgerlas Cladilla) | 10/0 | 2007 | U1 | FV | |
| | | | | c1 | | |
| | Cynodontium suecicum | 1981 | 2013 | | U1x | |
| | | | 2007 | | U1 nc | |
| | Dichelyma capillaceum | 1383 | 2013 | | U1= | |
| | | | 2007 | | U1 | |
| | Dianana sinida | 4004 | 0040 | | nc | |
| | Dicranum viride | 1381 | 2013 2007 | | U1x U1 | |
| | | | 2007 | | nc | |
| | Drepanocladus vernicosus | 1393 | 2013 | FV | U1= | |
| | | | 2007 | XX | U1 | |
| | Encalypta mutica | 1982 | 2013 | b1 U1= | nc U1x | |
| | | 1302 | 2013 | U1 | U1 | |
| | 1 | | | nc | nc | |

| Group | Name | Code | Year | ALP | BOR | MBAL |
|-----------------|-------------------------------------|------|--------------|-----------------|------------------|------|
| | Hamatocaulis lapponicus | 1983 | 2013 2007 | | U2= U1 b1 | |
| | Herzogiella turfacea | 1984 | 2013 2007 | | U1- U1 nc | |
| | Leucobryum glaucum | 1400 | 2013 2007 | | FV FV | |
| | Meesia longiseta | 1389 | 2013 2007 | FV | U1= U2 b1 | |
| | Orthothecium lapponicum | 1986 | 2013 2007 | U1= XX b1 | | |
| | Plagiomnium drummondii | 1987 | 2013 2007 | | U1- U1- | |
| | Sphagnum spp. | 1409 | 2013 2007 | FV FV | FV FV | |
| Vascular plants | Agrimonia pilosa | 1939 | 2013 2007 | | U1- U1 a | |
| | Alisma wahlenbergii | 1940 | 2013 2007 | | U1+ U1 b1 | |
| | Arctagrostis latifolia | 1941 | 2013 2007 | | FV FV | |
| | Arctophila fulva | 1942 | 2013 2007 | | U2- U2 a | |
| | Arenaria ciliata ssp. pseudofrigida | 1943 | 2013 2007 | | FV FV | |
| | Artemisia campestris ssp. bottnica | 1945 | 2013 2007 | | U2+ U2 c1 | |
| | Asplenium adulterinum | 4066 | 2013 2007 | | FV FV | |
| | Botrychium simplex | 1419 | 2013 2007 | | U2x U2- b1 | |
| | Calypso bulbosa | 1949 | 2013 2007 | | U1- U1 b1 | |
| | Carex holostoma | 1950 | 2013 2007 | FV FV | FV FV | |
| | Cinna latifolia | 1951 | 2013 2007 | | U1= U1 nc | |
| | Crepis tectorum ssp. nigrescens | 1953 | 2013 2007 | U1+ U2 b1 | U1= FV c1 | |
| | Cypripedium calceolus | 1902 | 2013 2007 | | U1x U1 nc | |
| | Diplazium sibiricum | 1955 | 2013 2007 | FV FV | FV FV | |
| | Draba cinerea | 1957 | 2013 2007 | | FV FV | |
| | Dryopteris fragans | 1958 | 2013 2007 | FV FV | | |
| | Hippuris tetraphylla | 1960 | 2013 2007 | | U1- U2 b1 | |
| | Liparis loeselii | 1903 | 2013 2007 | | U2- U2 a | |

| Group | Name | Code | Year | ALP | BOR | MBAL |
|------------|----------------------------------|------|--------------|-----------------|------------------|------|
| | Lycopodium spp. | 1413 | 2013 2007 | FV FV | FV FV | |
| | Moehringia lateriflora | 1962 | 2013 2007 | | U1x U1 nc | |
| | Najas flexilis | 1833 | 2013 2007 | | U1- U1 nc | |
| | Najas tenuissima | 1963 | 2013 2007 | | U1- U1 nc | |
| | Persicaria foliosa | 1966 | 2013 2007 | | U1- U1 a | |
| | Primula nutans | 1968 | 2013 2007 | | U1= U1- b1 | |
| | Puccinellia phryganodes | 1971 | 2013 2007 | | U2x U2 nc | |
| | Pulsatilla patens | 1477 | 2013 2007 | | U2- U2- | |
| | Ranunculus lapponicus | 1972 | 2013 2007 | FV FV | FV FV | |
| | Saxifraga hirculus | 1528 | 2013 2007 | FV FV | U1x U1 nc | |
| | Silene furcata ssp. angustiflora | 1975 | 2013 2007 | | U2- U1- a | |
| | Sorbus teodori | 1976 | 2013 2007 | | FV | |
| | Trisetum subalpestre | 1977 | 2013 2007 | FV FV | | |
| | Viola rupestris ssp. relicta | 1978 | 2013 2007 | FV FV | | |
| Molluscs | Margaritifera margaritifera | 1029 | 2013 2007 | U2x FV b1 | U2- U1 b1 | |
| | Unio crassus | 1032 | 2013 2007 | | FV FV | |
| | Vertigo angustior | 1014 | 2013 2007 | | FV XX b1 | |
| | Vertigo genesii | 1015 | 2013 2007 | | XX XX | |
| | Vertigo geyeri | 1013 | 2013 2007 | XX b1 | FV XX b1 | |
| Arthropods | Aeshna viridis | 1048 | 2013 2007 | | U1= U1- b1 | |
| | Agathidium pulchellum | 1919 | 2013 2007 | | U1= U1 nc | |
| | Agriades glandon aquilo | 1930 | 2013 2007 | FV FV | | |
| | Aradus angularis | 1929 | 2013 2007 | | U1= U1 nc | |
| | Astacus astacus | 1091 | 2013 2007 | | FV FV | |
| | Boros schneideri | 1920 | 2013 2007 | | U1- U1 nc | |

| Group | Name | Code | Year | ALP | BOR | MBAL |
|-------|---------------------------|------|--------------|----------------|------------------|------|
| | Clossiana improba | 1931 | 2013 2007 | FV FV | | |
| | Corticaria planula | 1921 | 2013 2007 | | XX XX | |
| | Cucujus cinnaberinus | 1086 | 2013 2007 | | U2= U2 nc | |
| | Dytiscus latissimus | 1081 | 2013 2007 | | FV FV | |
| | Erebia medusa polaris | 1932 | 2013 2007 | FV U1 b1 | | |
| | Euphydryas aurinia | 1065 | 2013 2007 | | U1- U1 nc | |
| | Graphoderus bilineatus | 1082 | 2013 2007 | | FV FV | |
| | Hesperia comma catena | 1933 | 2013 2007 | FV FV | | |
| | Hypodryas maturna | 1052 | 2013 2007 | | FV FV | |
| | Leucorrhinia albifrons | 1038 | 2013 2007 | | FV FV | |
| | Leucorrhinia caudalis | 1035 | 2013 2007 | | FV FV | |
| | Leucorrhinia pectoralis | 1042 | 2013 2007 | | FV FV | |
| | Lopinga achine | 1067 | 2013 2007 | | FV U1+ a | |
| | Lycaena dispar | 1060 | 2013 2007 | | FV U1+ a | |
| - | Lycaena helle | 4038 | 2013 2007 | | U2- U2- | |
| - | Macroplea pubipennis | 1922 | 2013 2007 | | U1- U1- | |
| - | Maculinea arion | 1058 | 2013 2007 | | U2= U2- c1 | |
| - | Mesosa myops | 1923 | 2013 2007 | | U1= U1 nc | |
| | Ophiogomphus cecilia | 1037 | 2013 2007 | | FV U1 b1 | |
| | Osmoderma eremita | 1084 | 2013 2007 | | FV FV b2 | |
| | Oxyporus mannerheimii | 1924 | 2013 2007 | | XX U1 c1 | |
| | Parnassius apollo | 1057 | 2013 2007 | | U1= U1 nc | |
| | Parnassius mnemosyne | 1056 | 2013 2007 | | U1= U1 nc | |
| | Phryganophilus ruficollis | 4021 | 2013 2007 | | U1x U1 nc | |
| | Pytho kolwensis | 1925 | 2013 2007 | | U1= U1- c1 | |

| Group | Name | Code | Year | ALP | BOR | MBAL |
|------------|----------------------------|------|--------------|-----------------|------------------|----------|
| | Stephanopachys linearis | 1926 | 2013 2007 | | U1= U1+ c1 | |
| | Stephanopachys substriatus | 1927 | 2013 2007 | | U1= U1+ c1 | |
| | Xestia borealis | 1934 | 2013 2007 | | U1x U1 nc | |
| | Xestia brunneopicta | 1935 | 2013 2007 | | XX XX | |
| | Xyletinus tremulicola | 1928 | 2013 2007 | | U1= U1 nc | |
| Fish | Aspius aspius | 1130 | 2013 2007 | | FV FV | |
| | Cobitis taenia | 1149 | 2013 2007 | | FV XX b1 | |
| | Coregonus albula | 2492 | 2013 2007 | | FV FV | |
| | Coregonus lavaretus | 2494 | 2013 2007 | FV FV | U1= FV a | |
| | Cottus gobio | 1163 | 2013 2007 | | FV nhd FV | |
| | Lampetra fluviatilis | 1099 | 2013 2007 | | FV FV | |
| | Lampetra planeri | 1096 | 2013 2007 | | FV nhd FV | |
| | Salmo salar | 1106 | 2013 2007 | FV FV | FV FV | |
| | Thymallus thymallus | 1109 | 2013 2007 | FV FV | U1= FV a | |
| Amphibians | Rana arvalis | 1214 | 2013 2007 | | FV FV | |
| | Rana temporaria | 1213 | 2013 2007 | FV FV | FV FV | |
| | Triturus cristatus | 1166 | 2013 2007 | | U1x U1 nc | |
| Reptiles | Coronella austriaca | 1283 | 2013 2007 | | U1- XX nc | |
| Mammals | Alopex lagopus | 1911 | 2013 2007 | U2= U2- a | | |
| | Canis lupus | 1352 | 2013 2007 | U1= FV c1 | U1- FV a | |
| | Castor fiber | 1337 | 2013 2007 | | U1= FV c2 | |
| | Eptesicus nilssonii | 1313 | 2013 2007 | XX XX | FV FV | |
| | Gulo gulo | 1912 | 2013 2007 | FV FV | U1+ U1+ | |
| | Halichoerus grypus | 1364 | 2013 2007 | | | FV FV |
| | Lepus timidus | 1334 | 2013 2007 | FV FV | U1- FV a | |

| Group | Name | Code | Year | ALP | BOR | MBAL |
|---------------------|----------------------------|------|--------------|----------|-----------------|-----------------|
| | Lutra lutra | 1355 | 2013 2007 | FV FV | FV FV | |
| | Lynx lynx | 1361 | 2013 2007 | FV FV | FV FV | |
| | Martes martes | 1357 | 2013 2007 | FV FV | FV FV | |
| | Mustela putorius | 1358 | 2013 2007 | | U1- XX nc | |
| | Myotis brandtii | 1320 | 2013 2007 | | XX XX | |
| | Myotis daubentonii | 1314 | 2013 2007 | | FV FV | |
| | Myotis mystacinus | 1330 | 2013 2007 | | XX XX | |
| | Myotis nattereri | 1322 | 2013 2007 | | XX XX | |
| | Phoca hispida botnica | 1938 | 2013 2007 | | | U1x U1+ a |
| | Phoca hispida saimensis | 1913 | 2013 2007 | | U2+ U2+ | |
| | Pipistrellus nathusii | 1317 | 2013 2007 | | XX XX | |
| | Plecotus auritus | 1326 | 2013 2007 | | FV FV | |
| | Pteromys volans | 1910 | 2013 2007 | | U1- U1 nc | |
| | Rangifer tarandus fennicus | 1937 | 2013 2007 | | U1+ FV a | |
| | Sicista betulina | 1343 | 2013 2007 | | FV FV | |
| | Ursus arctos | 1354 | 2013 2007 | FV FV | FV FV | |
| Other invertebrates | Hirudo medicinalis | 1034 | 2013 2007 | | XX XX | |

Species reported as occasional (OCC), newly arriving (ARR), extinct prior the Habitats Directive came into force (PEX), marginal (MAR), invalid report in marine region (IRM) or introduced (INT) etc. (only listed when an occasional species etc has been reported). In addition species with optional reports (OP) and scientific reserves (SR) are listed here.

| Group | Name | Code | Year | ALP | BOR | MBAL |
|------------------------|-------------------------------------|------|--------------|-----------|-----|------|
| Non-vascular plants | Cynodontium suecicum | 1981 | 2013 2007 | MAR XX | | |
| | Hygrohypnum montanum | 1985 | 2013 2007 | | ARR | |
| | Scapania massalongii | 1394 | 2013 2007 | | OCC | |
| Vascular plants | Platanthera obtusata ssp. oligantha | 1967 | 2013 2007 | OCC | | |

| Group | Name | Code | Year | ALP | BOR | MBAL |
|------------|---------------------------|------|--------------|-----|-----------------|------|
| Arthropods | Nymphalis vaualbum | 4039 | 2013 2007 | | OCC | |
| | Sympecma braueri | 1039 | 2013 2007 | | ARR XX a | |
| | Xylomoia strix | 4044 | 2013 2007 | | 220 | |
| Fish | Coregonus albula | 2492 | 2013 2007 | MAR | | |
| | Cottus gobio | 1163 | 2013 2007 | MAR | | |
| | Pelecus cultratus | 2522 | 2013 2007 | | ARR | |
| Mammals | Eptesicus serotinus | 1327 | 2013 2007 | | 000 | |
| | Myotis dasycneme | 1318 | 2013 2007 | | 000 | |
| | Nyctalus noctula | 1312 | 2013 2007 | | OCC XX nc | |
| | Phocoena phocoena | 1351 | 2013 2007 | | | 000 |
| | Pipistrellus pipistrellus | 1309 | 2013 2007 | | 000 | |
| | Pipistrellus pygmaeus | 5009 | 2013 2007 | | 000 | |
| | Vespertilio murinus | 1332 | 2013 2007 | | 000 | |