

## Information on measures and related costs in relation to species considered for inclusion on the Union list: *Lygodium japonicum*

This note has been drafted by IUCN within the framework of the contract No 07.0202/2017/763436/SER/ENV.D2 “Technical and Scientific support in relation to the Implementation of Regulation 1143/2014 on Invasive Alien Species”. The information and views set out in this note do not necessarily reflect the official opinion of the Commission, or IUCN. The Commission does not guarantee the accuracy of the data included in this note. Neither the Commission nor IUCN or any person acting on the Commission’s behalf, including any authors or contributors of the notes themselves, may be held responsible for the use which may be made of the information contained therein. Reproduction is authorised provided the source is acknowledged. This document shall be cited as:

Bohn, K.K. 2018. Information on measures and related costs in relation to species considered for inclusion on the Union list: *Lygodium japonicum*. Technical note prepared by IUCN for the European Commission.

Date of completion: 04/10/2018

Comments which could support improvement of this document are welcome. Please send your comments by e-mail to [ENV-IAS@ec.europa.eu](mailto:ENV-IAS@ec.europa.eu).

<b>Species (scientific name)</b>	<i>Lygodium japonicum</i> (Thunb.) Sw
<b>Species (common name)</b>	Japanese climbing fern
<b>Author(s)</b>	Kimberly K. Bohn, Pennsylvania State University, USA
<b>Date Completed</b>	04/10/2018
<b>Reviewer</b>	Johan van Valkenburg, National Plant Protection Organization, Wageningen, Netherlands.

### Summary

Highlight of measures that provide the most cost-effective options to prevent the introduction, achieve early detection, rapidly eradicate and manage the species, including significant gaps in information or knowledge to identify cost-effective measures.

*Lygodium japonicum* (Japanese climbing fern) is a broadleaved, herbaceous perennial vine and true fern which is native to much of south-eastern Asia, through India and as far south as Papua New Guinea (Uddin et al., 1997, - as cited in CABI, 2017; Maideen et al., 2004; Singh et al., 2012; Lindsay & Middleton, 2013; Chang et al., 2014; Flora of China Editorial Committee, 2014). It was introduced outside of its native range on several continents. North American establishment was first recorded in the early 1900’s in the State of Georgia (Clute, 1903), where it was introduced as an ornamental plant (Clarke, 1936; Correll, 1938). *L. japonicum* has since spread throughout the south-eastern USA. In addition to North America, it has also been recognized as an introduced species in South Africa (Henderson, 2007), Singapore (Chong et al., 2009) and Australia, where it is classified as an environmental weed (Orchard & McCarthy, 1998; Hosking et al. 2011).

Where introduced in the USA, *L. japonicum* occupies a broad range of natural and disturbed habitats. While *L. japonicum* has a strong preference for moist soils it can sometimes occupy xeric sites (van Loan, 2006). It is invasive in diverse habitats throughout the southeastern USA ranging from floodplain forests, swamps, marshes, river and stream banks, to pine flatwoods, hardwood hammocks, and upland woodlands (Wunderlin & Hansen 2003; van Loan, 2006; Miller et al., 2010). It commonly invades disturbed areas, including along roads and particularly ditches and culverts. *L. japonicum* can smother groundcover and shrubs, leading to a reduction in the abundance of native species within a local ecosystem (Bohn et al., 2011; Leichty et al., 2011). It has been documented to reduce native plant richness in both mesic and upland pine ecosystems of the U.S. (Ulrich, 2012). It also can climb trees, creating dense patches that limit photosynthesis of tree leaves or needles. In South Africa, it has been found in moist forest, scrub and road edges, and in Australia it has been documented outside of its cultivated area in wet forests and riparian areas (CABI, 2017). It is expected it would invade similar ecosystems in Europe, in areas with climate suitable to its development.

Across its native range, *L. japonicum* tolerates subtropical to tropical conditions with both wet and dry seasons (Koppen-Geiger climate codes Af, Am, and Aw; Kottek et al., 2006), and prefers warm temperate conditions that are fully humid with warm to hot summers. Predictive models of a risk assessment of *L. japonicum* (EPPO, 2017) currently show most suitable climates to be in Spain ( Canary Islands), Portugal( the Azores), France, Italy and coastlines of the Adriatic and Black Sea (Turkey Georgia and Russia). However, climate change models do suggest that much of central and northern Europe is predicted to become suitable, potentially allowing substantial establishment within the Atlantic, Continental, Black Sea, Mediterranean and Boreal biogeographical Regions (including: Portugal, France, Germany, the Netherlands, Belgium, Austria, Hungary, Czech Republic, Great Britain, Poland, Lithuania, Latvia, Italy, Croatia, southern Sweden and Denmark).

Prevention: The primary pathway for introduction of *L. japonicum* into the European Union is through the ornamental trade industry, either intentionally through direct breeding and selling of plants, or unintentionally as a contaminant in the soil or growing medium of other plants for planting. In fact, in the Netherlands gametophytes (small germinants) of *L. japonicum* have been detected in growing media of bonsai plants imported from China (J. van Valkenburg pers. comm. 2017) on more than one occasion.

Therefore, the primary methods for preventing its intentional introduction would be a ban on keeping, importing, selling, breeding and growing this plant. Although a recent web search suggests some limited trade of this species, it is not very popular and a ban would likely not have a significant economic impact on the ornamental trade industry. Prevention of unintentional spread, through the importation and movement of gametophyte contaminants in soils of other plants for planting would require inspection of imported ornamental plants, particularly from facilities where ornamental *L. japonicum* is grown, or from outdoor nurseries in countries where *L. japonicum* is native (Asia) or invasive (southeastern United States, Australia). Inspections can be costly, though made cost-effective by incorporation into an overall inspection plan for Invasive Alien Species. Awareness campaigns to both the industry and buyers may also be a cost-effective method of prevention. Note also that the EPPO recognizes the introduction of *L. japonicum* via transport on imported equipment/machinery from invaded countries as a potential unintentional pathway into the E.U., though this has not been detected or documented thus far.

Prevention of the secondary spread of *L. japonicum*, if established in natural areas within the E.U., would include inspection and cleaning of clothing, equipment, and machinery used in invaded sites, particularly during periods where *L. japonicum* is reproducing. Spores adhere readily to soil, water, and clothing, and plant fragments have been noted to become lodged in equipment and machinery.

Early Detection: Scouting and monitoring of sentinel sites by natural resource professionals and/or in combination with citizen scientist volunteers will be necessary for early detection. As *L. japonicum* is not currently found in the E.U. outside of a few known botanical and private gardens, initial monitoring could be limited to areas just outside those known locations or near large ornamental/nursery industries from where unintentionally introduced *L. japonicum* may spread. Monitoring in larger natural

areas by resource professionals should focus on ecosystems most likely to be invaded within geographical regions that have climates suitable to its establishment now or with projected climate change (see areas listed above, as defined by projections in EPPO, 2017).

**Rapid Eradication:** Plant protection products (post-emergent herbicides) are one of the most effective and efficient ways to eradicate early stages of an infestation of *L. japonicum*. At early stages, chemicals can be applied as a directed foliar spray using hand pumps or backpack sprayers. Glyphosate applied at low rates of 2% v:v have been shown to be effective in several studies (van Loan 2006; Minogue et al., 2010; Bohn et al., 2011) however it was also noted that metsulfuron methyl treatments at 0.75 g/L were least damaging to surrounding native vegetation, particularly graminoids (Zeller & Leslie 2004). Herbicides must be applied according to manufacturer instructions and in accordance with EU and national regulations (particularly near waterways), and precautions must be applied to prevent non-target damage.

At early stages, particularly in the first year of establishment, mechanically removing *L. japonicum* is also feasible but must include complete removal of the below-ground root and rhizome system. Below-ground biomass will need to be dug out, by hand in loose soil or with a small shovel. Roots and rhizomes should be separated from soil and disposed of properly to prevent further spread, either by burning or desiccating and killing plant material in plastic bags or under tarps. Above-ground plant material with spores present should be disposed of in a similar manner.

**Management:** Herbicide application is the primary method by which *L. japonicum* is managed in widespread and highly invaded areas. Glyphosate at a rate between 2-4% v:v solution is recommended for large scale infestations, though metsulfuron methyl at a rate of 0.038-0.75 g/L applied may at least provide short-term (1-2yr) control. A tank mix of the two chemicals has been found to be effective for controlling both adult plants and spore viability. Mechanical treatments alone are not effective for controlling *L. japonicum*, and in fact may be more detrimental. However, cutting of large climbing vines in combination with herbicide application to the lower portion of the vine as an Integrated Vegetation Management approach has been successful for controlling some *Lygodium* species. An integrated approach of utilizing prescribed fire and herbicides has also been effective in the southeastern United States, and would be appropriate in fire-dependent ecosystems of the E.U. where prescribed fire may be utilized, such as Mediterranean pine forests and shrublands in southern Portugal, France, and Italy (Fernandes et al, 2013) which are also climatically suitable for *L. japonicum*.

**Prevention of intentional introductions and spread** – measures for preventing the species being introduced intentionally. **This table is repeated for each of the prevention measures identified.**

<p><b>Measure description</b> Provide a description of the measure, and identify its objective</p>	<p><b>The measure that can best prevent the <i>intentional</i> introduction of <i>Lygodium japonicum</i> is a ban on keeping, importing, selling, breeding and growing this plant.</b></p> <p>The ornamental trade has been the main pathway of introduction of <i>L. japonicum</i> on other continents where it has become invasive. It was imported for use in home gardens and landscaping in the southern United States in the early 1930's (Clarke, 1936; Correll, 1938), and for planting in botanical gardens in Australia in the early 1900's (Orchard &amp; McCarthy, 1998). A recent internet search suggests it is available for purchase and import into the European Union (EPPO, 2017), and indications from plant exchange website forums (for example <a href="https://davesgarden.com/community/trading/search.php">https://davesgarden.com/community/trading/search.php</a>) suggest the species is currently available and already exists in the European Union but to a low extent. For example, one supplier in Ireland has been highlighted as supplying the species in the past, and another supplier is also listed on the Royal Horticultural Society (GB) website.</p>
--	--

<p><b>Effectiveness of the measure</b> Is it effective in relation to its objective? Has the measure previously worked, failed?</p> <p>Please select one of the categories of effectiveness (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<table border="1" data-bbox="645 153 1888 220"> <tr> <td><b>Effectiveness of measures</b></td> <td><i>Effective</i></td> <td><input checked="" type="checkbox"/></td> <td><i>Neutral</i></td> <td><input type="checkbox"/></td> <td><i>Ineffective</i></td> <td><input type="checkbox"/></td> </tr> </table> <p>Rationale: In the United States, <i>L. japonicum</i> was listed as a noxious weed in two states, Florida and Alabama, in 1999. This measure has been effective at stopping the selling, growing, and importation of new plants into these states; however the listing occurred well after this species had spread aggressively throughout the southeastern U.S. Further, it is not listed as a federal noxious weed, and therefore may be purchased and grown in other nearby states.</p> <p>A trade ban implemented across the European Union could be expected to be an effective measure because this species has not been widely detected or documented in the E.U. and would truly be a preventative measure. It currently only appears to be of limited interest to home gardeners. Preventing its trade and breeding as an ornamental now, before it has spread throughout the region, will be more effective than has been done on other continents. This measure will only be effective, however, if enforced across the entire E.U. region.</p>	<b>Effectiveness of measures</b>	<i>Effective</i>	<input checked="" type="checkbox"/>	<i>Neutral</i>	<input type="checkbox"/>	<i>Ineffective</i>	<input type="checkbox"/>														
<b>Effectiveness of measures</b>	<i>Effective</i>	<input checked="" type="checkbox"/>	<i>Neutral</i>	<input type="checkbox"/>	<i>Ineffective</i>	<input type="checkbox"/>																
<p><b>Side effects (incl. potential) – both positive and negative</b> i.e. positive or negative side effects of the measure on public health, environment including non-targeted species, etc.</p> <p>For each of the side effect types please select one of the impact categories (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<table border="1" data-bbox="645 644 1888 751"> <tr> <td><b>Environmental effects</b></td> <td><i>Positive</i></td> <td><input type="checkbox"/></td> <td><i>Neutral or mixed</i></td> <td><input checked="" type="checkbox"/></td> <td><i>Negative</i></td> <td><input type="checkbox"/></td> </tr> <tr> <td><b>Social effects</b></td> <td><i>Positive</i></td> <td><input type="checkbox"/></td> <td><i>Neutral or mixed</i></td> <td><input checked="" type="checkbox"/></td> <td><i>Negative</i></td> <td><input type="checkbox"/></td> </tr> <tr> <td><b>Economic effects</b></td> <td><i>Positive</i></td> <td><input type="checkbox"/></td> <td><i>Neutral or mixed</i></td> <td><input checked="" type="checkbox"/></td> <td><i>Negative</i></td> <td><input type="checkbox"/></td> </tr> </table> <p>Rationale: Environmental: No side effects are anticipated, though preventing the intentional introduction of <i>L japonicum</i> into the E.U. would be a key measure to limit the potential for it to establish in and impact natural areas.</p> <p>Social: <i>L. japonicum</i> is recorded as having medicinal value in its native range. For example, it is used as a diuretic in China and for treatment of colds, inflammation, and renal ailments; and in India to treat snakebites, diabetes, and ulcers (Puri, 1970; Eisenberg et al., 2009; Yumkham &amp; Singh, 2011- as cited in CABI 2017). However, no conclusive evidence of its health benefits has been documented except for it having several antioxidants (Duan, 2012). In the E.U., it appears to be purchased primarily for ornamental use, and a ban on trade, breeding, and growing, would only have limited or negligible social impact.</p> <p>Economic: Costs associated with enforcing a ban on trade, breeding, or growing would likely offset any cost associated with treatment of this species were it to escape and establish throughout the E.U. (see Eradication and Management sections for cost of <i>L. japonicum</i> treatment). Because <i>L. japonicum</i> is not widely sold or traded as an ornamental plant, a ban on this species would have little economic side effects on the ornamental/horticulture industries.</p>	<b>Environmental effects</b>	<i>Positive</i>	<input type="checkbox"/>	<i>Neutral or mixed</i>	<input checked="" type="checkbox"/>	<i>Negative</i>	<input type="checkbox"/>	<b>Social effects</b>	<i>Positive</i>	<input type="checkbox"/>	<i>Neutral or mixed</i>	<input checked="" type="checkbox"/>	<i>Negative</i>	<input type="checkbox"/>	<b>Economic effects</b>	<i>Positive</i>	<input type="checkbox"/>	<i>Neutral or mixed</i>	<input checked="" type="checkbox"/>	<i>Negative</i>	<input type="checkbox"/>
<b>Environmental effects</b>	<i>Positive</i>	<input type="checkbox"/>	<i>Neutral or mixed</i>	<input checked="" type="checkbox"/>	<i>Negative</i>	<input type="checkbox"/>																
<b>Social effects</b>	<i>Positive</i>	<input type="checkbox"/>	<i>Neutral or mixed</i>	<input checked="" type="checkbox"/>	<i>Negative</i>	<input type="checkbox"/>																
<b>Economic effects</b>	<i>Positive</i>	<input type="checkbox"/>	<i>Neutral or mixed</i>	<input checked="" type="checkbox"/>	<i>Negative</i>	<input type="checkbox"/>																
<p><b>Acceptability to stakeholders</b> e.g. impacted economic activities, animal welfare considerations, public perception, etc.</p>	<table border="1" data-bbox="645 1276 1888 1343"> <tr> <td><b>Acceptability to stakeholders</b></td> <td><i>Acceptable</i></td> <td><input checked="" type="checkbox"/></td> <td><i>Neutral or mixed</i></td> <td><input type="checkbox"/></td> <td><i>Unacceptable</i></td> <td><input type="checkbox"/></td> </tr> </table> <p>Rationale:</p>	<b>Acceptability to stakeholders</b>	<i>Acceptable</i>	<input checked="" type="checkbox"/>	<i>Neutral or mixed</i>	<input type="checkbox"/>	<i>Unacceptable</i>	<input type="checkbox"/>														
<b>Acceptability to stakeholders</b>	<i>Acceptable</i>	<input checked="" type="checkbox"/>	<i>Neutral or mixed</i>	<input type="checkbox"/>	<i>Unacceptable</i>	<input type="checkbox"/>																

<p>Please select one of the categories of acceptability (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<p>The social and economic impact on the ornamental plant industry is expected to be low since a recent review of plant forums and websites suggest is has only been sought after and sold in small quantities. Additionally, nursery professionals have become more cognizant of selling plants that have been identified as invasive, and increasingly are willing to forego importation, breeding, and selling of those plants, (Burt et al., 2007), particularly when those plants are not a major economic commodity such as is the case with <i>Lygodium japonicum</i>.</p>								
<p><b>Additional cost information</b><sup>1</sup> When not already included above, or in the species Risk Assessment.</p> <ul style="list-style-type: none"> <li>- the cost of inaction</li> <li>- the cost-effectiveness</li> <li>- the socio-economic aspects</li> </ul> <p>Include quantitative &amp;/or qualitative data, and case studies (incl. from countries outside the EU).</p>	<p>As stated above, costs associated with enforcing a ban on trade, breeding, or growing would likely more than offset any cost associated with treatment of this species were it to escape and establish throughout the E.U. Exact costs for enforcement of a ban in the E.U. are unknown. Providing awareness of this species to the ornamental trade industry and emphasizing internal voluntary compliance could increase cost effectiveness in terms of insuring or enforcing compliance with the ban through monitoring and inspections.</p>								
<p><b>Level of confidence on the information provided</b><sup>2</sup></p> <p>Please select one of the confidence categories along with a statement to support the category chosen. See <i>Notes</i> section at the bottom of this document.</p> <p><b>NOTE – this is not related to the effectiveness of the measure</b></p>	<i>Inconclusive</i>		<i>Unresolved</i>		<i>Established but incomplete</i>		<i>Well established</i>	x	
		<p>Rationale: It is expected that regulations to ban importing, selling, breeding and growing this plant would be effective as a means to prevent additional introductions into the E.U.</p>							

<p><b>Prevention of intentional introductions and spread</b> – measures for preventing the species being introduced intentionally. <b>This table is repeated for each of the prevention measures identified.</b></p>	
<p><b>Measure description</b> Provide a description of the measure, and identify its objective</p>	<p><b>Awareness campaigns targeted to key stakeholder groups to reduce the risk of intentional introductions, and also unintentional introductions as a contaminant in other plants for planting (see table below).</b></p> <p>Awareness campaigns targeted at both ornamental industries (growers and sellers) and to the general public or gardeners in particular (buyers), about the invasive potential of <i>L. japonicum</i>, is another measure that could help prevent new intentional and</p>

	<p>unintentional introductions and spread. Advising or providing training to industry professionals regarding best management practices to restrict trade of <i>L. japonicum</i> and to dispose of plant growing medium unintentionally contaminated with <i>L. japonicum</i> could be incorporated into this measure. Education and awareness campaigns to the general public could include fliers, announcements, or other educational materials specifically focusing on <i>L. japonicum</i> or incorporated into existing materials educating the public about invasive plants in general.</p>													
<p><b>Effectiveness of the measure</b> Is it effective in relation to its objective? Has the measure previously worked, failed?</p> <p>Please select one of the categories of effectiveness (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<p><b>Effectiveness of measures</b></p>	<p>Effective</p>	<p>x</p>	<p>Neutral</p>		<p>Ineffective</p>								
<p><i>Rationale:</i> Awareness of biosecurity issues has been identified as one of the top 20 priorities for invasive species awareness in the European Union (Caffrey et al., 2014). Few studies have documented the impact and effectiveness of awareness campaigns at limiting intentional introductions of invasive species; though one recent study of the aquatics trade (Verbrugge et al., 2014) suggests that retailers are more likely to be aware and act on the need to control potential impacts of invasive species than buyers. This suggests that additional education and outreach may be needed for the latter stakeholders.</p> <p>It is more difficult to assess the effectiveness of awareness campaigns regarding limiting unintentional introductions through contaminants. Even with awareness and education, effectiveness would more likely be related to the frequency of inspections and ability to detect <i>L. japonicum</i> within other plants or in plant growing mediums.</p>														
<p><b>Side effects (incl. potential) – both positive and negative</b> i.e. positive or negative side effects of the measure on public health, environment including non-targeted species, etc.</p> <p>For each of the side effect types please select one of the impact categories (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<p><b>Environmental effects</b></p>	<p>Positive</p>	<p>X</p>	<p>Neutral or mixed</p>		<p>Negative</p>								
<p><b>Social effects</b></p>									<p>Positive</p>		<p>Neutral or mixed</p>	<p>x</p>	<p>Negative</p>	
<p><b>Economic effects</b></p>									<p>Positive</p>		<p>Neutral or mixed</p>	<p>x</p>	<p>Negative</p>	
<p>One potential positive side effect would be that those targeted by any awareness raising activities, may become more aware of invasive alien species in general, and therefore the measure may potentially reduce the risk of introductions of other species.</p>														
<p><b>Acceptability to stakeholders</b> e.g. impacted economic activities, animal welfare considerations, public perception, etc.</p>	<p><b>Acceptability to stakeholders</b></p>	<p>Acceptable</p>	<p>x</p>	<p>Neutral or mixed</p>		<p>Unacceptable</p>								
<p>Both the trade industry and general public are becoming more aware of the impact of invasive alien plants and are generally willing to learn more about non-native plants. As noted in the Effectiveness section above, consistent outreach and education through awareness campaigns may be needed to change behaviour.</p>														

Please select one of the categories of acceptability (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.									
<b>Additional cost information <sup>1</sup></b> When not already included above, or in the species Risk Assessment.  - the cost of inaction - the cost-effectiveness - the socio-economic aspects  Include quantitative &/or qualitative data, and case studies (incl. from countries outside the EU).	No additional cost information available.								
<b>Level of confidence on the information provided <sup>2</sup></b>  Please select one of the confidence categories along with a statement to support the category chosen. See <i>Notes</i> section at the bottom of this document. <b>NOTE – this is not related to the effectiveness of the measure</b>	<i>Inconclusive</i>		<i>Unresolved</i>		<i>Established but incomplete</i>	x	<i>Well established</i>		While awareness campaigns are generally received positively, few studies have evaluated the degree to which such campaigns lead to the reduction in trade or unintended introductions in comparison to other efforts such as regulations or inspections.

<b>Prevention of <u>un-intentional</u> introductions and spread</b> – measures for preventing the species being introduced un-intentionally (cf. Article 13 of the IAS Regulation). <b>This table is repeated for each of the prevention measures identified.</b>	
<b>Measure description</b> Provide a description of the measure, and identify its objective	<b>A measure to prevent the <u>un-intentional</u> introduction and spread of <i>L. japonicum</i> is through the inspection of the growing medium of other species imported as plants for planting.</b>  <i>L. japonicum</i> spores are small, ranging in size from 64-80 µm, and readily disperse by wind. In nurseries where <i>L. japonicum</i> is grown alongside other plants, or where plants are grown outdoors in the vicinity of naturally occurring <i>L. japonicum</i> , spores have been known to disperse and germinate in the growing medium of other plants. Spores may also remain viable for several years.

	<p>In the Netherlands, gametophytes (i.e. young germinants) have been detected in growing media of bonsai plants imported from China (J. van Valkenburg pers. comm. 2017) on more than one occasion. Contaminated soil of other plants for planting may therefore act as a pathway for its unintentional introduction and spread into natural areas within the E.U. In the United States, <i>L. japonicum</i> spores, as well as reproducing vegetative rhizomes, have also been known to be unintentionally spread in contaminated mulch products. While spores themselves may be undetectable, inspection of growing medium or mulch of other plants for <i>L. japonicum</i> gametophytes or sporophytes might be an effective method for limiting unintentional introductions and spread.</p>																					
<p><b>Scale of application</b> At what scale is the measure applied? What is the largest scale at which it has been successfully used? Please provide examples, with areas (km<sup>2</sup> or ha) if possible.</p>	<p>To be effective, inspection of plants for planting for possible <i>L. japonicum</i> contaminants would be required across the entire E.U. region.</p>																					
<p><b>Effectiveness of the measure</b> Is it effective in relation to its objective? Has the measure previously worked, failed?  Please select one of the categories of effectiveness (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<table border="1" data-bbox="645 651 1888 715"> <tr> <td><i>Effectiveness of measures</i></td> <td><i>Effective</i></td> <td></td> <td><i>Neutral</i></td> <td>X</td> <td><i>Ineffective</i></td> <td></td> </tr> </table> <p>Rationale: The Florida Department of Agriculture and Consumer Affairs has effectively restricted and regulated the movement of <i>L. japonicum</i> plant parts from landscaping materials such as mulch that is transported into and throughout the state. The greatest efficacy has been in detecting root fragments or live fronds interspersed with other landscaping material. The ability to inspect all plants entering the E.U. and to detect small germinants of <i>L. japonicum</i> is likely challenging because of the small size of the germinants, and it is very likely spores will not have germinated during the shipment period and initial entry into the E.U. To be effective this measure would require thorough and continual inspection over at least 6-8 weeks (Ulrich, 2012).</p>	<i>Effectiveness of measures</i>	<i>Effective</i>		<i>Neutral</i>	X	<i>Ineffective</i>															
<i>Effectiveness of measures</i>	<i>Effective</i>		<i>Neutral</i>	X	<i>Ineffective</i>																	
<p><b>Effort required</b> e.g. period of time over which measure need to be applied to have results</p>	<p>Inspection of all plants for planting must be done indefinitely across the entire E.U. region for this measure to be effective.</p>																					
<p><b>Resources required</b><sup>1</sup> e.g. cost, staff, equipment etc.</p>	<p>While the cost of initiating new inspections can often be high, where inspection of nursery stock is already occurring for other invasive plant propagules, little additional cost may be needed to also inspect for <i>L. japonicum</i>. Staff will need training on the identification of <i>L. japonicum</i> gametophytes and sporophytes.</p>																					
<p><b>Side effects (incl. potential) – both positive and negative</b> i.e. positive or negative side effects of the measure on public health,</p>	<table border="1" data-bbox="645 1284 1888 1385"> <tr> <td><i>Environmental effects</i></td> <td><i>Positive</i></td> <td></td> <td><i>Neutral or mixed</i></td> <td>X</td> <td><i>Negative</i></td> <td></td> </tr> <tr> <td><i>Social effects</i></td> <td><i>Positive</i></td> <td></td> <td><i>Neutral or mixed</i></td> <td>x</td> <td><i>Negative</i></td> <td></td> </tr> <tr> <td><i>Economic effects</i></td> <td><i>Positive</i></td> <td></td> <td><i>Neutral or mixed</i></td> <td>x</td> <td><i>Negative</i></td> <td></td> </tr> </table>	<i>Environmental effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>	X	<i>Negative</i>		<i>Social effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>	x	<i>Negative</i>		<i>Economic effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>	x	<i>Negative</i>	
<i>Environmental effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>	X	<i>Negative</i>																	
<i>Social effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>	x	<i>Negative</i>																	
<i>Economic effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>	x	<i>Negative</i>																	



<p>environment including non-targeted species, etc.</p> <p>For each of the side effect types please select one of the impact categories (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<p>Rationale: See information described in section above regarding Preventing Intentional Spread for similar considerations. Additionally, because of the relatively few cases in which <i>L. japonicum</i> has previously been identified as a contaminant in growing medium of other plants, the measure may have little economic impact to the ornamental and horticultural trades, in terms of disposing of contaminated product, unless regular inspections significantly increases the number of positive detections.</p>															
<p><b>Acceptability to stakeholders</b> e.g. impacted economic activities, animal welfare considerations, public perception, etc.</p> <p>Please select one of the categories of acceptability (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<table border="1" data-bbox="645 424 1888 488"> <tr> <td data-bbox="645 424 925 488"><b>Acceptability to stakeholders</b></td> <td data-bbox="925 424 1151 488"><i>Acceptable</i></td> <td data-bbox="1151 424 1211 488" style="text-align: center;">x</td> <td data-bbox="1211 424 1509 488"><i>Neutral or mixed</i></td> <td data-bbox="1509 424 1570 488"></td> <td data-bbox="1570 424 1832 488"><i>Unacceptable</i></td> <td data-bbox="1832 424 1888 488"></td> </tr> </table> <p>Rationale: Nursery inspections have become more common practice with the increase in international trade and the known problems of nursery stock contamination with pests, pathogens, and invasive plants (Liebhold et al., 2012). Therefore, it is unlikely that inspections for an additional potentially invasive plant will be viewed negatively.</p>								<b>Acceptability to stakeholders</b>	<i>Acceptable</i>	x	<i>Neutral or mixed</i>		<i>Unacceptable</i>		
<b>Acceptability to stakeholders</b>	<i>Acceptable</i>	x	<i>Neutral or mixed</i>		<i>Unacceptable</i>											
<p><b>Additional cost information</b><sup>1</sup> When not already included above, or in the species Risk Assessment.</p> <ul style="list-style-type: none"> <li>- implementation cost for Member States</li> <li>- the cost of inaction</li> <li>- the cost-effectiveness</li> <li>- the socio-economic aspects</li> </ul> <p>Include quantitative &amp;/or qualitative data, and case studies (incl. from countries outside the EU).</p>	<p>The cost effectiveness should be high assuming inspections for Invasive Alien Species in general are already occurring. Screening and plant-quarantines of incoming plant material that contain non-native plant contaminants has been estimated to save \$1.67 billion over 50 years in Australia (Keller et al., 2007), though it is unclear what the actual cost benefits would be in the E.U. for this particular species. The payoff could be high if potential unintentional introductions are prevented; however, it is difficult to know or quantify how many cases where unintentional introductions of <i>L. japonicum</i>, particularly as gametophytes, could be detected as a contaminant in growing medium on other plants imported into the E.U.</p>															
<p><b>Level of confidence on the information provided</b><sup>2</sup></p> <p>Please select one of the confidence categories along with a statement to support the category chosen. See <i>Notes</i> section at the bottom of this document.</p>	<table border="1" data-bbox="645 1121 1924 1185"> <tr> <td data-bbox="645 1121 913 1185"><i>Inconclusive</i></td> <td data-bbox="913 1121 963 1185"></td> <td data-bbox="963 1121 1171 1185"><i>Unresolved</i></td> <td data-bbox="1171 1121 1227 1185"></td> <td data-bbox="1227 1121 1509 1185"><i>Established but incomplete</i></td> <td data-bbox="1509 1121 1570 1185" style="text-align: center;">X</td> <td data-bbox="1570 1121 1868 1185"><i>Well established</i></td> <td data-bbox="1868 1121 1924 1185"></td> </tr> </table> <p>Rationale: Inspection of plants for planting has been successful for detecting other known pests, but there has been no documentation of its efficacy for <i>L. japonicum</i>.</p>								<i>Inconclusive</i>		<i>Unresolved</i>		<i>Established but incomplete</i>	X	<i>Well established</i>	
<i>Inconclusive</i>		<i>Unresolved</i>		<i>Established but incomplete</i>	X	<i>Well established</i>										

<b>NOTE – this is not related to the effectiveness of the measure</b>	
---	--

**Prevention of secondary spread of the species** – measures for preventing the species spreading once they have been introduced (cf. Article 13 of the IAS Regulation). **This table is repeated for each of the prevention measures identified.**

<b>Measure description</b> Provide a description of the measure, and identify its objective	<p><b>Inspection and cleaning of machinery, equipment, and clothing in areas where <i>L. japonicum</i> is documented as established and reproducing. As the species is not yet established within the EU, this measure is equally applicable for addressing un-intentional introductions.</b></p> <p><i>L. japonicum</i> plant parts, including fertile fronds or roots/rhizomes, have been detected on both clothing and equipment of workers treating invaded areas (Hutchinson &amp; Langeland, 2006). Inspection and cleaning of machinery, equipment, and clothing may therefore limit secondary spread by humans from invaded to non-invaded areas. This measure could apply to both professional settings (e.g. where agriculture or forestry is practiced) as well as in recreational areas (e.g. hiking boots, bike or ATV tires). Cleaning of equipment and machinery is typically done by pressure washing to ensure that soil and plant parts are removed from any part of the equipment. This can be done on-site or at designated cleaning stations. Another common practice for both professionals and in recreation areas is the use of boot brushes to remove dirt and plant parts on-site.</p>					
<b>Scale of application</b> At what scale is the measure applied? What is the largest scale at which it has been successfully used? Please provide examples, with areas (km <sup>2</sup> or ha) if possible.	<p>Guidelines for Best Management Practices (BMPs) and/or specific protocols for inspecting and cleaning equipment and clothing for invasive species typically occurs within an agency/industry, or at regional (state-level) and national scales. The actual implementation of these BMPs would have the most environmental impact at local and possibly regional scales, depending on the distance at which equipment and machinery is typically moved.</p>					
<b>Effectiveness of the measure</b> Is it effective in relation to its objective? Has the measure previously worked, failed?  Please select one of the categories of effectiveness (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.	<i>Effectiveness of measures</i>	<i>Effective</i>	x	<i>Neutral</i>	<i>Ineffective</i>	
	<p>Rationale: Best management practices for various industries, including agriculture and forestry, are more commonly recommending sanitizing equipment and clothing for limiting the spread of many invasive plants, including <i>L. japonicum</i> (Ledoux &amp; Martin, 2012). However, there has been very little research to evaluate or quantify the effectiveness and efficiency of cleaning equipment as a means of preventing the secondary spread of invasive plants by human. As <i>L. japonicum</i> spores can be transported naturally by wind and water over several kilometres (Ferriter, 2001) this measure would be most effective as a means of preventing spread to</p>					

	<p>non-contiguous uninvaded areas at a greater distance. Additionally, to be an effective measure in areas where <i>L. japonicum</i> is reproducing, any run-off from cleaned equipment/machinery must be contained on-site as spores will readily disperse in water.</p>																					
<p><b>Effort required</b> e.g. period of time over which measure need to be applied to have results</p>	<p>Because <i>L. japonicum</i> is absent from natural areas in the E.U., these efforts are not an immediate need for <i>L. japonicum</i> but should be suggested in context to preventing the secondary spread of a large number of existing invasive plants in the region. Additionally, equipment and machinery that may be imported from countries/continents where <i>L. japonicum</i> is native (i.e. Asia) or invasive (i.e. southeastern United States, Australia, or South Africa) should be considered for inspection and/or sanitation before used in natural areas throughout the E.U., particularly in climates suitable to <i>L. japonicum</i> establishment (addressing unintentional introductions). It is only recently that an ISPM Standard, no. 41 (IPPC, 2017) has been drafted and adopted on 'International movement of used vehicles, machinery and equipment'. This focuses on reducing the risks of transporting contaminants (soil, seeds, plant debris, pests) associated with the international movement (either traded or for operational relocation) of vehicles, machinery and equipment (VME) that may have been used in agriculture, forestry, as well as for construction, industrial purposes, mining and waste management, and military.</p> <p>In areas where <i>L. japonicum</i> has established, these measures must be applied indefinitely or up to 5 years after the plant has been removed from its introduced location since spores may remain viable under certain conditions for that length of time.</p>																					
<p><b>Resources required</b><sup>1</sup> e.g. cost, staff, equipment etc.</p>	<p>Pressure washing equipment in a quarantined area and staff to conduct inspections would account for the greatest costs in implementing this measure. Boot brush stations and bike cleaning stations are more regularly being installed in many parks and recreation areas.</p>																					
<p><b>Side effects (incl. potential) – both positive and negative</b> i.e. positive or negative side effects of the measure on public health, environment including non-targeted species, etc.</p> <p>For each of the side effect types please select one of the impact categories (with an 'X'), and provide a</p>	<table border="1" data-bbox="645 1046 1888 1147"> <tr> <td><i>Environmental effects</i></td> <td><i>Positive</i></td> <td>x</td> <td><i>Neutral or mixed</i></td> <td></td> <td><i>Negative</i></td> <td></td> </tr> <tr> <td><i>Social effects</i></td> <td><i>Positive</i></td> <td></td> <td><i>Neutral or mixed</i></td> <td>x</td> <td><i>Negative</i></td> <td></td> </tr> <tr> <td><i>Economic effects</i></td> <td><i>Positive</i></td> <td></td> <td><i>Neutral or mixed</i></td> <td>x</td> <td><i>Negative</i></td> <td></td> </tr> </table> <p>Rationale: The measure will address other potential invasive alien species that could also be introduced via the same pathway. There are likely no substantial negative side-effects of this measure, aside from the additional costs identified in the Resources section. Additionally, as stated in the Effectiveness section, care must be given to collect and dispose of any potentially contaminated run-off from cleaned equipment or clothing to prevent spread into nearby locations.</p>	<i>Environmental effects</i>	<i>Positive</i>	x	<i>Neutral or mixed</i>		<i>Negative</i>		<i>Social effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>	x	<i>Negative</i>		<i>Economic effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>	x	<i>Negative</i>	
<i>Environmental effects</i>	<i>Positive</i>	x	<i>Neutral or mixed</i>		<i>Negative</i>																	
<i>Social effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>	x	<i>Negative</i>																	
<i>Economic effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>	x	<i>Negative</i>																	

<p>rationale, with supporting evidence and examples if possible.</p>									
<p><b>Acceptability to stakeholders</b> e.g. impacted economic activities, animal welfare considerations, public perception, etc.</p> <p>Please select one of the categories of acceptability (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<p><b>Acceptability to stakeholders</b></p>	<p><i>Acceptable</i></p>	<p><i>Neutral or mixed</i></p>	<p>x</p>	<p><i>Unacceptable</i></p>				
<p><b>Additional cost information</b><sup>1</sup> When not already included above, or in the species Risk Assessment. - implementation cost for Member States - the cost of inaction - the cost-effectiveness - the socio-economic aspects</p> <p>Include quantitative &amp;/or qualitative data, and case studies (incl. from countries outside the EU).</p>		<p>No additional cost information available.</p>							
<p><b>Level of confidence on the information provided</b><sup>2</sup></p> <p>Please select one of the confidence categories along with a statement to support the category chosen. See <i>Notes</i> section at the bottom of this document. <b>NOTE – this is not related to the effectiveness of the measure</b></p>	<p><i>Inconclusive</i></p>	<p><i>Unresolved</i></p>	<p>x</p>	<p><i>Established but incomplete</i></p>	<p><i>Well established</i></p>				
		<p>Rationale: Best management practices for invasive plants such as cleaning clothing and equipment are just gaining wider usage, but few studies have quantified their impact on reducing spread of invasive plants from introduced locations to new, uninfested areas and at what distances this measure is most effective.</p>							

**Surveillance measures to support early detection** - Measures to run an effective surveillance system for achieving an early detection of a new occurrence (cf. Article 16). This section assumes that the species is not currently present in a Member State, or part of a Member State's territory. **This table is repeated for each of the early detection measures identified.**

<p><b>Measure description</b> Provide a description of the measure, and identify its objective</p>	<p><b>Early detection monitoring of sentinel (high risk) sites by natural resource professionals.</b></p> <p>Early detection is often a key measure in preventing the ecological and economic harm to ecosystems associated with larger infestations. Monitoring can be incorporated into day-to-day activities of natural resource professionals. Because <i>L. japonicum</i> is not established in natural areas in the E.U. yet, the most efficient implementation of this measure would be to limit monitoring to high risk ecosystems in areas with climates suitable to its establishment. In its invaded range, <i>L. japonicum</i> tends to be most problematic in wetter ecosystems such as floodplain forests, swamps, marshes, river and stream banks, but it can invade upland pine and hardwood forests as well (Wunderlin &amp; Hansen 2003; van Loan, 2006; Miller et al., 2010). It commonly first invades disturbed areas, including along roads and particularly ditches and culverts.</p> <p>Additionally, across its native range, <i>L. japonicum</i> tolerates subtropical to tropical conditions with both wet and dry seasons (Koppen-Geiger climate codes Af, Am, and Aw, Kottek et al., 2006) and prefers warm temperate conditions that are fully humid with warm to hot summers. Predictive models of a risk assessment of <i>L. japonicum</i> (EPPO, 2017) currently show the most suitable climates across the European continent to be in Spain (the Canary Islands), Portugal (the Azores), France, Italy and coastlines of the Adriatic and Black Sea. However, climate change models do suggest that much of central and northern Europe is predicted to become suitable, potentially allowing substantial establishment within the Atlantic, Continental, Black Sea, Mediterranean and Boreal biogeographical Regions (including: Portugal, France, Germany, Netherlands, Belgium, Austria, Hungary, Czech Republic, Great Britain, Poland, Lithuania, Latvia, Italy, Croatia, southern Sweden and Denmark).</p>								
<p><b>Scale of application</b> At what scale is the measure applied? What is the largest scale at which it has been successfully used? Please provide examples, with areas (km<sup>2</sup> or ha) if possible.</p>	<p>Early detection programs have been applied at a variety of scales from the local to regional, the latter particularly so when implemented as a collaboration between multiple organizations and incorporating citizen science (see next section). For example, in the United States, "Cooperative Invasive Species Management Areas" can range in size from several hundred to tens of thousands of square kilometres, and often utilize web mapping tools to record observations across a large area. Individual organizations can also work at the scale of their individual land ownerships.</p>								
<p><b>Effectiveness of the measure</b> Is it effective in relation to its objective? Has the measure previously worked, failed?  Please select one of the categories of effectiveness (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<p><i>Effectiveness of measures</i></p>	<p><i>Effective</i></p>	<p>X</p>	<p><i>Neutral</i></p>		<p><i>Ineffective</i></p>			
<p>Rationale: Early detection has the potential to be highly effective. Though not evaluated for <i>Lygodium</i>, some case studies report that early detection of just small patches of invasive species and rapid response led to eradication in just 2 years, in contrast to a several year delay elsewhere that resulted in spread to thousands of hectares (Simberloff et al., 2013). Effectiveness will depend on consistency of monitoring and ability to positively identify <i>L. japonicum</i> as it first establishes. Early detection is critical as this species can increase by 20-30% cover in one year, and fronds can have reproductive spore production rates of 5,900 spores per square cm within the first year (van Loan, 2006).</p>									

<p><b>Effort required</b> e.g. period of time over which measure need to be applied to have results</p>	<p>While it would be challenging and time consuming to identify gametophytes (new germinants) in a natural ecosystem, adult foliage of <i>L. japonicum</i> should be relatively easy to identify since no other fern species grow as a vine plant form. Additionally there are no other native <i>Lygodium</i> species in Europe with which it could be misidentified. Because aboveground fronds of <i>L. japonicum</i> dies back after multiple frosts, monitoring can be limited to once a year during the early part of the active growing season but preferably before spore production in order to initiate rapid eradication treatments. In the U.S., and in sub-temperate climates, the best period for monitoring would be between late May and early July.</p> <p>As stated previously, because <i>L. japonicum</i> is not currently found in the E.U outside of a few known botanical and private gardens, monitoring in natural areas can be restricted to key ecosystems in geographical regions with climates suitable to its establishment now or with projected climate change (see areas listed above, as defined by projections in <a href="#">EPPO, 2017</a>).</p>																											
<p><b>Resources required</b><sup>1</sup> e.g. cost, staff, equipment etc.</p>	<p>No additional staff or equipment would be needed to carry out early detection monitoring, although it would require additional hours of labour. Monitoring for this species may be incorporated with monitoring of other invasive species or integrated into other environmental data collection that is done routinely.</p>																											
<p><b>Side effects (incl. potential) – both positive and negative</b> i.e. positive or negative side effects of the measure on public health, environment including non-targeted species, etc.</p> <p>For each of the side effect types please select one of the impact categories (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<table border="1" data-bbox="645 683 1888 786"> <tr> <td><i>Environmental effects</i></td> <td><i>Positive</i></td> <td></td> <td><i>Neutral or mixed</i></td> <td>X</td> <td><i>Negative</i></td> <td></td> </tr> <tr> <td><i>Social effects</i></td> <td><i>Positive</i></td> <td></td> <td><i>Neutral or mixed</i></td> <td>x</td> <td><i>Negative</i></td> <td></td> </tr> <tr> <td><i>Economic effects</i></td> <td><i>Positive</i></td> <td></td> <td><i>Neutral or mixed</i></td> <td>x</td> <td><i>Negative</i></td> <td></td> </tr> </table> <p>Rationale: No side effects are expected for surveillance measures for early detection. Though a positive environmental <i>direct</i> effect will ensue if early detection AND rapid eradication occurs.</p>							<i>Environmental effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>	X	<i>Negative</i>		<i>Social effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>	x	<i>Negative</i>		<i>Economic effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>	x	<i>Negative</i>	
<i>Environmental effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>	X	<i>Negative</i>																							
<i>Social effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>	x	<i>Negative</i>																							
<i>Economic effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>	x	<i>Negative</i>																							
<p><b>Acceptability to stakeholders</b> e.g. impacted economic activities, animal welfare considerations, public perception, etc.</p> <p>Please select one of the categories of acceptability (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<table border="1" data-bbox="645 1086 1888 1155"> <tr> <td><i>Acceptability to stakeholders</i></td> <td><i>Acceptable</i></td> <td>x</td> <td><i>Neutral or mixed</i></td> <td></td> <td><i>Unacceptable</i></td> <td></td> </tr> </table> <p>Rationale: Knowledge of invasive plant impacts have increased greatly in the last few decades, and most natural resource managers may already be monitoring for other species. The biggest limitation is manpower and time.</p>							<i>Acceptability to stakeholders</i>	<i>Acceptable</i>	x	<i>Neutral or mixed</i>		<i>Unacceptable</i>															
<i>Acceptability to stakeholders</i>	<i>Acceptable</i>	x	<i>Neutral or mixed</i>		<i>Unacceptable</i>																							

<p><b>Additional cost information</b> <sup>1</sup></p> <p>When not already included above, or in the species Risk Assessment.</p> <ul style="list-style-type: none"> <li>- implementation cost for Member States</li> <li>- the cost of inaction</li> <li>- the cost-effectiveness</li> <li>- the socio-economic aspects</li> </ul> <p>Include quantitative &amp;/or qualitative data, and case studies (incl. from countries outside the EU).</p>	<p>Costs associated with scouting for a species can vary widely based on region, terrain, habitat, vegetation type, and the number of species being scouted simultaneously. Often early detection surveillance by natural resource professionals is incorporated with other day-to-day activities, which can reduce the extra time and labor needed to accomplish that task. The cost of inaction may be high since the likelihood of successful eradication decreases the longer a new invasive species goes undetected. The costs of eradication also increase with time, often exponentially (Simberloff, 2003).</p>								
<p><b>Level of confidence on the information provided</b> <sup>2</sup></p> <p>Please select one of the confidence categories along with a statement to support the category chosen. See <i>Notes</i> section at the bottom of this document.</p> <p><b>NOTE – this is not related to the effectiveness of the measure</b></p>	<i>Inconclusive</i>		<i>Unresolved</i>		<i>Established but incomplete</i>	x	<i>Well established</i>		<p>Rationale:</p> <p>Early detection/rapid response programs have been evaluated for other species, and it is assumed would be effective for <i>L. japonicum</i> as well.</p>

<p><b>Surveillance measures to support early detection</b> - Measures to run an effective surveillance system for achieving an early detection of a new occurrence (cf. Article 16). This section assumes that the species is not currently present in a Member State, or part of a Member State's territory. <b>This table is repeated for each of the early detection measures identified.</b></p>	
<p><b>Measure description</b></p> <p>Provide a description of the measure, and identify its objective</p>	<p><b>Early detection monitoring by citizen scientists.</b></p> <p>The use of citizen science programs for monitoring invasive plants has gained in popularity in recent years, particularly as new technologies and use of "smart phone" apps has increased. In the context of monitoring for <i>L. japonicum</i> in Europe, citizen scientist might best be utilized in areas not frequented by natural resource managers, or to assist resource managers in parks and recreational areas either casually or as targeted events to detect (and remove) invasive species.</p>

	Another key area where citizen scientists could be effective would be near major nurseries or horticultural areas <i>where L. japonicum</i> may be unintentionally introduced and established, and near the few known locations where <i>L. japonicum</i> was introduced in the E.U. in private gardens and botanical gardens. Historical accounts suggest <i>L. japonicum</i> was cultivated in a number of private collections across the E.U. including in Germany, Portugal, and Italy (EPPO, 2017).																					
<b>Scale of application</b> At what scale is the measure applied? What is the largest scale at which it has been successfully used? Please provide examples, with areas (km <sup>2</sup> or ha) if possible.	A comprehensive review of citizen science programs across multiple continents indicate that monitoring activities range in scale from local to regional (Aceves-Bueno et al., 2015). In this review, 41% of all identified projects were classified as local and 57% of projects were classified as regional.																					
<b>Effectiveness of the measure</b> Is it effective in relation to its objective? Has the measure previously worked, failed?  Please select one of the categories of effectiveness (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.	<table border="1" data-bbox="645 555 1888 619"> <tr> <td><b>Effectiveness of measures</b></td> <td>Effective</td> <td>x</td> <td>Neutral</td> <td></td> <td>Ineffective</td> <td></td> </tr> </table> <p>Rationale: Several studies indicate that citizen science monitoring can be effective and in many cases as accurate as professional monitoring (Crall et al., 2011; Gardiner et al, 2012; Aceves-Bueno et al., 2015). Effectiveness of monitoring programs using citizen scientists is also improved in projects where data can be 'verified' by professionals (Gardiner et al., 2012).</p> <p>On the regional scale, citizen science monitoring can be most efficient and effective when citizens have training and access to a web database where pictures, locations, and other descriptors can be uploaded and available across the region. In Europe, there are a few such apps already developed, including those by the JRC European Alien Species Information Network (EASIN) as well as the UK Plant Tracker app.</p>	<b>Effectiveness of measures</b>	Effective	x	Neutral		Ineffective															
<b>Effectiveness of measures</b>	Effective	x	Neutral		Ineffective																	
<b>Effort required</b> e.g. period of time over which measure need to be applied to have results	<i>L. japonicum</i> is currently not a highly visible or recognized species so effort would be needed in terms of engaging citizens through awareness campaigns and training. If and when the species is detected in greater numbers (either as plantings or escaped) then increased effort would be needed to monitor larger areas. In this case, monitoring would need to occur consistently and in perpetuity.																					
<b>Resources required</b> <sup>1</sup> e.g. cost, staff, equipment etc.	Few resources are needed by citizen scientists themselves except smart phones, internet access, etc. where monitoring is used in conjunction with apps or web-based databases. Most mobile apps likely do not currently contain <i>L. japonicum</i> , and would need to be updated should significant monitoring on a regional level be warranted. Additional staff resources and time would be needed by organizations to oversee training and coordination of volunteers as well as data verification. This could be incorporated into an overall invasive species program.																					
<b>Side effects (incl. potential) – both positive and negative</b>	<table border="1" data-bbox="645 1313 1888 1412"> <tr> <td><b>Environmental effects</b></td> <td>Positive</td> <td>x</td> <td>Neutral or mixed</td> <td></td> <td>Negative</td> <td></td> </tr> <tr> <td><b>Social effects</b></td> <td>Positive</td> <td>x</td> <td>Neutral or mixed</td> <td></td> <td>Negative</td> <td></td> </tr> <tr> <td><b>Economic effects</b></td> <td>Positive</td> <td></td> <td>Neutral or mixed</td> <td>x</td> <td>Negative</td> <td></td> </tr> </table>	<b>Environmental effects</b>	Positive	x	Neutral or mixed		Negative		<b>Social effects</b>	Positive	x	Neutral or mixed		Negative		<b>Economic effects</b>	Positive		Neutral or mixed	x	Negative	
<b>Environmental effects</b>	Positive	x	Neutral or mixed		Negative																	
<b>Social effects</b>	Positive	x	Neutral or mixed		Negative																	
<b>Economic effects</b>	Positive		Neutral or mixed	x	Negative																	



<p>i.e. positive or negative side effects of the measure on public health, environment including non-targeted species, etc.</p> <p>For each of the side effect types please select one of the impact categories (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<p>Citizen Science can have positive effects on society by increasing engagement of the community with important issues. Awareness of the potential impacts of the species through education and training are all suggested to increase the motivation and intent of the public to engage in invasive species monitoring efforts (Crall et al., 2012). A listing of current Citizen Science Projects can be found at European Alien Species Information Network's website at <a href="https://easin.jrc.ec.europa.eu/CitizenScience/Projects">https://easin.jrc.ec.europa.eu/CitizenScience/Projects</a>.</p> <p>Total costs of training and oversight of Citizen Science projects can be minimized by monitoring several invasive species jointly.</p>															
<p><b>Acceptability to stakeholders</b> e.g. impacted economic activities, animal welfare considerations, public perception, etc.</p> <p>Please select one of the categories of acceptability (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<table border="1" data-bbox="645 491 1888 555"> <tr> <td><b>Acceptability to stakeholders</b></td> <td><i>Acceptable</i></td> <td><input checked="" type="checkbox"/></td> <td><i>Neutral or mixed</i></td> <td><input type="checkbox"/></td> <td><i>Unacceptable</i></td> <td><input type="checkbox"/></td> </tr> </table> <p>Rationale: Generally both natural resource agencies and the public view Citizen Science projects positively.</p>								<b>Acceptability to stakeholders</b>	<i>Acceptable</i>	<input checked="" type="checkbox"/>	<i>Neutral or mixed</i>	<input type="checkbox"/>	<i>Unacceptable</i>	<input type="checkbox"/>	
<b>Acceptability to stakeholders</b>	<i>Acceptable</i>	<input checked="" type="checkbox"/>	<i>Neutral or mixed</i>	<input type="checkbox"/>	<i>Unacceptable</i>	<input type="checkbox"/>										
<p><b>Additional cost information</b><sup>1</sup> When not already included above, or in the species Risk Assessment. - implementation cost for Member States - the cost of inaction - the cost-effectiveness - the socio-economic aspects</p> <p>Include quantitative &amp;/or qualitative data, and case studies (incl. from countries outside the EU).</p>	<p>Data collection by Citizen Scientists has been documented to save money for researchers and natural resource agencies. Some estimates suggest that verified citizen science data reduced data collection costs from \$126/hour to \$40/hour (Gardiner et al., 2012), and another study of a forest monitoring project estimated monitoring costs were four times less using citizen scientists (Holck, 2008).</p>															
<p><b>Level of confidence on the information provided</b><sup>2</sup></p> <p>Please select one of the confidence categories along with a statement to support the category chosen. See</p>	<table border="1" data-bbox="645 1185 1924 1249"> <tr> <td><i>Inconclusive</i></td> <td><input type="checkbox"/></td> <td><i>Unresolved</i></td> <td><input type="checkbox"/></td> <td><i>Established but incomplete</i></td> <td><input type="checkbox"/></td> <td><i>Well established</i></td> <td><input checked="" type="checkbox"/></td> </tr> </table> <p>Rationale: Citizen Science programs have gained wide popularity and usage in the last few decades, and their effectiveness has also been well studied and documented.</p>								<i>Inconclusive</i>	<input type="checkbox"/>	<i>Unresolved</i>	<input type="checkbox"/>	<i>Established but incomplete</i>	<input type="checkbox"/>	<i>Well established</i>	<input checked="" type="checkbox"/>
<i>Inconclusive</i>	<input type="checkbox"/>	<i>Unresolved</i>	<input type="checkbox"/>	<i>Established but incomplete</i>	<input type="checkbox"/>	<i>Well established</i>	<input checked="" type="checkbox"/>									

Notes section at the bottom of this document.

**NOTE – this is not related to the effectiveness of the measure**

**Rapid eradication for new introductions** - Measures to achieve eradication at an early stage of invasion, after an early detection of a new occurrence (cf. Article 17). This section assumes that the species is not currently present in a Member State, or part of a Member State's territory. **This table is repeated for each of the eradication measures identified.**

**Measure description**

Provide a description of the measure, and identify its objective

**Application of chemicals (plant protection products).**

Post-emergent herbicides are one of the most effective and efficient ways to eradicate early stages of an infestation of *L. japonicum*. At early stages, chemicals can be applied as a directed foliar spray using hand pumps, backpack sprayers, or other mounted tanks and sprayers that direct application from atop an ATV or small vehicle. Directed sprayer wands can be fitted with an adjustable cone nozzle set to produce a relatively narrow (approximately 20°) cone pattern to direct the spray to very small areas. Herbicides should be applied according to manufacturer instructions and in accordance with EU and national regulations. It is important to note that EU/national/local legislation on the use of plant protection products and biocides needs to be respected and authorities should check to ensure chemicals are licensed for use in their respective countries/regions. Some chemicals described below may be restricted in certain nations and/or certain ecosystems (particularly near water).

Herbicide treatment of *L. japonicum* fern has been investigated in multiple studies in its invaded range in the southeastern United States (Valenta et al., 2001; van Loan, 2006; Minogue et al., 2010; Bohn et al., 2011). The effectiveness of glyphosate treatments was observed in these studies, however it was also noted that metsulfuron treatments were least damaging to surrounding native vegetation, particularly graminoids (Zeller & Leslie 2004). Minogue et al. (2010) and Bohn et al. (2011) examined a range of herbicide types and rates for efficacy in controlling *L. japonicum* and their impact on associated vegetation in both bottomland (wet) hardwood forests and upland (xeric) pine forests. Early control of Japanese climbing fern improved linearly as the glyphosate product rate was increased from 1 percent to 4 percent of the spray solution, though at 2 years after treatment, effectiveness of different glyphosate rates did not differ. Given that re-treatment was needed to control re-sprouting plants and new germinates, use of a 2% glyphosate solution was recommended to reduce costs and potentially avoid adverse effects to associated vegetation. Combinations of glyphosate (2% v:v) and metsulfuron methyl (at 0.375-0.75 g/L) were also effective. A non-ionic surfactant should also be used if not already incorporated into the herbicide product, except in areas near waterways, streams, and wetlands.

From operational experience, best results are obtained with application of foliar herbicides in the later stages of the growing season, from July through early October. However, studies evaluating the timing of treatments to better control spore maturation found reduced spore viability from spores collected off of adult fronds treated between July and no later than early September. Metsulfuron methyl was shown to be more effective than glyphosate for reducing spore viability on individually treated plants

	(Bohn & Thetford 2014), though both herbicides applied individually or in combination similarly reduced spore viability when applied at a larger patch and stand scale (Bohn et al., 2015).						
<b>Scale of application</b> At what scale is the measure applied? What is the largest scale at which it has been successfully used? Please provide examples, with areas (km <sup>2</sup> or ha) if possible.	Directed foliar application by hand pump or backpack sprayer is most efficient for treating individual plants or patches at small scales of tens to hundreds of hectares. The number of hectares treated is a function of number of personnel and hours available for treatment. ATV/small vehicle mounted tanks and directed spray wands can cover larger swaths of land but require existing trails and easy access.						
<b>Effectiveness of the measure</b> Is it effective in relation to its objective? Has the measure previously worked, failed?  Please select one of the categories of effectiveness (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.	<b>Effectiveness of measures</b>	<i>Effective</i>	<input checked="" type="checkbox"/>	<i>Neutral</i>	<input type="checkbox"/>	<i>Ineffective</i>	<input type="checkbox"/>
	Rationale: Several studies have shown that low rates of glyphosate have resulted in 90% or greater control of <i>L. japonicum</i> up to 2 years after treatment on large infestations (van Loan, 2006; Minogue et al., 2010; Bohn et al., 2011). Successful (complete) eradication is possible on smaller, early stage infestations. Individual plants that were germinated from spores and grown for 2 years in contained pots were treated with glyphosate, which resulted in complete dieback of above-ground fronds with no observed resprouting from rhizome biomass 2 years after the treatment (Bohn & Thetford, 2013).						
<b>Effort required</b> e.g. period of time over which measure need to be applied to have results	The level of effort required in early eradication would be determined primarily by the terrain and accessibility to the site and the density and distribution of <i>L. japonicum</i> patches, since directed herbicide applications are generally easy to apply on small populations. Additional effort should be made for follow-up monitoring and treatments to control vegetative resprouts or new gametophytes (germinants).						
<b>Resources required</b> <sup>1</sup> e.g. cost, staff, equipment etc.	Depending on EU and national regulations, trained (and often certified or licensed) professionals will be required to mix and apply herbicides either on public or private land. For early infestations, costs in the United States have generally ranged from \$200-300/hectare (ca. €170 to €260) (S. Miller pers. comm. 2017).						
<b>Side effects (incl. potential) – both positive and negative</b> i.e. positive or negative side effects of the measure on public health, environment including non-targeted species, etc.  For each of the side effect types please select one of the impact categories (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.	<b>Environmental effects</b>	<i>Positive</i>	<input type="checkbox"/>	<i>Neutral or mixed</i>	<input checked="" type="checkbox"/>	<i>Negative</i>	<input type="checkbox"/>
	<b>Social effects</b>	<i>Positive</i>	<input type="checkbox"/>	<i>Neutral or mixed</i>	<input checked="" type="checkbox"/>	<i>Negative</i>	<input type="checkbox"/>
	<b>Economic effects</b>	<i>Positive</i>	<input type="checkbox"/>	<i>Neutral or mixed</i>	<input checked="" type="checkbox"/>	<i>Negative</i>	<input type="checkbox"/>
	Rationale: Environmental: Early eradication efforts should mostly be positive, particularly in contrast to the 'environmental cost' of doing nothing which can over time lead to reduced native plant cover. Care must be given to preventing non-target exposure to broad-spectrum herbicides vegetation as a result of drift or mis-application. However, Bohn et al. (2011) found that when herbicides were directed to <i>L. japonicum</i> , even when intermingling with other native vegetation, effects to non-target native vegetation was minimal. In fact, control of <i>L. japonicum</i> resulted in an increase of 20 to 35 % cover of native plants two years after 90-95% <i>L. japonicum</i> control using glyphosate, compared to an almost 25% loss in native plant cover on untreated plots.						

	<p>Care must also be used when herbicides are applied near water such as in wetlands, near streambanks, or in ditches, if allowed at all under national regulation. Some formulations of glyphosate (and other chemicals) are known to cause harm to aquatic organisms, primarily from the surfactants and additives in the products (Tsui &amp; Chu, 2003), so only herbicide formulations approved for aquatic use (as indicated on the manufacturer’s label and by national regulation) should be used.</p> <p>Social: There are few if any scientifically rigorous studies indicating side effects of small-scale herbicide applications on public health, particularly at the rates required to control early infestations of <i>L. japonicum</i>. However, herbicide applicators do need to wear appropriate personal protective equipment (PPE) such as safety glasses, gloves, and other outerwear to prevent accidental exposure to chemicals. Attention must also be paid to apply herbicides in low wind conditions to avoid accidental exposure.</p> <p>Economic: Though there is a cost associated with eradication at early stages of establishment, it is likely far less than the cost of control efforts for widespread and heavy infestations, particularly in regard to the cost of purchasing herbicides needed for small versus large applications. Herbicide applications of small-scale infestations are quite low compared to treatment costs of large-scale infestations; often reducing costs by as much as \$300/ hectare (ca. €260) (S. Miller pers. comm. 2017).</p>								
<p><b>Acceptability to stakeholders</b> e.g. impacted economic activities, animal welfare considerations, public perception, etc.</p> <p>Please select one of the categories of acceptability (with an ‘X’), and provide a rationale, with supporting evidence and examples if possible.</p>	<p><b>Acceptability to stakeholders</b></p>	<p>Acceptable</p>		<p>Neutral or mixed</p>	<p>x</p>	<p>Unacceptable</p>			
<p><b>Additional cost information</b><sup>1</sup> When not already included above, or in the species Risk Assessment. - implementation cost for Member States - the cost of inaction - the cost-effectiveness - the socio-economic aspects</p> <p>Include quantitative &amp;/or qualitative data, and case studies (incl. from countries outside the EU).</p>	<p>Rationale: Given the efficiency and efficacy of herbicides for eradication of <i>L. japonicum</i>, most natural resource professionals, at least in countries where <i>L. japonicum</i> is already well-established and invasive, approve of the use of chemical methods. Public stakeholders may be more wary of herbicide use, unless educated about the limited side effects with proper application.</p> <p>See comments on side-effects above for information on cost of inaction.</p>								
<p><b>Level of confidence on the information provided</b><sup>2</sup></p>	<p>Inconclusive</p>		<p>Unresolved</p>		<p>Established but incomplete</p>		<p>Well established</p>	<p>x</p>	

<p>Please select one of the confidence categories along with a statement to support the category chosen. See <i>Notes</i> section at the bottom of this document.</p> <p><b>NOTE – this is not related to the effectiveness of the measure</b></p>	<p>Rationale:</p> <p>As noted above, there is clear evidence of the application of herbicides, particularly glyphosate and/or metsulfuron methyl.</p>
--	---

<p><b>Rapid eradication for new introductions</b> - Measures to achieve eradication <u>at an early stage of invasion</u>, after an early detection of a new occurrence (cf. Article 17). This section assumes that the species is not currently present in a Member State, or part of a Member State's territory. <b>This table is repeated for each of the eradication measures identified.</b></p>	
<p><b>Measure description</b> Provide a description of the measure, and identify its objective</p>	<p><b>Manual removal of the plant and its root/rhizome system.</b></p> <p>For small infestations, particularly in the first year of establishment, mechanically removing <i>L. japonicum</i> may be possible but must include complete removal of the belowground root and rhizome system, since it readily spreads and resprouts by rhizomes. For plants detected in its first year of establishment, the aboveground vines can be gently pulled out or can be cut near the base if intertwined with other vegetation. Below-ground biomass will need to be dug out, by hand in loose soil or with a small shovel or trowel. Roots and rhizomes should be separated from soil and disposed of properly, either by burning or desiccating and killing plant material in plastic bags or under tarps. Above-ground plant material with spores present should be disposed of in a similar manner. This method of rapid eradication may be an alternative approach where chemical use is restricted or undesirable.</p>
<p><b>Scale of application</b> At what scale is the measure applied? What is the largest scale at which it has been successfully used? Please provide examples, with areas (km<sup>2</sup> or ha) if possible.</p>	<p>This method of eradication is best applied at very small scale, from individual plants to small scattered patches (less than a few m<sup>2</sup>). However total area (ha) of individual plants that can be treated in this manner would be a function of number of personnel or volunteers available. This measure would not be effective or efficient for large-scale or heavy infestations.</p>

<p><b>Effectiveness of the measure</b> Is it effective in relation to its objective? Has the measure previously worked, failed?</p> <p>Please select one of the categories of effectiveness (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<table border="1" data-bbox="640 151 1888 220"> <tr> <td><b>Effectiveness of measures</b></td> <td><i>Effective</i></td> <td>x</td> <td><i>Neutral</i></td> <td></td> <td><i>Ineffective</i></td> <td></td> </tr> </table> <p>Rationale: It is very likely that such mechanical treatments can be effective at very small scales, particularly for individual plants or very small patches. It would be appropriate for homeowners, gardeners, recreationists, and volunteers in small areas. Manual treatments would likely be less effective and efficient in larger natural areas where <i>L. japonicum</i> may be more intertwined with native vegetation and separation of <i>L. japonicum</i> roots from roots of native vegetation would be difficult. Manual treatments have typically not been used in other countries where <i>L. japonicum</i> has invaded, so there is little documentation of the effectiveness and efficiency of this treatment method, even at small scales.</p>	<b>Effectiveness of measures</b>	<i>Effective</i>	x	<i>Neutral</i>		<i>Ineffective</i>															
<b>Effectiveness of measures</b>	<i>Effective</i>	x	<i>Neutral</i>		<i>Ineffective</i>																	
<p><b>Effort required</b> e.g. period of time over which measure need to be applied to have results</p>	<p>As with chemical treatments, the level of effort required in early eradication would be determined primarily by the time and effort to detect small-scale patches across a site, since hand pulling, clipping, and digging out of roots is relatively easy. Additional effort would be needed for follow-up monitoring and treatments to control vegetative resprouts or new gametophytes.</p>																					
<p><b>Resources required</b><sup>1</sup> e.g. cost, staff, equipment etc.</p>	<p>The main resource and cost will be in the personnel needed to conduct treatments (or oversee volunteers). Other resources needed include simple tools for cutting and digging as well as bags, tarps, etc for proper disposal of dead plant material.</p>																					
<p><b>Side effects (incl. potential) – both positive and negative</b> i.e. positive or negative side effects of the measure on public health, environment including non-targeted species, etc.</p> <p>For each of the side effect types please select one of the impact categories (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<table border="1" data-bbox="640 756 1888 863"> <tr> <td><b>Environmental effects</b></td> <td><i>Positive</i></td> <td></td> <td><i>Neutral or mixed</i></td> <td>x</td> <td><i>Negative</i></td> <td></td> </tr> <tr> <td><b>Social effects</b></td> <td><i>Positive</i></td> <td>x</td> <td><i>Neutral or mixed</i></td> <td></td> <td><i>Negative</i></td> <td></td> </tr> <tr> <td><b>Economic effects</b></td> <td><i>Positive</i></td> <td></td> <td><i>Neutral or mixed</i></td> <td>x</td> <td><i>Negative</i></td> <td></td> </tr> </table> <p>Rationale: Environmental: While eradication of <i>L. japonicum</i> in the early stages of infestation will have beneficial long-term effects on native vegetation by preventing the vine from creating dense mats and smothering vegetation, there are some potential negative effects of the soil disturbance created by digging out roots as well as potential damage to native plants due to inadvertent clipping of foliage or roots where <i>L. japonicum</i> is intertwined. This negative impact would be minimal however, in very early stages where the plant is small and isolated.</p> <p>Social: Volunteers tend to prefer mechanical treatments and derive satisfaction from removing invasive plants.</p> <p>Economic: As with chemical treatments, there is a cost associated with eradication at early stages of establishment, but it is likely far less than the cost of control efforts for wide-spread and heavy infestations. Exact costs for small scale manual treatments are not available.</p>	<b>Environmental effects</b>	<i>Positive</i>		<i>Neutral or mixed</i>	x	<i>Negative</i>		<b>Social effects</b>	<i>Positive</i>	x	<i>Neutral or mixed</i>		<i>Negative</i>		<b>Economic effects</b>	<i>Positive</i>		<i>Neutral or mixed</i>	x	<i>Negative</i>	
<b>Environmental effects</b>	<i>Positive</i>		<i>Neutral or mixed</i>	x	<i>Negative</i>																	
<b>Social effects</b>	<i>Positive</i>	x	<i>Neutral or mixed</i>		<i>Negative</i>																	
<b>Economic effects</b>	<i>Positive</i>		<i>Neutral or mixed</i>	x	<i>Negative</i>																	
<p><b>Acceptability to stakeholders</b></p>	<table border="1" data-bbox="640 1321 1888 1390"> <tr> <td><b>Acceptability to stakeholders</b></td> <td><i>Acceptable</i></td> <td>x</td> <td><i>Neutral or mixed</i></td> <td></td> <td><i>Unacceptable</i></td> <td></td> </tr> </table>	<b>Acceptability to stakeholders</b>	<i>Acceptable</i>	x	<i>Neutral or mixed</i>		<i>Unacceptable</i>															
<b>Acceptability to stakeholders</b>	<i>Acceptable</i>	x	<i>Neutral or mixed</i>		<i>Unacceptable</i>																	

<p>e.g. impacted economic activities, animal welfare considerations, public perception, etc.</p> <p>Please select one of the categories of acceptability (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<p>Rationale: Manual treatments tend to be more acceptable to the general public than herbicides and may be more appropriate for use near waterways or other sensitive areas.</p>								
<p><b>Additional cost information</b> <sup>1</sup></p> <p>When not already included above, or in the species Risk Assessment.</p> <ul style="list-style-type: none"> <li>- implementation cost for Member States</li> <li>- the cost of inaction</li> <li>- the cost-effectiveness</li> <li>- the socio-economic aspects</li> </ul> <p>Include quantitative &amp;/or qualitative data, and case studies (incl. from countries outside the EU).</p>	<p>As stated in other sections, the cost of inaction in terms of potential detriment to native vegetation and habitat far exceeds the cost and effort of early eradication. There are trade-offs in cost and cost-effectiveness between manual and chemical treatments. While there is an additional cost associated with purchasing herbicides, chemical applications are also quite effective for eradication, and more so as infestations grow in size as compared to manual treatments. However, manual treatments require little training and/or licensing to implement in contrast to chemical applications.</p>								
<p><b>Level of confidence on the information provided</b> <sup>2</sup></p> <p>Please select one of the confidence categories along with a statement to support the category chosen. See <i>Notes</i> section at the bottom of this document.</p> <p><b>NOTE – this is not related to the effectiveness of the measure</b></p>	<i>Inconclusive</i>	<input type="checkbox"/>	<i>Unresolved</i>	<input type="checkbox"/>	<i>Established but incomplete</i>	<input checked="" type="checkbox"/>	<i>Well established</i>	<input type="checkbox"/>	
<p>Rationale: Mechanical removal of invasive plants has been documented to be effective, though more so for shallow-rooted species (eg. <a href="#">Flory &amp; Lewis, 2009</a>). It is assumed that as long as entire root systems are removed that this will be effective for <i>L. japonicum</i> at early stages of establishment as well, but effectiveness of manual treatments has not been studied or documented.</p>									

**Management** - Measures to achieve management of the species once it has become widely spread within a Member State, or part of a Member State's territory. (cf. Article 19), i.e. **not** at an early stage of invasion (see Rapid eradication table above). These measures can be aimed at eradication, population control or containment of a population of the species. **This table is repeated for each of the management measures identified.**

<p><b>Measure description</b> Provide a description of the measure, and identify its objective</p>	<p><b>Application of chemicals (plant protection products)</b></p> <p>Herbicide application is the primary method by which <i>L. japonicum</i> is managed in widespread and highly invaded areas. Glyphosate at a rate between 2-4% v:V solution is recommended for large scale infestations, though metsulfuron methyl at a rate of 0.75 g/L applied may at least provide short-term (1-2yr) control.</p> <p>Broadcast applications of herbicides in wide-spread areas can be made using backpack sprayer, mist blowers mounted on ATV's or other vehicles, and even by aerial application for extensive monocultures. Herbicides should be applied according to manufacturer instructions and in accordance with EU and national regulations. It is important to note that EU/national/local legislation on the use of plant protection products and biocides needs to be respected and authorities should check to ensure chemicals are licensed for use in their respective countries/regions. Some chemicals may be restricted in certain nations and/or certain ecosystems (particularly near water). Most critical with broadcast applications is the prevention of drift into non-target areas or onto non-target plants.</p>						
<p><b>Scale of application</b> At what scale is the measure applied? What is the largest scale at which it has been successfully used? Please provide examples, with areas (km<sup>2</sup> or ha) if possible.</p>	<p>Broadcast foliar herbicide treatments can be applied at scales ranging anywhere from tens to hundreds of hectares, and the scale of application depends primarily on the amount of personnel and hours available for treatment, equipment or vehicles available, as well as terrain and accessibility to the site.</p>						
<p><b>Effectiveness of the measure</b> Is it effective in relation to its objective? Has the measure previously worked, failed?</p> <p>Please select one of the categories of effectiveness (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<p><i>Effectiveness of measures</i></p>	<p><i>Effective</i></p>	<p>x</p>	<p><i>Neutral</i></p>		<p><i>Ineffective</i></p>	
<p><b>Effort required</b> e.g. period of time over which measure need to be applied to have results</p>	<p>Rationale: Several studies have documented the effectiveness of controlling (though not eradicating) <i>L. japonicum</i> with glyphosate, or a combination glyphosate and metsulfuron methyl, at least as small plot and stand levels. In both bottomland (wet) hardwood forests and upland (xeric) pine forests where <i>L. japonicum</i> cover ranged from 40-60% prior to treatment, <a href="#">Minogue et al. (2010)</a> and <a href="#">Bohn et al. (2011)</a> found 90-95% reduction in cover of fern two years after treatment. However, the degree to which small-scale control studies translate to large-scale invasive plant management programs has not been well studied (<a href="#">Kettenring &amp; Adams, 2011</a>). Successful <i>L. japonicum</i> control at large scales will depend on the effectiveness of the broadcast application in making complete contact with target foliage across a wide area.</p> <p>Monitoring and repeated treatments will be necessary as it often takes several treatments to kill the entire root system. Additionally, because spores may travel by wind for several kilometres, repeated treatments of new gametophytes and growth will likely be needed if <i>L. japonicum</i> is still present off-site.</p>						



<p><b>Resources required</b><sup>1</sup> e.g. cost, staff, equipment etc.</p>	<p>Depending on EU and national regulations, trained (and often certified or licensed) professionals will be required to mix and apply herbicides either on public or private land. Costs based on experience in the southeastern United States, from six years of data and 147 treatment records, indicate that to effectively treat and control <i>L. japonicum</i> ranges from \$200 to as much as \$500/hectare (ca. €173 to €433) (S. Miller pers. comm. 2017).</p>						
<p><b>Side effects (incl. potential) – both positive and negative</b> i.e. positive or negative side effects of the measure on public health, environment including non-targeted species, etc.</p> <p>For each of the side effect types please select one of the impact categories (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<p><b>Environmental effects</b></p>		<p>Positive</p>	<p>Neutral or mixed</p>	<p>x</p>	<p>Negative</p>	
	<p><b>Social effects</b></p>		<p>Positive</p>	<p>Neutral or mixed</p>	<p>x</p>	<p>Negative</p>	
	<p><b>Economic effects</b></p>		<p>Positive</p>	<p>Neutral or mixed</p>	<p>x</p>	<p>Negative</p>	
	<p>Rationale: Side effects are generally the same as those noted in the section on Rapid Eradication using chemicals. The positive environmental effects of managing <i>L. japonicum</i> are thought to outweigh any potential non-target impacts. However, with widespread infestations and application of broad spectrum chemicals, increased diligence must be given to avoid non-target damage and/or human exposure from drift. This can be avoided with proper application rates under appropriate environmental conditions (e.g. low wind) using appropriate personal protective equipment.</p>						
<p><b>Acceptability to stakeholders</b> e.g. impacted economic activities, animal welfare considerations, public perception, etc.</p> <p>Please select one of the categories of acceptability (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<p><b>Acceptability to stakeholders</b></p>		<p>Acceptable</p>	<p>Neutral or mixed</p>	<p>x</p>	<p>Unacceptable</p>	
<p><b>Additional cost information</b><sup>1</sup> When not already included above, or in the species Risk Assessment. - implementation cost for Member States - the cost of inaction - the cost-effectiveness - the socio-economic aspects</p> <p>Include quantitative &amp;/or qualitative data, and case studies (incl. from countries outside the EU).</p>	<p>See comments in section on Rapid Eradication by chemical application.</p>						

<p><b>Level of confidence on the information provided</b><sup>2</sup></p> <p>Please select one of the confidence categories along with a statement to support the category chosen. See <i>Notes</i> section at the bottom of this document.</p> <p><b>NOTE – this is not related to the effectiveness of the measure</b></p>	<i>Inconclusive</i>		<i>Unresolved</i>		<i>Established but incomplete</i>		<i>Well established</i>	x
	<p>Rationale: As noted above, there is clear evidence of the effectiveness of herbicides, particularly glyphosate and/or metsulfuron methyl.</p>							

<p><b>Management</b> - Measures to achieve management of the species once it has become widely spread within a Member State, or part of a Member State's territory. (cf. Article 19), i.e. <b>not</b> at an early stage of invasion (see Rapid eradication table above). These measures can be aimed at eradication, population control or containment of a population of the species. <b>This table is repeated for each of the management measures identified.</b></p>	
<p><b>Measure description</b> Provide a description of the measure, and identify its objective</p>	<p><b>Integrated Vegetation Management: Mechanical (Cutting) and Chemical Application</b></p> <p>For widespread and heavily invaded sites, mechanical treatments such as pulling or cutting are ineffectual as stand-alone treatments for controlling <i>L. japonicum</i> because of the inability of these methods to control resprouting from rhizomes; however mechanical treatments may be used in conjunction with chemical treatments as an integrated vegetation management approach. For example, for large infestations where <i>L. japonicum</i> is climbing trees or tall vegetation, the vines may be clipped at about 0.5-1m from the ground and herbicide only applied to the foliage on just the lower part of the cut vines. This approach has been used more extensively in the southeastern United States on <i>Lygodium microphyllum</i> and is known as a "poodle cut". This integrated approach thus limits the volume of herbicide applied in the landscape. However, this approach may not work where <i>L. japonicum</i> forms a dense low-lying mat over low-lying vegetation as it would be difficult to distinguish the cut fronds unless raked away.</p>
<p><b>Scale of application</b> At what scale is the measure applied? What is the largest scale at which it has been successfully used? Please provide examples, with areas (km<sup>2</sup> or ha) if possible.</p>	<p>Integrated vegetation management programs can be implemented at the same large scales as chemical-only programs (see Section above for more detail).</p>

<p><b>Effectiveness of the measure</b> Is it effective in relation to its objective? Has the measure previously worked, failed?</p> <p>Please select one of the categories of effectiveness (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<table border="1" data-bbox="640 151 1888 220"> <tr> <td><b>Effectiveness of measures</b></td> <td><i>Effective</i></td> <td><input checked="" type="checkbox"/></td> <td><i>Neutral</i></td> <td><input type="checkbox"/></td> <td><i>Ineffective</i></td> <td><input type="checkbox"/></td> </tr> </table> <p>Rationale: Several studies have documented the effectiveness of integrated vegetation management of <i>L. japonicum</i>, so long as herbicides are included as one of the treatments. <a href="#">Hutchinson and Langeland (2007)</a> found the greatest translocation of herbicides, especially glyphosate, to the root system of <i>Lygodium</i> was when applying herbicides only to the base foliage of cut vines rather than a complete foliar application, resulting in 85-90% control.</p>	<b>Effectiveness of measures</b>	<i>Effective</i>	<input checked="" type="checkbox"/>	<i>Neutral</i>	<input type="checkbox"/>	<i>Ineffective</i>	<input type="checkbox"/>														
<b>Effectiveness of measures</b>	<i>Effective</i>	<input checked="" type="checkbox"/>	<i>Neutral</i>	<input type="checkbox"/>	<i>Ineffective</i>	<input type="checkbox"/>																
<p><b>Effort required</b> e.g. period of time over which measure need to be applied to have results</p>	<p>Even with integrated approaches, monitoring and repeated treatments will be necessary as it often takes several treatments to kill the entire root system and suppress spore germination over time.</p>																					
<p><b>Resources required</b><sup>1</sup> e.g. cost, staff, equipment etc.</p>	<p>Incorporating mechanical cutting treatments into control methods can reduce the amount and therefore costs of herbicides, but it does require more personnel and labor, which might offset those savings. In the southeastern United States, estimates for manual cutting and herbicide treatments can be as high as \$800 /ha (ca. €695) in large-scale infestations (<a href="#">Hutchinson &amp; Langeland, 2007</a>).</p>																					
<p><b>Side effects (incl. potential) – both positive and negative</b> i.e. positive or negative side effects of the measure on public health, environment including non-targeted species, etc.</p> <p>For each of the side effect types please select one of the impact categories (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<table border="1" data-bbox="640 751 1888 858"> <tr> <td><b>Environmental effects</b></td> <td><i>Positive</i></td> <td><input checked="" type="checkbox"/></td> <td><i>Neutral or mixed</i></td> <td><input type="checkbox"/></td> <td><i>Negative</i></td> <td><input type="checkbox"/></td> </tr> <tr> <td><b>Social effects</b></td> <td><i>Positive</i></td> <td><input checked="" type="checkbox"/></td> <td><i>Neutral or mixed</i></td> <td><input type="checkbox"/></td> <td><i>Negative</i></td> <td><input type="checkbox"/></td> </tr> <tr> <td><b>Economic effects</b></td> <td><i>Positive</i></td> <td><input type="checkbox"/></td> <td><i>Neutral or mixed</i></td> <td><input checked="" type="checkbox"/></td> <td><i>Negative</i></td> <td><input type="checkbox"/></td> </tr> </table> <p>Rationale: As noted previously, management and control of widespread infestations will have beneficial long-term effects on sites where native vegetation can recover or be restored. Further, a reduction in the volume of herbicides being applied using integrated vegetation management may limit potential impacts of broad spectrum herbicides to non-target, desirable species as well as decrease potential human exposure. Economic effects remain neutral or mixed because of the increased costs of implementing the combination of mechanical and chemical applications, though this may offset potential negative impacts of <i>L. japonicum</i> to agricultural products or timber.</p>	<b>Environmental effects</b>	<i>Positive</i>	<input checked="" type="checkbox"/>	<i>Neutral or mixed</i>	<input type="checkbox"/>	<i>Negative</i>	<input type="checkbox"/>	<b>Social effects</b>	<i>Positive</i>	<input checked="" type="checkbox"/>	<i>Neutral or mixed</i>	<input type="checkbox"/>	<i>Negative</i>	<input type="checkbox"/>	<b>Economic effects</b>	<i>Positive</i>	<input type="checkbox"/>	<i>Neutral or mixed</i>	<input checked="" type="checkbox"/>	<i>Negative</i>	<input type="checkbox"/>
<b>Environmental effects</b>	<i>Positive</i>	<input checked="" type="checkbox"/>	<i>Neutral or mixed</i>	<input type="checkbox"/>	<i>Negative</i>	<input type="checkbox"/>																
<b>Social effects</b>	<i>Positive</i>	<input checked="" type="checkbox"/>	<i>Neutral or mixed</i>	<input type="checkbox"/>	<i>Negative</i>	<input type="checkbox"/>																
<b>Economic effects</b>	<i>Positive</i>	<input type="checkbox"/>	<i>Neutral or mixed</i>	<input checked="" type="checkbox"/>	<i>Negative</i>	<input type="checkbox"/>																
<p><b>Acceptability to stakeholders</b> e.g. impacted economic activities, animal welfare considerations, public perception, etc.</p> <p>Please select one of the categories of acceptability (with an 'X'), and</p>	<table border="1" data-bbox="640 1155 1888 1224"> <tr> <td><b>Acceptability to stakeholders</b></td> <td><i>Acceptable</i></td> <td><input checked="" type="checkbox"/></td> <td><i>Neutral or mixed</i></td> <td><input type="checkbox"/></td> <td><i>Unacceptable</i></td> <td><input type="checkbox"/></td> </tr> </table> <p>Rationale: Natural resource managers likely will find the integrated approach acceptable, and both professionals and the public typically support efforts to reduce herbicide use.</p>	<b>Acceptability to stakeholders</b>	<i>Acceptable</i>	<input checked="" type="checkbox"/>	<i>Neutral or mixed</i>	<input type="checkbox"/>	<i>Unacceptable</i>	<input type="checkbox"/>														
<b>Acceptability to stakeholders</b>	<i>Acceptable</i>	<input checked="" type="checkbox"/>	<i>Neutral or mixed</i>	<input type="checkbox"/>	<i>Unacceptable</i>	<input type="checkbox"/>																

provide a rationale, with supporting evidence and examples if possible.									
<p><b>Additional cost information</b> <sup>1</sup></p> <p>When not already included above, or in the species Risk Assessment.</p> <ul style="list-style-type: none"> <li>- implementation cost for Member States</li> <li>- the cost of inaction</li> <li>- the cost-effectiveness</li> <li>- the socio-economic aspects</li> </ul> <p>Include quantitative &amp;/or qualitative data, and case studies (incl. from countries outside the EU).</p>	<p>Integrated vegetation management approaches are not necessarily more effective than herbicide-only treatments in widespread and heavy infestations, and the costs are often similar if not greater. The trade-off and advantages are mostly environmental and social in regards to the reduction in herbicide use.</p>								
<p><b>Level of confidence on the information provided</b> <sup>2</sup></p> <p>Please select one of the confidence categories along with a statement to support the category chosen. See <i>Notes</i> section at the bottom of this document.</p> <p><b>NOTE – this is not related to the effectiveness of the measure</b></p>	<i>Inconclusive</i>	x	<i>Unresolved</i>	x	<i>Established but incomplete</i>	x	<i>Well established</i>	x	
		<p>Rationale:</p> <p>As noted above, there is documented evidence of the effectiveness of integrated vegetation management for <i>L. japonicum</i>.</p>							

<p><b>Management</b> - Measures to achieve management of the species once it has become widely spread within a Member State, or part of a Member State's territory. (cf. Article 19), i.e. <b>not</b> at an early stage of invasion (see Rapid eradication table above). These measures can be aimed at eradication, population control or containment of a population of the species. <b>This table is repeated for each of the management measures identified.</b></p>	
<p><b>Measure description</b></p> <p>Provide a description of the measure, and identify its objective</p>	<p><b>Integrated Vegetation Management: Cultural (Prescribed Fire) and Chemical Application</b></p> <p>Prescribed fire alone tends to promote resprouting of <i>L. japonicum</i> from rhizomes; however when implemented in conjunction with chemical treatments, it can be effective as an integrated vegetation management approach. In heavily infested areas, prescribed burns can reduce thick mats of <i>L. japonicum</i>, and chemicals would be applied to resprouting vegetation. This approach could reduce the volume of chemicals applied to an area. <a href="#">Bohn and Minogue (2011)</a> found that a period of six weeks of growth</p>

	<p>post-fire was sufficient for herbicide absorption and translocation, applied at similar rates as mentioned in the Chemical Application section above. Chemicals can also be applied before prescribed fire application, but should be done at least a month after the active growing season begins and at least 6 weeks prior to burning to allow time for translocation of herbicides throughout the entire plant.</p> <p>This integrated approach would be most appropriate in fire-dependent ecosystems. It should only be considered where <i>L. japonicum</i> has formed dense mats over groundcover or low lying shrubs. It should not be used where <i>L. japonicum</i> has climbed up and around trees, as this could lead to severe crown fires.</p>																					
<p><b>Scale of application</b> At what scale is the measure applied? What is the largest scale at which it has been successfully used? Please provide examples, with areas (km<sup>2</sup> or ha) if possible.</p>	<p>Integrated vegetation management programs can be implemented at the same large scales as chemical-only programs.</p>																					
<p><b>Effectiveness of the measure</b> Is it effective in relation to its objective? Has the measure previously worked, failed?</p> <p>Please select one of the categories of effectiveness (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<table border="1" data-bbox="645 687 1888 751"> <tr> <td><i>Effectiveness of measures</i></td> <td><i>Effective</i></td> <td><input checked="" type="checkbox"/></td> <td><i>Neutral</i></td> <td><input type="checkbox"/></td> <td><i>Ineffective</i></td> <td><input type="checkbox"/></td> </tr> </table> <p>Rationale: Use of chemicals in conjunction with prescribed fire has been used effectively for control of <i>L. japonicum</i> in the southeastern United States. In a replicated study of timing of herbicides with prescribed fire, <a href="#">Bohn and Minogue (2010)</a> found that plots treated either in July before burning, or in August or September six weeks after burning, resulted in between 85-100% reductions in fern cover. Greater reductions were observed with hotter burns.</p>	<i>Effectiveness of measures</i>	<i>Effective</i>	<input checked="" type="checkbox"/>	<i>Neutral</i>	<input type="checkbox"/>	<i>Ineffective</i>	<input type="checkbox"/>														
<i>Effectiveness of measures</i>	<i>Effective</i>	<input checked="" type="checkbox"/>	<i>Neutral</i>	<input type="checkbox"/>	<i>Ineffective</i>	<input type="checkbox"/>																
<p><b>Effort required</b> e.g. period of time over which measure need to be applied to have results</p>	<p>Even with integrated approaches, monitoring and repeated treatments will be necessary as it often takes several treatments to kill the entire root system and suppress spore germination over time.</p>																					
<p><b>Resources required</b><sup>1</sup> e.g. cost, staff, equipment etc.</p>	<p>This integrated approach can reduce the amount and therefore costs of chemical control but it does require more personnel and labor, which might offset those cost savings. Use of prescribed fire has been shown to reduce the amount of herbicide by half (<a href="#">Stocker et al., 2008</a>). Costs for implementing prescribed fire can add an additional \$50 to \$500 per hectare (ca. €43 to €434) to the cost of chemical application for invasive plant control (<a href="#">Cleaves et al., 1999</a>).</p>																					
<p><b>Side effects (incl. potential) – both positive and negative</b> i.e. positive or negative side effects of the measure on public health,</p>	<table border="1" data-bbox="645 1291 1888 1390"> <tr> <td><i>Environmental effects</i></td> <td><i>Positive</i></td> <td><input type="checkbox"/></td> <td><i>Neutral or mixed</i></td> <td><input checked="" type="checkbox"/></td> <td><i>Negative</i></td> <td><input type="checkbox"/></td> </tr> <tr> <td><i>Social effects</i></td> <td><i>Positive</i></td> <td><input type="checkbox"/></td> <td><i>Neutral or mixed</i></td> <td><input checked="" type="checkbox"/></td> <td><i>Negative</i></td> <td><input type="checkbox"/></td> </tr> <tr> <td><i>Economic effects</i></td> <td><i>Positive</i></td> <td><input type="checkbox"/></td> <td><i>Neutral or mixed</i></td> <td><input checked="" type="checkbox"/></td> <td><i>Negative</i></td> <td><input type="checkbox"/></td> </tr> </table>	<i>Environmental effects</i>	<i>Positive</i>	<input type="checkbox"/>	<i>Neutral or mixed</i>	<input checked="" type="checkbox"/>	<i>Negative</i>	<input type="checkbox"/>	<i>Social effects</i>	<i>Positive</i>	<input type="checkbox"/>	<i>Neutral or mixed</i>	<input checked="" type="checkbox"/>	<i>Negative</i>	<input type="checkbox"/>	<i>Economic effects</i>	<i>Positive</i>	<input type="checkbox"/>	<i>Neutral or mixed</i>	<input checked="" type="checkbox"/>	<i>Negative</i>	<input type="checkbox"/>
<i>Environmental effects</i>	<i>Positive</i>	<input type="checkbox"/>	<i>Neutral or mixed</i>	<input checked="" type="checkbox"/>	<i>Negative</i>	<input type="checkbox"/>																
<i>Social effects</i>	<i>Positive</i>	<input type="checkbox"/>	<i>Neutral or mixed</i>	<input checked="" type="checkbox"/>	<i>Negative</i>	<input type="checkbox"/>																
<i>Economic effects</i>	<i>Positive</i>	<input type="checkbox"/>	<i>Neutral or mixed</i>	<input checked="" type="checkbox"/>	<i>Negative</i>	<input type="checkbox"/>																

<p>environment including non-targeted species, etc.</p> <p>For each of the side effect types please select one of the impact categories (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<p>Rationale: As noted previously, management and control of widespread infestations will have beneficial long-term effects on sites where native vegetation can recover or be restored. Further, a reduction in the volume of herbicides being applied using integrated vegetation management may limit potential impacts of broad spectrum herbicides to non-target, desirable species, but there are still limited risks. There are also certain risks associated with using prescribed fire (e.g. where fire might escape due to unpredicted weather or other circumstances). Additionally, while this integrated approach can also reduce the volume of chemicals used and potential exposure of herbicides to people, prescribed fire can create smoke management issues that must be addressed. Economic effects remain neutral or mixed because of the increased costs of implementing an Integrated Vegetation Management system; though this may offset potential negative impacts of <i>L. japonicum</i> to agricultural products or timber.</p>															
<p><b>Acceptability to stakeholders</b> e.g. impacted economic activities, animal welfare considerations, public perception, etc.</p> <p>Please select one of the categories of acceptability (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<table border="1" data-bbox="645 448 1888 517"> <tr> <td data-bbox="645 448 927 517"><b>Acceptability to stakeholders</b></td> <td data-bbox="927 448 1151 517"><i>Acceptable</i></td> <td data-bbox="1151 448 1211 517"></td> <td data-bbox="1211 448 1509 517"><i>Neutral or mixed</i></td> <td data-bbox="1509 448 1570 517">x</td> <td data-bbox="1570 448 1832 517"><i>Unacceptable</i></td> <td data-bbox="1832 448 1888 517"></td> </tr> </table> <p>Rationale: Natural resource managers likely will find the integrated approach acceptable, and both professionals and the public typically support efforts to reduce herbicide use. However, the general public may not favour the use of prescribed fire unless this cultural treatment has been a long-standing practice in the community (Fernandes, 2013). High wildfire seasons may also contribute to concern over the use of prescribed fire.</p>								<b>Acceptability to stakeholders</b>	<i>Acceptable</i>		<i>Neutral or mixed</i>	x	<i>Unacceptable</i>		
<b>Acceptability to stakeholders</b>	<i>Acceptable</i>		<i>Neutral or mixed</i>	x	<i>Unacceptable</i>											
<p><b>Additional cost information</b><sup>1</sup> When not already included above, or in the species Risk Assessment.</p> <ul style="list-style-type: none"> <li>- implementation cost for Member States</li> <li>- the cost of inaction</li> <li>- the cost-effectiveness</li> <li>- the socio-economic aspects</li> </ul> <p>Include quantitative &amp;/or qualitative data, and case studies (incl. from countries outside the EU).</p>	<p>Integrated vegetation management approaches are not necessarily more effective than herbicide-only treatments in widespread and heavy infestations, and the costs are often similar if not greater. The trade-off and advantage is mostly environmental and social in regards to the reduction in herbicide use.</p>															
<p><b>Level of confidence on the information provided</b><sup>2</sup></p> <p>Please select one of the confidence categories along with a statement to support the category chosen. See <i>Notes</i> section at the bottom of this document.</p>	<table border="1" data-bbox="645 1145 1924 1214"> <tr> <td data-bbox="645 1145 913 1214"><i>Inconclusive</i></td> <td data-bbox="913 1145 965 1214"></td> <td data-bbox="965 1145 1173 1214"><i>Unresolved</i></td> <td data-bbox="1173 1145 1227 1214"></td> <td data-bbox="1227 1145 1509 1214"><i>Established but incomplete</i></td> <td data-bbox="1509 1145 1570 1214"></td> <td data-bbox="1570 1145 1868 1214"><i>Well established</i></td> <td data-bbox="1868 1145 1924 1214">x</td> </tr> </table> <p>Rationale: As noted above, there is documented evidence of the effectiveness of integrated vegetation management for <i>L. japonicum</i>.</p>								<i>Inconclusive</i>		<i>Unresolved</i>		<i>Established but incomplete</i>		<i>Well established</i>	x
<i>Inconclusive</i>		<i>Unresolved</i>		<i>Established but incomplete</i>		<i>Well established</i>	x									

**NOTE – this is not related to the effectiveness of the measure**

**Management** - Measures to achieve management of the species once it has become widely spread within a Member State, or part of a Member State's territory. (cf. Article 19), i.e. **not** at an early stage of invasion (see Rapid eradication table above). These measures can be aimed at eradication, population control or containment of a population of the species. **This table is repeated for each of the management measures identified.**

<p><b>Measure description</b> Provide a description of the measure, and identify its objective</p>	<p><b>Biological control</b></p> <p>At the present time, biological control agents have not been sufficiently tested for <i>L. japonicum</i> as a viable control option. A rust-forming organism (<i>Puccinia lygodii</i>) is currently being evaluated in central Florida, USA as a potential biological control.</p> <p><i>Neomusotima conspurcatalis</i> (a <i>Lygodium</i> defoliator moth) was released in peninsular Florida as a biocontrol of <i>L. microphyllum</i> (Smith et al., 2014). This moth is genus specific and also will feed on <i>L. japonicum</i>, however, it is sensitive to cold temperatures. A second species in the genus, <i>N. fuscolinealis</i>, may also be a potential biocontrol agent (Bennett &amp; Pemberton, 2008). It feeds on <i>L. japonicum</i> in its native range but was not released and tested in the southeastern USA, because it was found to also impact a sympatric species, <i>L. palmatum</i>, which is native to the U.S. However, <i>L. palmatum</i> is not native to Europe, so <i>N. fuscolinalis</i> may have potential for trial in the European Union. It is important to note that the release of macro- (or micro- in this case) organisms as biological control agents is currently not regulated at EU level. Nevertheless national/regional laws are to be respected. Before any release of an alien species as a biological control agent an appropriate risk assessment should be made.</p>
<p><b>Scale of application</b> At what scale is the measure applied? What is the largest scale at which it has been successfully used? Please provide examples, with areas (km<sup>2</sup> or ha) if possible.</p>	<p>In initial phases of the release of a biological control agent, it is suggested that release of large populations in a few locations across a broad landscape may be most effective to establish populations of the control agent in the ecosystem (Shea &amp; Possingham, 2001). Following establishment, smaller populations can be released at multiple sites within a region.</p>

<p><b>Effectiveness of the measure</b> Is it effective in relation to its objective? Has the measure previously worked, failed?</p> <p>Please select one of the categories of effectiveness (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<table border="1"> <tr> <td><b>Effectiveness of measures</b></td> <td><i>Effective</i></td> <td></td> <td><i>Neutral</i></td> <td>x</td> <td><i>Ineffective</i></td> <td></td> </tr> </table>	<b>Effectiveness of measures</b>	<i>Effective</i>		<i>Neutral</i>	x	<i>Ineffective</i>						
<b>Effectiveness of measures</b>	<i>Effective</i>		<i>Neutral</i>	x	<i>Ineffective</i>								
	<p>Rationale: Effectiveness is rated as neutral because it is unknown at this time whether biocontrol agents who have been found to be effective for <i>L. microphyllum</i>, such as <i>Neomusotima conspurcatalis</i> and <i>N. fuscolinealis</i>, will also be effective for <i>L. japonicum</i>. It is suggested that these agents feed on all species in the <i>Lygodium</i> genus, but this has not been thoroughly tested or documented, nor is it known how these agents would adapt to the climate and ecosystems of Europe.</p>												
<p><b>Effort required</b> e.g. period of time over which measure need to be applied to have results</p>	<p>Multiple releases over multiple years are needed to establish populations of biological control agents in the wild. Additionally, host specific biological control agents require a constant supply of its host for populations to survive, so multiple release events would be necessary if <i>L. japonicum</i> were not widespread across a region.</p>												
<p><b>Resources required</b><sup>1</sup> e.g. cost, staff, equipment etc.</p>	<p>Testing and release of biological control agents in new areas can require significant upfront resources and costs. Worldwide estimates of establishing biological control programs, which includes breeding, lab testing, and field trials, range from \$160,000 to \$460,000 US\$ per year (ca. €139,000 to €400,000) for several years (Paynter et al., 2015).</p>												
<p><b>Side effects (incl. potential) – both positive and negative</b> i.e. positive or negative side effects of the measure on public health, environment including non-targeted species, etc.</p> <p>For each of the side effect types please select one of the impact categories (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<table border="1"> <tr> <td><b>Environmental effects</b></td> <td><i>Positive</i></td> <td></td> <td><i>Neutral or mixed</i></td> <td>x</td> <td><i>Negative</i></td> <td></td> </tr> </table>	<b>Environmental effects</b>	<i>Positive</i>		<i>Neutral or mixed</i>	x	<i>Negative</i>						
<b>Environmental effects</b>	<i>Positive</i>		<i>Neutral or mixed</i>	x	<i>Negative</i>								
	<table border="1"> <tr> <td><b>Social effects</b></td> <td><i>Positive</i></td> <td></td> <td><i>Neutral or mixed</i></td> <td>x</td> <td><i>Negative</i></td> <td></td> </tr> </table>	<b>Social effects</b>	<i>Positive</i>		<i>Neutral or mixed</i>	x	<i>Negative</i>						
<b>Social effects</b>	<i>Positive</i>		<i>Neutral or mixed</i>	x	<i>Negative</i>								
	<table border="1"> <tr> <td><b>Economic effects</b></td> <td><i>Positive</i></td> <td></td> <td><i>Neutral or mixed</i></td> <td>x</td> <td><i>Negative</i></td> <td></td> </tr> </table>	<b>Economic effects</b>	<i>Positive</i>		<i>Neutral or mixed</i>	x	<i>Negative</i>						
<b>Economic effects</b>	<i>Positive</i>		<i>Neutral or mixed</i>	x	<i>Negative</i>								
	<p>Rationale: Because there are currently no evaluations of potential biocontrol agents for <i>L. japonicum</i>, the potential side effects are also not known.</p>												
<p><b>Acceptability to stakeholders</b> e.g. impacted economic activities, animal welfare considerations, public perception, etc.</p> <p>Please select one of the categories of acceptability (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<table border="1"> <tr> <td><b>Acceptability to stakeholders</b></td> <td><i>Acceptable</i></td> <td></td> <td><i>Neutral or mixed</i></td> <td>x</td> <td><i>Unacceptable</i></td> <td></td> </tr> </table>	<b>Acceptability to stakeholders</b>	<i>Acceptable</i>		<i>Neutral or mixed</i>	x	<i>Unacceptable</i>						
<b>Acceptability to stakeholders</b>	<i>Acceptable</i>		<i>Neutral or mixed</i>	x	<i>Unacceptable</i>								
	<p>Rationale: Public perception of the use of biological control agents can be mixed. While there is much interest in reducing the use of chemical plant protection products, biological controls have not been widely used in the E.U. for invasive plants. Education and awareness campaigns could aid in the acceptability of this control method.</p>												



<p><b>Additional cost information</b> <sup>1</sup></p> <p>When not already included above, or in the species Risk Assessment.</p> <ul style="list-style-type: none"> <li>- implementation cost for Member States</li> <li>- the cost of inaction</li> <li>- the cost-effectiveness</li> <li>- the socio-economic aspects</li> </ul> <p>Include quantitative &amp;/or qualitative data, and case studies (incl. from countries outside the EU).</p>	<p>No information available.</p>							
<p><b>Level of confidence on the information provided</b> <sup>2</sup></p> <p>Please select one of the confidence categories along with a statement to support the category chosen. See <i>Notes</i> section at the bottom of this document.</p> <p><b>NOTE – this is not related to the effectiveness of the measure</b></p>	<p><i>Inconclusive</i></p>	<p>x</p>	<p><i>Unresolved</i></p>		<p><i>Established but incomplete</i></p>		<p><i>Well established</i></p>	
<p>Rationale:</p> <p>The scientific literature (Bennett &amp; Pemberton, 2008; Smith et al., 2014) suggests that the brown <i>Lygodium</i> moths including <i>Neomusotima conspurcatalis</i> and <i>N. fuscolinealis</i> will feed on <i>L. japonicum</i>, but they have only been tested for control with <i>L. microphyllum</i>. Sensitivity to cold may limit their use in the full range of <i>L. japonicum</i>'s possible distribution.</p>								

<p><b>Bibliography</b> <sup>3</sup></p>
<p>Aceves-Bueno, E., Adeleye, A.S., Bradley, D., Brandt, W.T., Callery, P., Feraud, M., Garner, K.L., Gentry, R., Huang, Y., McCullough, I., Pearlman, I., Sutherland, S.A., Wilkinson, W., Yang, Y., Zink, T., Anderson, S.E. &amp; Tague, C. (2015). Citizen Science as an Approach for Overcoming Insufficient Monitoring and Inadequate Stakeholder Buy-in in Adaptive Management: Criteria and Evidence. <i>Ecosystems</i>, 18, 493-506.</p> <p>Bennett, C.A. &amp; Pemberton, R.W. (2008). <i>Neomusotima fuscolinealis</i> (Lepidoptera: Pryalidae) Is an Unsuitable Biological Control Agent of <i>Lygodium japonicum</i>. <i>The Florida Entomologist</i>, 91(1), 26-29.</p>

- Bohn, K.K. & Minogue, P.J. (2010). Assessing the timing and sequence of prescribed fire and herbicide applications on the control of the invasive Japanese climbing fern in Florida's natural areas. Florida Fish and Wildlife Cons. Comm. Final Report. 9 pp.
- Bohn, K.K., Minogue, P.J. & Pieterse, E.C. (2011). Control of invasive Japanese climbing fern (*Lygodium japonicum*) and response of native ground cover during restoration of a disturbed longleaf pine ecosystem. *Ecological Restoration*, 29, 346-356.
- Bohn, K.K. & Thetford, M. (2013). *Using herbicides to control Japanese climbing fern reproduction: effects on rhizome regeneration and spore germination*. Florida Fish and Wildlife Cons. Comm. Final Report. 13 pp.
- Bohn, K.K. & Thetford, M. (2014). *Effect of herbicide type, application timing, and light intensity on Japanese climbing fern reproduction*. Florida Fish and Wildlife Cons. Comm. Final Report. 13 pp.
- Bohn, K.K., Thetford, M. & MacDonald, G. (2015). *Broad-scale herbicide applications to control Japanese climbing fern reproduction and growth across Florida's climatic zones*. Florida Fish and Wildlife Cons. Comm. Final Report. 12 pp.
- Burt, J.W., Muir, A.A., Piovia-Scott, J., Veblen, K.E., Chang, A.L., Grossman J.D. & Weiskel, H.W. (2007). Preventing horticultural introductions of invasive plants: potential efficacy of voluntary initiatives. *Biological Invasions*, 9, 909-923.
- CABI. (2017). *Lygodium japonicum*. Retrieved from [www.cabi.org/isc/datasheet/31783](http://www.cabi.org/isc/datasheet/31783). Accessed February, 21 2017.
- Caffrey, J.M., Baars, J.R., Barbour, J.H. Boets, P., Boon, P., Davenport, K., Dick, J.T.A., Early, J., Edsman, L., Gallagher, C., Gross, J., Heinimaa, P., Horrill, C., Hudin, S. & MacIscac, H.J. (2014). Tackling invasive alien species in Europe: The top 20 issues. *Mngt Biological Invasions*, 5(1), 1-20.
- Clarke, H.M. (1936). The morphology and anatomy of *Lygodium japonicum*. *American Journal of Botany*, 23(6), 405-413.
- Cleaves, D.A., Haines, T.K. & Martinez, J. (1999). *Prescribed Burning Costs: Trends and Influences in the National Forest System*. USDA Forest Service Gen. Tech. Rep. PSW-GTR-173.
- Clute, W.N. (1903). Fernwort notes - IV. *Fern Bulletin*, 11, 105-107.
- Correll, D.S. 1938. A county check-list of Florida ferns and fern allies. *American Fern Journal*, 28(2), 46-54.
- Chang, C-S., Kim, H. & Chang, K. (2014). *Provisional checklist of vascular plants for the Korea peninsula flora (KPF)*. Seoul, South Korea: Design Post, 561 pp.
- Chong, K.Y., Tan, H.T. & Corlett, R.T. (2009). *A checklist of the total vascular plant flora of Singapore: native, naturalised and cultivated species*. Singapore: Raffles Museum of Biodiversity Research, National University of Singapore, 273 pp.
- Crall, A.W., Newman, G.J., Stohlgren, T.J., Holfelder, K.A., Graham, J. & Waller, D.M. (2011). Assessing citizen science data quality: an invasive species case study. *Conservation Letters*, 4(6), 433-442.
- Crall, A.W., Renz, M., Panke, B.J., Newman, G.J., Chapin, C., Graham, J. & Barger, C. (2012). Developing cost-effective early detection networks for regional invasions. *Biological Invasions*, 14(12), 2461-2469.
- Duan, Y., Yi, D., RongRong, H., Kurihara, H., YaoLan, L. & XinSheng, Y. (2012). A new phenylpropanoid glucoside from the aerial parts of *Lygodium japonicum*. *Journal of Asian Natural Products Research*, 14(3), 286-292.
- Eisenberg, A., Amato, P.J. & Tao, D. (2009). Kam Guilzhoh nyim Guangxih di Benxtux Wenchual nyim Zihyuanc dih Gonxliix: Kam local indigenous knowledge and sustainable resource management in Guizhou and Guangxi provinces, China. *Ethnobotany Research and Applications*, 7, 67-113.
- EPPO. (2017). *Pest risk analysis for Lygodium japonicum*. Paris: EPPO. Retrieved from <https://circabc.europa.eu/>. Accessed June, 14 2018
- Fernandes, P.M., Davies, M., Ascoli, D., Fernández, C. Moreira, F., Rigolot, E., Stoof, C.R., Vega, J.A., and Molina, d. (2013). Prescribed burning in southern Europe: developing fire management in a dynamic landscape. *Frontiers Ecol Env*, 11(1), e4-e14.
- Ferriter, A., ed. (2001). *Lygodium management plan for Florida*. 1st Ed. Florida Exotic Pest Plant Council, Lygodium Task Force.
- Flora of China Editorial Committee. (2014). *Flora of China*. St. Louis, Missouri and Cambridge, Massachusetts, USA: Missouri Botanical Garden and Harvard University Herbaria. Retrieved from [http://www.efloras.org/flora\\_page.aspx?flora\\_id=2](http://www.efloras.org/flora_page.aspx?flora_id=2) Accessed February 21, 2018.
- Flory, S.L. & Lewis, J. (2009). Nonchemical methods for managing Japanese stiltgrass (*Microstegium vimineum*). *Invasive Plant Science and Management*, 2(4), 301-308.


- Gardiner, M.M., Allee, L.L., Brown, P.M.J., Losey, J.E., Roy, H.E. & Smyth, R. (2012). Lessons from lady beetles: accuracy of monitoring data from US and UK citizen-science programs. *Front Ecol Environ*, 10(9), 471-476. <https://doi.org/10.1890/110185> Accessed February 21, 2018.
- Henderson, L. (2007). Invasive, naturalized and casual alien plants in southern Africa: a summary based on the Southern African Plant Invaders Atlas (SAPIA). *Bothalia*, 37(2), 215-248
- Holck, M.H. (2008). Participatory forest monitoring: an assessment of the accuracy of simple cost-effective methods. *Biodiversity and Conservation*, 17(8), 2023-2036.
- Hosking, J.R., Conn, B.J., Lepschi, B.J. & Barker, C. (2011). Plant species first recognised as naturalised or naturalising for New South Wales in 2004 and 2005. *Cunninghamia: A Journal of Plant Ecology for Eastern Australia*, 12(1), 85-114.
- Hutchinson, J.T. & Langeland, K.A. (2006). Potential spread of *Lygodium microphyllum* spores by herbicide applicators. *Wildland Weeds* (Spring), 13-16.
- Hutchinson, J.T. and Langeland, K.A. (2007). Herbicide control of *Lygodium microphyllum*. EISS presentation. Retrieved from [www.pesticide.ifas.ufl.edu/Lygodium/.../EISS-2007-herb-control-lygodium-hutchinson.ppt](http://www.pesticide.ifas.ufl.edu/Lygodium/.../EISS-2007-herb-control-lygodium-hutchinson.ppt). Accessed August, 24 2017.
- IPPC (2017). ISMP 41 International movement of used vehicles, machinery and equipment. 12 pp. FAO, Rome. ([https://www.ippc.int/static/media/files/publication/en/2017/05/ISPM\\_41\\_2017\\_En\\_2017-05-15.pdf](https://www.ippc.int/static/media/files/publication/en/2017/05/ISPM_41_2017_En_2017-05-15.pdf))
- Keller, R.P., Lodge, D.M. & Finnoff, D.C. (2007). Risk assessment for invasive species produces net bioeconomic benefits. *PNAS*, 104 (1), 203–207.
- Kettenring, K.M. & Adams, C.R. (2011). Lessons learned from invasive plant control experiments: a systematic review and meta-analysis. *Journal of Applied Ecology*, 48, 970-979.
- Kottek, M., Grieser, J., Beck, C., Rudolf, B. & Rubel, F. (2006). World map of the Koppen-Geiger climate classification updated. *Meteorologische Zeitschrift*, 15(3), 259-263.
- Ledoux, C.B. & Martin, D.K. (2012). *Proposed BMPs for invasive plant mitigation during timber harvesting operations*. USDA For Serv Gen Tech Rep NRS-118. 16p.
- Leichty, E.R., Carmichael, B.J. & Platt, W.J. (2011). Invasion of a southeastern pine savanna by Japanese climbing fern. *Castanea*, 76, 293-299.
- Liebhold, A.M., Brockerhoff, E.G., Garrett, L.J., Parke, J.L., & Britton, K.O. (2012). Live plant imports: the major pathway for forest insect and pathogen invasions of the US. *Frontiers in Ecology and the Environment*, 10(3), 135-143.
- Lindsay, S. & Middleton, D. (2013). Ferns of Thailand, Laos and Cambodia. Retrieved from <http://rbg-web2.rbge.org.uk/thaiferns> Accessed August, 24 2017.
- Maideen, H., Latiff, A., Bidin, A., Damanhuri, A. & Jaman, R. (2004). Notes on the genus *Lygodium* Sw. (Schizaeaceae) in Malaysia. *Fern Gazette*, 17(2), 53-57.
- Miller, J.H., Manning, S.T., Enloe, S.F., United, U., Forest, F. & Southern, R.R. (2010). *A management guide for invasive plants in southern forests*. General Technical Report - Southern Research Station, USDA Forest Service, No.SRS-131:v + 120 pp.
- Minogue, P.J., Bohn, K.K., Osiecka, A. & Lauer, D.K. (2010). Japanese climbing fern (*Lygodium japonicum*) management in Florida's bottomland hardwood forests. *Invasive Plant Science and Management*, 3, 246-252.
- Orchard, A.E. & McCarthy, P.M. (1998). *Flora of Australia. Volume 48. Ferns, Gymnosperms and allied groups*. CSIRO. Melbourne, Australia.
- Paynter, Q.P., Fowler, S.V., Hayes, L. & Hill, R.L. (2015). Factors affecting the cost of weed biocontrol programs in New Zealand. *Biological Control*, 80, 119–127.
- Puri, H.S. (1970). Indian Pteridophytes used in folk remedies. *American Fern Journal*, 60(4), 137-143.
- Shea, K. and Possingham, H.P. (2001). Optimal release strategies for biological control agents: an application of stochastic dynamic programming to population management. *Journal of Applied Ecology*, 37(1), 77-86.
- Simberloff, D. (2003). How much information on population biology is needed to manage introduced species? *Cons. Biol.*, 17, 83–92.
- Simberloff, D., Martin, J-L., Genovesi, P., Maris, V., Wardle, D.A., Aronson, J., Courchamp, F., Galil B., Berthou, G., Pascal, M., Pysek, P., Sousa, R., Tabacchi, E. & Vila, M. (2013). Impacts of biological invasions: what's what and the way forward. *Trends Ecol Evol*, 28(1), 58-66.
- Singh, B., Singh, V.N., Phukan, S.J., Sinha, B.K. & Borthakur, S.K. (2012). Contribution to the pteridophytic flora of India: Nokrek Biosphere Reserve, Meghalaya. *Journal of Threatened Taxa*, 4(1), 2277-2294.
- Smith, M.C., Lake, E.C., Pratt, P.D., Boughton, A.J. & Pemberton, R.W. (2014). Current status of the biological control agent *Neomusotima conspurcatalis* (Lepidoptera: Crambidae), on *Lygodium microphyllum* (Polypodiales: Lygodiaceae) in Florida. *The Florida Entomologist*, 97(2), 817-820.

- Stocker, R.K., Miller, R.E. Jr., Black, D.W., Ferriter, A.P. & Thayer, D.D. (2008). Using Fire and Herbicide to Control *Lygodium microphyllum* and Effects on a Pine Flatwoods Plant Community in South Florid. *Natural Areas Journal*, 28(2), 144-154.
- Tsui, M.T.K and Chu, L.M. (2003). Aquatic toxicity of glyphosate-based formulations: comparison between different organisms and the effects of environmental factors. *Chemosphere*, 52, 1189-1197.
- Uddin, M.G., Mahal, M. & Pasha, M.K. (1997). Taxonomic account of genus *Lygodium* Sw. (Lygodiaceae) in Bangladesh. *Indian Fern Journal*, 14(1-2), 33-37.
- Ulrich, J. 2012. Japanese climbing fern (*Lygodium japonicum*) propagation and invasion: Effect of environmental variables on spore germination and consequences of establishment for understory plant communities and habitat quality. Master's Thesis, University of Florida. 70 pp.
- Valenta, J.T., Zeller, M., and Leslie, A. (2001). Glyphosate controls Japanese climbing fern in experimental plots (Florida). *Ecological Restoration*, 19(2), 118-119.
- van Loan, A.N. (2006). Aspects of the invasion and management of Japanese climbing fern (*Lygodium japonicum*) in Southeastern forests. Masters thesis. University of Florida.
- Verbrugge, L.N.H., Leuven, R.S.E., van Valkenburg, J.L.C.H. & van den Born, R.J.G. (2014). Evaluating stakeholder awareness and involvement in risk prevention of aquatic invasive plant species by a national code of conduct. *Aquatic Invasions*, 9(3), 369–381.
- Wunderlin, R.P. & Hansen, B.F. (2003). *Guide to the vascular plants of Florida*. Gainesville: University Press of Florida. 2<sup>nd</sup> ed. 806pp.
- Yumkham, S.D. & Singh, P.K. (2011). Less known ferns and fern-allies of Manipur with ethnobotanic uses. *Indian Journal of Traditional Knowledge*, 10(2), 287-291.
- Zeller, M., and Leslie, D. (2004). Japanese climbing fern controls in planted pine. *Wildland Weeds*, 7, 6-9.

## Notes

**1. Costs information.** The assessment of the potential costs shall describe those costs quantitatively and/or qualitatively depending on what information is available. This can include case studies from across the Union or third countries.

**2. Level of confidence<sup>1</sup>:** based on the quantity, quality and level of agreement in the evidence.



- **Well established:** comprehensive meta-analysis<sup>2</sup> or other synthesis or multiple independent studies that agree.
- **Established but incomplete:** general agreement although only a limited number of studies exist but no comprehensive synthesis and, or the studies that exist imprecisely address the question.
- **Unresolved:** multiple independent studies exist but conclusions do not agree.
- **Inconclusive:** limited evidence, recognising major knowledge gaps

**3. Citations and bibliography.** The APA formatting style for citing references in the text and in the bibliography is used.

e.g. Peer review papers will be written as follows:

In text citation: (Author & Author, Year)

In bibliography: Author, A. A., & Author, B. B. (Publication Year). Article title. *Periodical Title*, Volume(Issue), pp.-pp.

(see <http://www.waikato.ac.nz/library/study/referencing/styles/apa>)

<sup>1</sup> Assessment of confidence methodology is taken from IPBES. 2016. Guide on the production and integration of assessments from and across all scales (IPBES-4-INF-9), which is adapted from Moss and Schneider (2000).

<sup>2</sup> A statistical method for combining results from different studies which aims to identify patterns among study results, sources of disagreement among those results, or other relationships that may come to light in the context of multiple studies.