

Information on measures and related costs in relation to species considered for inclusion on the Union list: *Ehrharta calycina*

This note has been drafted by IUCN within the framework of the contract No 07.0202/2017/763436/SER/ENV.D2 “Technical and Scientific support in relation to the Implementation of Regulation 1143/2014 on Invasive Alien Species”. The information and views set out in this note do not necessarily reflect the official opinion of the Commission, or IUCN. The Commission does not guarantee the accuracy of the data included in this note. Neither the Commission nor IUCN or any person acting on the Commission’s behalf, including any authors or contributors of the notes themselves, may be held responsible for the use which may be made of the information contained therein. Reproduction is authorised provided the source is acknowledged. This document shall be cited as:

Visser, V. 2018. *Information on measures and related costs in relation to species considered for inclusion on the Union list: Ehrharta calycina*. Technical note prepared by IUCN for the European Commission.

Date of completion: 25/10/2018

Comments which could support improvement of this document are welcome. Please send your comments by e-mail to ENV-IAS@ec.europa.eu.

Species (scientific name)	<i>Ehrharta calycina</i> Sm. Pl. Ic. Ined. t. 33.
Species (common name)	Perennial veldt grass, purple veldt grass, veldt grass, common ehrharta, gewone ehrharta (Afrikaans), rooisaadgras (Afrikaans).
Author(s)	Vernon Visser, African Climate & Development Institute
Date Completed	25/10/2018
Reviewer	Courtenay A. Ray, Arizona State University

Summary

Highlight of measures that provide the most cost-effective options to prevent the introduction, achieve early detection, rapidly eradicate and manage the species, including significant gaps in information or knowledge to identify cost-effective measures.

Perennial veldt grass, *Ehrharta calycina* Sm. (Poaceae), is a perennial (sometimes annual) grass species native to South Africa and southern Namibia (Fish et al., 2015). It is a tufted rhizomatous grass that primarily reproduces from seed, and rarely from rhizomes (Chimera, 2015; Fish et al. 2015; Wittkuhn, 2010). It is a prolific producer of seed, which are primarily dispersed by wind, but also by rodents, large herbivores and water (DiTomaso et al., 2013; Newsome et al., 2008; Trunzo, 2015; Wittkuhn 2010). This species was introduced into many regions for pasture or for erosion control (Pickart, 2000; Quattrocchi, 2006). *E. calycina* is already established in the EU in Portugal (Bacelar et al., 1989; GBIF, 2018) and Spain (Bacelar et al., 1989; Charpin & Zarco, 1982; Fraga-Arguimbau, 2014; Valdés et al., 1987; Valdés, 2015). This species is invasive in Australia and California and occurs in a variety of habitats, but is most common in sandy soils (Frey, 2005; Pickart, 2000; Western Australian Herbarium, 1998).

In Australia and California, *E. calycina* is an ecosystem transformer, causing the conversion of native shrublands and woodlands into monospecific grasslands, either by preventing the growth of native plants or via a positive-feedback grass-fire cycle (Fisher et al., 2009; Pickart, 2000). In Australia, *E. calycina* has also caused a shift in phosphorous nutrient cycling in Banksia woodlands (Fisher et al., 2006).

Prevention: The most appropriate measure for preventing entry of *E. calycina* into a Member State is *a ban on keeping, importing, selling, breeding and growing* of this species. Seeds for pasture are the most likely life stage to be introduced and should be banned. *Phytosanitary measures* are likely to be ineffective for preventing entry via the principal pathways. The *establishment of containment areas around current introduction sites* should be investigated for its cost-effectiveness and suitability relative to eradication.

The use of *citizen science and resource managers' data* is a low-cost option as a **surveillance measure for early detection** with a high chance of success. Citizen science networks have been very successful in supporting early detection programs and suitable networks, databases and apps already exist in the EU.

Physical control, chemical control, grazing, and prescribed burning have all been proposed as control measures for *E. calycina* and most of these could be applied for either **rapid eradication for new introductions** or **management of widespread invasions**. However, there is a distinct lack of experimental trials on control measures for *E. calycina*, making it difficult to provide sound recommendations for the control of this species. Chemical control appears to be the measure of choice for *E. calycina* control (DiTomaso et al., 2013), but this species' long-lived seedbank (Smith et al., 1999) makes control difficult.

Prevention of intentional introductions and spread – measures for preventing the species being introduced intentionally. **This table is repeated for each of the prevention measures identified.**

<p>Measure description Provide a description of the measure, and identify its objective</p>	<p>A ban on keeping, importing, selling, breeding and growing as required under Article 7 of the IAS Regulation.</p> <p><i>Ehrharta calycina</i> has been planted as a forage plant and also for erosion control (Pickart, 2000; Quattrocchi, 2006). This species is available for commercial purposes in Australia. It is promoted as a forage grass in Australia and New Zealand, was previously tested as a forage grass in Tunisia (Greuter & Raus, 1998) and was introduced to California as a forage grass from Australia (DiTomaso et al., 2013). <i>E. calycina</i> is already established in the EU in Portugal (Bacelar et al., 1989; GBIF, 2018) and Spain (Bacelar et al., 1989; Charpin & Zarco, 1982; Fraga-Arguimbau, 2014; Valdés et al., 1987; Valdés, 2015), but it is uncertain how it was introduced there.</p> <p>In California and Australia, <i>E. calycina</i> can dominate plant communities excluding native plant species and transforming shrubland into grasslands (Fisher et al., 2009; Frey, 2005; Milberg & Lamont, 1995; Pickart, 2000). It can initiate an enhanced grass-fire cycle, promoting more frequent fires, which in turn favour this fire-adapted species at the expense of native plant species (Fisher et al., 2009; Milberg & Lamont, 1995). In eutrophic Australian Mediterranean-type environments, <i>E. calycina</i> has been shown to cause a shift in phosphorous nutrient cycling, with vegetation transformation coinciding with a shift of phosphorus from biomass to soils (Fisher et al., 2006).</p>
--	---

	<p>The objective of this measure would be to prevent the intentional re-introduction and spread of this species by banning its import, selling, and growing of the species. The fact that <i>E. calycina</i> has become invasive in Australia, California and New Zealand, where it was introduced as a forage grass, strongly supports implementing this measure.</p>																											
<p>Effectiveness of the measure Is it effective in relation to its objective? Has the measure previously worked, failed?</p>	<table border="1" data-bbox="616 290 1854 352"> <tr> <td data-bbox="616 290 893 352">Effectiveness of measures</td> <td data-bbox="902 290 1122 352"><i>Effective</i></td> <td data-bbox="1131 290 1178 352"></td> <td data-bbox="1187 290 1480 352"><i>Neutral</i></td> <td data-bbox="1489 290 1536 352">X</td> <td data-bbox="1545 290 1800 352"><i>Ineffective</i></td> <td data-bbox="1809 290 1854 352"></td> </tr> </table> <p data-bbox="616 391 1995 480"><i>Rationale:</i> No specific information is available to suggest how effective banning the keeping, importing, selling, breeding and growing <i>E. calycina</i> will be in preventing its invasion. This species occurs in the EU in Portugal and Spain. Therefore, this measure can be effective in limiting further intentional introductions of this species within the EU.</p>							Effectiveness of measures	<i>Effective</i>		<i>Neutral</i>	X	<i>Ineffective</i>															
Effectiveness of measures	<i>Effective</i>		<i>Neutral</i>	X	<i>Ineffective</i>																							
<p>Side effects (incl. potential) – both positive and negative i.e. positive or negative side effects of the measure on public health, environment including non-targeted species, etc.</p> <p>For each of the side effect types please select one of the impact categories (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<table border="1" data-bbox="616 534 1854 639"> <tr> <td data-bbox="616 534 893 568">Environmental effects</td> <td data-bbox="902 534 1122 568"><i>Positive</i></td> <td data-bbox="1131 534 1178 568"></td> <td data-bbox="1187 534 1480 568"><i>Neutral or mixed</i></td> <td data-bbox="1489 534 1536 568">X</td> <td data-bbox="1545 534 1800 568"><i>Negative</i></td> <td data-bbox="1809 534 1854 568"></td> </tr> <tr> <td data-bbox="616 571 893 604">Social effects</td> <td data-bbox="902 571 1122 604"><i>Positive</i></td> <td data-bbox="1131 571 1178 604"></td> <td data-bbox="1187 571 1480 604"><i>Neutral or mixed</i></td> <td data-bbox="1489 571 1536 604">X</td> <td data-bbox="1545 571 1800 604"><i>Negative</i></td> <td data-bbox="1809 571 1854 604"></td> </tr> <tr> <td data-bbox="616 608 893 641">Economic effects</td> <td data-bbox="902 608 1122 641"><i>Positive</i></td> <td data-bbox="1131 608 1178 641"></td> <td data-bbox="1187 608 1480 641"><i>Neutral or mixed</i></td> <td data-bbox="1489 608 1536 641">X</td> <td data-bbox="1545 608 1800 641"><i>Negative</i></td> <td data-bbox="1809 608 1854 641"></td> </tr> </table>							Environmental effects	<i>Positive</i>		<i>Neutral or mixed</i>	X	<i>Negative</i>		Social effects	<i>Positive</i>		<i>Neutral or mixed</i>	X	<i>Negative</i>		Economic effects	<i>Positive</i>		<i>Neutral or mixed</i>	X	<i>Negative</i>	
Environmental effects	<i>Positive</i>		<i>Neutral or mixed</i>	X	<i>Negative</i>																							
Social effects	<i>Positive</i>		<i>Neutral or mixed</i>	X	<i>Negative</i>																							
Economic effects	<i>Positive</i>		<i>Neutral or mixed</i>	X	<i>Negative</i>																							
<p>Acceptability to stakeholders e.g. impacted economic activities, animal welfare considerations, public perception, etc.</p> <p>Please select one of the categories of acceptability (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<table border="1" data-bbox="616 938 1854 1000"> <tr> <td data-bbox="616 938 893 1000">Acceptability to stakeholders</td> <td data-bbox="902 938 1122 1000"><i>Acceptable</i></td> <td data-bbox="1131 938 1178 1000">X</td> <td data-bbox="1187 938 1480 1000"><i>Neutral or mixed</i></td> <td data-bbox="1489 938 1536 1000"></td> <td data-bbox="1545 938 1800 1000"><i>Unacceptable</i></td> <td data-bbox="1809 938 1854 1000"></td> </tr> </table> <p data-bbox="616 1038 1995 1128"><i>Rationale:</i> Despite the use of this species for pasture in Australia, <i>E. calycina</i> does not appear to be used (or at least promoted) for this purpose in the EU. Therefore, this measure is likely to be acceptable to the agricultural community. This measure will also be acceptable due to its likelihood of reducing re-introductions and further spread of this species.</p>							Acceptability to stakeholders	<i>Acceptable</i>	X	<i>Neutral or mixed</i>		<i>Unacceptable</i>															
Acceptability to stakeholders	<i>Acceptable</i>	X	<i>Neutral or mixed</i>		<i>Unacceptable</i>																							
<p>Additional cost information¹ When not already included above, or in the species Risk Assessment.</p> <p>- the cost of inaction</p>	<p data-bbox="616 1236 1995 1299">Implementation costs for Member States will be dependent on the cost of enforcing such a ban, but figures are not readily available in the public domain. Kettunen et al. (2014) suggest that costs for this type of measure will be relatively high.</p> <p data-bbox="616 1337 1995 1399">The costs of inaction can be estimated based on costs of controlling invasions in other regions around the world. However, there are few statistics even for this. In California, the cost of hand pulling and herbicide backpack-spraying of <i>E. calycina</i> is estimated</p>																											

<p>- the cost-effectiveness - the socio-economic aspects</p> <p>Include quantitative &/or qualitative data, and case studies (incl. from countries outside the EU).</p>	<p>to cost USD 300 (ca. EUR 262) per acre, while aerial application of herbicides is estimated to cost USD 30 (ca. EUR 26) per acre (Kinkade, 2015).</p> <p>It is unknown whether this measure would be cost-effective. It has been suggested that because of the high costs of implementation and the high administrative burden, bans such as those suggested by this measure, are highly unlikely to be cost-effective (Kettunen et al., 2014). However, theoretical models suggest that there are major net positive economic benefits to preventing the entry of invasive species (Keller et al, 2007). There are, however, no known cost-benefit studies specific to <i>E. calycina</i>.</p> <p>There is no known socio-economic cost information related to the species.</p>				
<p>Level of confidence on the information provided ²</p> <p>Please select one of the confidence categories along with a statement to support the category chosen. See <i>Notes</i> section at the bottom of this document.</p> <p>NOTE – this is not related to the effectiveness of the measure</p>	<p><i>Inconclusive</i></p>	<p><i>Unresolved</i></p>	<p><i>Established but incomplete</i></p>	<p>X</p>	<p><i>Well established</i></p>
<p><i>Rationale:</i> There is a large body of literature (not specific to <i>E. calycina</i>) that supports a ban on keeping, importing, selling, breeding and growing alien species. However, there is no information specific to <i>E. calycina</i> to support this measure, either in the EU or in third countries.</p>					

Prevention of un-intentional introductions and spread – measures for preventing the species being introduced un-intentionally (cf. Article 13 of the IAS Regulation). **This table is repeated for each of the prevention measures identified.**

<p>Measure description Provide a description of the measure, and identify its objective</p>	<p>Phytosanitary inspections and measures, in particular related to the movement of hay, animals and agricultural machinery, mowers, and vehicles.</p> <p><i>Ehrharta calycina</i> has the potential to be introduced as a contaminant of hay (EPPO, 2018), as hay is imported into the EU from invaded areas (e.g. California) and grass seeds have been shown to remain viable in hay imports (Conn et al., 2010). <i>E. calycina</i> seeds may be dispersed on animal fur and in their dung (Chimera, 2015; Newsome et al., 2008), and on agricultural machinery, mowers, and vehicles (CABI, 2018). <i>E. calycina</i> may also reproduce vegetatively from rhizomes, although rarely (Chimera, 2015).</p> <p>Hay imports into the EU from areas where <i>E. calycina</i> is known to occur should not be permitted unless hay has been certified to be weed free (e.g. https://www.naisma.org/weed-free-forage). Animals from areas where <i>E. calycina</i> is known to occur should be inspected for seeds attached to their fur prior to transport.</p> <p>Agricultural machinery, mowers, and vehicles from areas where <i>E. calycina</i> is known to occur should be properly cleaned to prevent contamination of <i>E. calycina</i> seeds and rhizomes. An ISPM Standard has recently been drafted and adopted on 'International movement of used vehicles, machinery and equipment' (IPPC, 2017). This focuses on reducing the risks of transporting contaminants (soil, seeds, plant debris, pests) associated with the international movement (either traded, or for operational relocation) of vehicles, machinery and equipment (VME) that may have been used in agriculture, forestry, as well as for construction, industrial, mining, waste management and military purposes. For those VMEs that represent a contaminant risk, the phytosanitary measures recommended are detailed in the ISPM, and cover cleaning, prevention and disposal requirements. These include, cleaning using pressure washing or compressed air cleaning, chemical or temperature treatments, storing and handling VMEs that prevent contact with soil, and keeping vegetation short around storage areas or ports.</p>
<p>Scale of application At what scale is the measure applied? What is the largest scale at which it has been successfully used? Please provide examples, with areas (km² or ha) if possible.</p>	<p>This measure would need to be applied across the EU, as once hay, animals or VMEs have been imported into the EU they could be moved to high risk areas. No phytosanitary measures currently exist for this species.</p>

<p>Effectiveness of the measure Is it effective in relation to its objective? Has the measure previously worked, failed?</p> <p>Please select one of the categories of effectiveness (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<table border="1"> <tr> <td><i>Effectiveness of measures</i></td> <td><i>Effective</i></td> <td></td> <td><i>Neutral</i></td> <td></td> <td><i>Ineffective</i></td> <td>X</td> </tr> </table>	<i>Effectiveness of measures</i>	<i>Effective</i>		<i>Neutral</i>		<i>Ineffective</i>	X														
<i>Effectiveness of measures</i>	<i>Effective</i>		<i>Neutral</i>		<i>Ineffective</i>	X																
<p>Effort required e.g. period of time over which measure need to be applied to have results</p>	<p>This measure would have to be applied indefinitely due to the possibility of viable seeds and rhizomes being imported.</p>																					
<p>Resources required¹ e.g. cost, staff, equipment etc.</p>	<p>Phytosanitary inspections and measures require trained staff and equipment, and suitable disposal facilities are also required especially if implemented within the EU. Facilities required for the inspection, cleaning, and treatment of VMEs may include: surfaces that prevent contact with soil, including soil traps and wastewater management systems, temperature treatment facilities, and fumigation or chemical treatment facilities (IPPC, 2017).</p>																					
<p>Side effects (incl. potential) – both positive and negative i.e. positive or negative side effects of the measure on public health, environment including non-targeted species, etc.</p> <p>For each of the side effect types please select one of the impact categories (with an 'X'), and provide a</p>	<table border="1"> <tr> <td><i>Environmental effects</i></td> <td><i>Positive</i></td> <td></td> <td><i>Neutral or mixed</i></td> <td>X</td> <td><i>Negative</i></td> <td></td> </tr> <tr> <td><i>Social effects</i></td> <td><i>Positive</i></td> <td></td> <td><i>Neutral or mixed</i></td> <td>X</td> <td><i>Negative</i></td> <td></td> </tr> <tr> <td><i>Economic effects</i></td> <td><i>Positive</i></td> <td></td> <td><i>Neutral or mixed</i></td> <td></td> <td><i>Negative</i></td> <td>X</td> </tr> </table> <p><i>Rationale:</i> Inspections would have an economic cost to those undertaking it, which may include both government and the private sector. There would also be economic costs associated with cleaning/treating infected materials, and with any delays in the transport of high risk materials due to inspections.</p>	<i>Environmental effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>	X	<i>Negative</i>		<i>Social effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>	X	<i>Negative</i>		<i>Economic effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>		<i>Negative</i>	X
<i>Environmental effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>	X	<i>Negative</i>																	
<i>Social effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>	X	<i>Negative</i>																	
<i>Economic effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>		<i>Negative</i>	X																

<p>rationale, with supporting evidence and examples if possible.</p>						
<p>Acceptability to stakeholders e.g. impacted economic activities, animal welfare considerations, public perception, etc.</p> <p>Please select one of the categories of acceptability (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<p>Acceptability to stakeholders</p>	<p>Acceptable</p>	<p>Neutral or mixed</p>	<p>Unacceptable</p>	<p>X</p>	<p><i>Rationale:</i> Due to the probable ineffectiveness of the measure, and potential costs, it is likely this measure would be seen as unacceptable, especially by those sectors involved in the transport of high risk materials.</p>
<p>Additional cost information¹ When not already included above, or in the species Risk Assessment.</p> <ul style="list-style-type: none"> - implementation cost for Member States - the cost of inaction - the cost-effectiveness - the socio-economic aspects <p>Include quantitative &/or qualitative data, and case studies (incl. from countries outside the EU).</p>	<p>Implementation costs for Member States will be high because of the need for trained staff and long-term implementation of this measure, but figures are not readily available in the public domain.</p> <p>For costs of inaction, see above table, <i>Prevention of intentional introductions and spread</i>.</p> <p>This measure is unlikely to be cost-effective because of the high costs of implementation, but there are no studies specific to <i>E. calycina</i> to support this.</p> <p>There are no known socio-economic aspects.</p>					
<p>Level of confidence on the information provided²</p> <p>Please select one of the confidence categories along with a statement to support the category chosen. See <i>Notes</i> section at the bottom of this document.</p> <p>NOTE – this is not related to the effectiveness of the measure</p>	<p>Inconclusive</p>	<p>Unresolved</p>	<p>Established but incomplete</p>	<p>X</p>	<p>Well established</p>	<p><i>Rationale:</i> There is a reasonable amount of evidence to support the use of phytosanitary measures to prevent the unintentional introduction of <i>E. calycina</i>, but there is no evidence on the cost-effectiveness of this approach and on the probability that this species could be transported in hay or in agricultural objects.</p>

Prevention of secondary spread of the species – measures for preventing the species spreading once they have been introduced (cf. Article 13 of the IAS Regulation).

<p>Measure description Provide a description of the measure, and identify its objective</p>	<p>Establishment of containment areas around current introduction sites.</p> <p><i>Ehrharta calycina</i> has only been recorded in a few sites in the EU: in Portugal in the vicinity of Lisbon and Setúbal (Bacelar et al., 1989; GBIF, 2018), and in Spain in and near the Doñana National Park (Valdés et al., 1987; Valdés, 2015), near Cañaveral (GBIF, 2018), near Seville (Charpin & Zarco, 1982), and on Menorca (Fraga-Arguimbau, 2014).</p> <p>Natural spread rates of <i>E. calycina</i> are quite low, with seeds being primarily dispersed by wind (Mashau, 2008; Wittkuhn, 2010), although rodent dispersal has also been recorded (Trunzo, 2015). Maximum dispersal distances observed have been up to ~5 m in wind (Wittkuhn, 2010) and ~25 m by rodents (Trunzo, 2015). Apart from these natural dispersal mechanisms, it is thought that <i>E. calycina</i> may be unintentionally spread in hay, on the fur of animals and in their dung, and attached to agricultural machinery, mowers, and vehicles (see <i>Prevention of un-intentional introductions and spread</i> table above).</p> <p>Given the short natural dispersal distances of this species, and the limited number of unintentional dispersal mechanisms, populations of <i>E. calycina</i> could be effectively contained through the establishment of buffer zones (Grice et al., 2013). These buffer zones could encompass a ca. 25 m wide area (the maximum natural dispersal distance) around <i>E. calycina</i> populations in which grass-selective herbicides are applied annually (Wittkuhn, 2010). In addition, it would be necessary to prevent the removal of hay, livestock grazing and the use of agricultural machinery, mowers and vehicles in these zones (or have these inspected for <i>E. calycina</i> seeds).</p> <p>A cost-benefit analysis would need to be performed to determine whether this measure should be applied instead of those detailed in the table, <i>Rapid eradication for new introductions</i> (see below).</p>
<p>Scale of application At what scale is the measure applied? What is the largest scale at which it has been successfully used? Please provide examples, with areas (km² or ha) if possible.</p>	<p>This measure would only need to be applied around the known populations of <i>E. calycina</i> in Portugal and Spain. This measure has only been recommended (Wittkuhn, 2010), but has not been applied before.</p>

<p>Effectiveness of the measure Is it effective in relation to its objective? Has the measure previously worked, failed?</p> <p>Please select one of the categories of effectiveness (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<table border="1"> <tr> <td><i>Effectiveness of measures</i></td> <td><i>Effective</i></td> <td></td> <td><i>Neutral</i></td> <td>X</td> <td><i>Ineffective</i></td> <td></td> </tr> </table>	<i>Effectiveness of measures</i>	<i>Effective</i>		<i>Neutral</i>	X	<i>Ineffective</i>															
<i>Effectiveness of measures</i>	<i>Effective</i>		<i>Neutral</i>	X	<i>Ineffective</i>																	
<p>Effort required e.g. period of time over which measure need to be applied to have results</p>	<p><i>Rationale:</i> It is unknown whether this measure will be effective as it has not been applied to <i>E. calycina</i> before. Invasive plant species containment programs however can be successful given sufficient resources, relatively small invaded areas, and high detectability of the invasive species (Moore et al., 2011).</p> <p>This measure would need to be implemented indefinitely to prevent the secondary spread of this species. As this species flowers in spring and summer (University of California, Berkeley, 2018; Western Australian Herbarium, 1998), preventing the removal of hay, livestock grazing and the use of agricultural machinery, mowers and vehicles in these zones would perhaps not be necessary during winter and early spring when there are unlikely to be many seeds on the plants.</p>																					
<p>Resources required¹ e.g. cost, staff, equipment etc.</p>	<p>The cost of implementing this measure is unknown. This measure would require trained staff for (1) plant identification, (2) delimitation of the containment zones, (3) herbicide application, (4) management of potential vectors of <i>E. calycina</i> seed (e.g. livestock and vehicles) from late spring to autumn when seeds are likely to be on the plants.</p> <p>In addition, (1) would require identification guides, and (3) would require all the necessary equipment for herbicide application (see <i>Management</i> table).</p>																					
<p>Side effects (incl. potential) – both positive and negative i.e. positive or negative side effects of the measure on public health, environment including non-targeted species, etc.</p>	<table border="1"> <tr> <td><i>Environmental effects</i></td> <td><i>Positive</i></td> <td></td> <td><i>Neutral or mixed</i></td> <td>X</td> <td><i>Negative</i></td> <td></td> </tr> <tr> <td><i>Social effects</i></td> <td><i>Positive</i></td> <td></td> <td><i>Neutral or mixed</i></td> <td></td> <td><i>Negative</i></td> <td>X</td> </tr> <tr> <td><i>Economic effects</i></td> <td><i>Positive</i></td> <td></td> <td><i>Neutral or mixed</i></td> <td></td> <td><i>Negative</i></td> <td>X</td> </tr> </table> <p><i>Rationale:</i> This measure will have negative social and economic side effects in that it could limit certain farming activities in the invaded areas (e.g. grazing of livestock at certain times of the year).</p>	<i>Environmental effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>	X	<i>Negative</i>		<i>Social effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>		<i>Negative</i>	X	<i>Economic effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>		<i>Negative</i>	X
<i>Environmental effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>	X	<i>Negative</i>																	
<i>Social effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>		<i>Negative</i>	X																
<i>Economic effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>		<i>Negative</i>	X																
<p>Acceptability to stakeholders e.g. impacted economic activities, animal welfare considerations, public perception, etc.</p> <p>Please select one of the categories of acceptability (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<table border="1"> <tr> <td><i>Acceptability to stakeholders</i></td> <td><i>Acceptable</i></td> <td></td> <td><i>Neutral or mixed</i></td> <td>X</td> <td><i>Unacceptable</i></td> <td></td> </tr> </table> <p><i>Rationale:</i> This measure may be seen as too costly and impractical to implement. Farmers, especially from within the containment zones, may perceive it negatively if it limits their farming activities and causes production losses. However, farmers, especially from outside the containment zones, may also perceive it positively if <i>E. calycina</i> is perceived as a poor pasture species that is outcompeting more nutritious pasture species.</p>	<i>Acceptability to stakeholders</i>	<i>Acceptable</i>		<i>Neutral or mixed</i>	X	<i>Unacceptable</i>															
<i>Acceptability to stakeholders</i>	<i>Acceptable</i>		<i>Neutral or mixed</i>	X	<i>Unacceptable</i>																	

<p>Additional cost information ¹</p> <p>When not already included above, or in the species Risk Assessment.</p> <ul style="list-style-type: none"> - implementation cost for Member States - the cost of inaction - the cost-effectiveness - the socio-economic aspects <p>Include quantitative &/or qualitative data, and case studies (incl. from countries outside the EU).</p>	<p>Implementation costs for such a containment measure are not readily available for <i>E. calycina</i>. Costs could be approximated based on the extent of the invaded areas, the labour hours needed to implement this measure, and herbicide costs.</p> <p>Costs of inaction associated with this measure are likely to be similar to those detailed in the <i>Prevention of intentional introductions and spread</i> table above.</p> <p>To determine the cost effectiveness of this measure, one would need to conduct a formal analysis along the lines of Moore et al. (2011).</p> <p>Socio-economic aspects include the potential loss of income to farmers in the invaded areas due to the implementation of this measure.</p>								
<p>Level of confidence on the information provided ²</p> <p>Please select one of the confidence categories along with a statement to support the category chosen. See <i>Notes</i> section at the bottom of this document.</p> <p>NOTE – this is not related to the effectiveness of the measure</p>	<i>Inconclusive</i>		<i>Unresolved</i>	<i>X</i>	<i>Established but incomplete</i>		<i>Well established</i>		
<p><i>Rationale:</i> While there is a large scientific literature on the containment of invasive species, the use of this strategy for <i>E. calycina</i> has never been investigated. Therefore, it is not known whether this strategy would be effective, and the data to conduct such an analysis would require a considerable investment of time and effort to acquire.</p>									

<p>Surveillance measures to support early detection - Measures to run an effective surveillance system for achieving an early detection of a new occurrence (cf. Article 16). This section assumes that the species is not currently present in a Member State, or part of a Member State's territory. This table is repeated for each of the early detection measures identified.</p>	
<p>Measure description</p> <p>Provide a description of the measure, and identify its objective</p>	<p>Use citizen science data for early detection, supported by awareness campaigns</p> <p>The objectives of this measure would be (1) to promote <i>Ehrharta calycina</i> as a target for identification to invasive species citizen science platforms, and (2) to provide citizen scientists, farmers and environmental managers with the knowledge to identify this species and means to notify the relevant authorities, thereby supporting its early detection. Citizen science programs need good quality assessment of the data collected, well designed and standardised methods of data collection, an explicit goal or hypothesis</p>

	<p>(i.e., in this case, the early detection of <i>E. calycina</i>), and feedback to participants on their contributions as a reward for their participation (Silvertown, 2009).</p> <p>Citizen science locality data has been shown to be very useful for the early detection of invasive species (Gallo et al. 2011; Maistrello et al., 2016). Numerous such databases currently exist, including EASIN (https://easin.jrc.ec.europa.eu/), which is the official EU platform for reporting alien species occurrences along with the accompanying smart phone application. There are also a number of other European and national IAS awareness and citizen science IAS monitoring programs into which <i>E. calycina</i> could be incorporated, which are important resources for environmental agencies, environmental managers and decision makers (Genovesi et al., 2010).</p> <p>Easy-to-use identification guides for <i>E. calycina</i> (e.g. University of California, Berkeley, 2018; Western Australian Herbarium, 1998) should be developed specifically for Europe to assist with identification of this species and made available online on citizen science platforms, and distributed to key interest groups such as farming and environmental management organisations.</p>					
<p>Scale of application At what scale is the measure applied? What is the largest scale at which it has been successfully used? Please provide examples, with areas (km² or ha) if possible.</p>	<p>This measure would need to be applied across the EU, but countries/regions with known populations, as well as those with high climatic suitability, should be prioritised (see EPPO, 2018 for details).</p>					
<p>Effectiveness of the measure Is it effective in relation to its objective? Has the measure previously worked, failed? Please select one of the categories of effectiveness (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<table border="1" data-bbox="618 855 1854 912"> <tr> <td data-bbox="618 855 896 912"><i>Effectiveness of measures</i></td> <td data-bbox="904 855 1120 912"><i>Effective</i></td> <td data-bbox="1128 855 1178 912">X</td> <td data-bbox="1187 855 1478 912"><i>Neutral</i></td> <td data-bbox="1487 855 1800 912"><i>Ineffective</i></td> </tr> </table> <p><i>Rationale:</i> This measure has the potential to be very effective. Citizen science locality data has been shown to increase the likelihood of success of arthropod eradication programs, and the authors suggest that awareness campaigns were pivotal in this regard (Tobin et al., 2014).</p>	<i>Effectiveness of measures</i>	<i>Effective</i>	X	<i>Neutral</i>	<i>Ineffective</i>
<i>Effectiveness of measures</i>	<i>Effective</i>	X	<i>Neutral</i>	<i>Ineffective</i>		
<p>Effort required e.g. period of time over which measure need to be applied to have results</p>	<p>This measure would need to be supported for the long term.</p>					
<p>Resources required¹ e.g. cost, staff, equipment etc.</p>	<p>This measure will require a well-designed and supported citizen science platform, including expertise to validate records. The use of EASIN and other established national invasive alien species platforms for this purpose is possible, but the promotion of recording <i>E. calycina</i> will be required. Identification guides, and engagement with key sectors of civil society to increase awareness will also be needed.</p>					

<p>Side effects (incl. potential) – both positive and negative i.e. positive or negative side effects of the measure on public health, environment including non-targeted species, etc.</p> <p>For each of the side effect types please select one of the impact categories (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<table border="1" data-bbox="611 188 1854 292"> <tr> <td><i>Environmental effects</i></td> <td><i>Positive</i></td> <td>X</td> <td><i>Neutral or mixed</i></td> <td></td> <td><i>Negative</i></td> <td></td> </tr> <tr> <td><i>Social effects</i></td> <td><i>Positive</i></td> <td>X</td> <td><i>Neutral or mixed</i></td> <td></td> <td><i>Negative</i></td> <td></td> </tr> <tr> <td><i>Economic effects</i></td> <td><i>Positive</i></td> <td></td> <td><i>Neutral or mixed</i></td> <td>X</td> <td><i>Negative</i></td> <td></td> </tr> </table> <p><i>Rationale:</i> Awareness of additional invasive alien species, and potentially their reporting, could be a side-effect of this measure.</p> <p>While citizen science projects cost money to develop and maintain, the return on investment is estimated to be substantial and much higher than the input costs (Tulloch et al., 2013).</p>	<i>Environmental effects</i>	<i>Positive</i>	X	<i>Neutral or mixed</i>		<i>Negative</i>		<i>Social effects</i>	<i>Positive</i>	X	<i>Neutral or mixed</i>		<i>Negative</i>		<i>Economic effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>	X	<i>Negative</i>	
<i>Environmental effects</i>	<i>Positive</i>	X	<i>Neutral or mixed</i>		<i>Negative</i>																	
<i>Social effects</i>	<i>Positive</i>	X	<i>Neutral or mixed</i>		<i>Negative</i>																	
<i>Economic effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>	X	<i>Negative</i>																	
<p>Acceptability to stakeholders e.g. impacted economic activities, animal welfare considerations, public perception, etc.</p> <p>Please select one of the categories of acceptability (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<table border="1" data-bbox="611 595 1854 659"> <tr> <td><i>Acceptability to stakeholders</i></td> <td><i>Acceptable</i></td> <td>X</td> <td><i>Neutral or mixed</i></td> <td></td> <td><i>Unacceptable</i></td> <td></td> </tr> </table> <p><i>Rationale:</i> The public is likely to perceive a citizen science measure favourably. Participants in citizen science programs report having an increased appreciation of the natural world, and greater scientific knowledge, among other benefits (Toomey & Domroese, 2013). However, it has been noted that participation in citizen science programs is often limited to wealthier segments of society (Toomey & Domroese, 2013). Environmental managers would likely welcome information on (potentially) invasive species.</p>	<i>Acceptability to stakeholders</i>	<i>Acceptable</i>	X	<i>Neutral or mixed</i>		<i>Unacceptable</i>															
<i>Acceptability to stakeholders</i>	<i>Acceptable</i>	X	<i>Neutral or mixed</i>		<i>Unacceptable</i>																	
<p>Additional cost information¹ When not already included above, or in the species Risk Assessment.</p> <ul style="list-style-type: none"> - implementation cost for Member States - the cost of inaction - the cost-effectiveness - the socio-economic aspects <p>Include quantitative &/or qualitative data, and case studies (incl. from countries outside the EU).</p>	<p>Implementation costs of setting up and running citizen science databases and awareness raising programs are large (Genovesi et al., 2010), but databases and programs like these are already running and funded by the EU (e.g. EASIN). Therefore, additional costs for promoting the collection of <i>E. calycina</i> records, and raising awareness of this species, are likely to be minimal.</p> <p>Costs of inaction associated with this measure are likely to be similar to those detailed in the <i>Prevention of intentional introductions and spread</i> table.</p> <p>Cost-effectiveness of citizen science programs is well established and justified elsewhere (e.g. Gallo et al., 2011; Genovesi et al., 2010; Maistrello et al., 2016).</p> <p>There are no known additional socio-economic aspects to consider.</p>																					
<p>Level of confidence on the information provided²</p>	<table border="1" data-bbox="611 1321 1899 1385"> <tr> <td><i>Inconclusive</i></td> <td></td> <td><i>Unresolved</i></td> <td></td> <td><i>Established but incomplete</i></td> <td></td> <td><i>Well established</i></td> <td>X</td> </tr> </table>	<i>Inconclusive</i>		<i>Unresolved</i>		<i>Established but incomplete</i>		<i>Well established</i>	X													
<i>Inconclusive</i>		<i>Unresolved</i>		<i>Established but incomplete</i>		<i>Well established</i>	X															

<p>Please select one of the confidence categories along with a statement to support the category chosen. See <i>Notes</i> section at the bottom of this document.</p> <p>NOTE – this is not related to the effectiveness of the measure</p>	<p><i>Rationale:</i> There is considerable evidence to support the use of citizen science for early detection of invasive species.</p>
--	--

<p>Rapid eradication for new introductions - Measures to achieve eradication <u>at an early stage of invasion</u>, after an early detection of a new occurrence (cf. Article 17). This section assumes that the species is not currently present in a Member State, or part of a Member State's territory. This table is repeated for each of the eradication measures identified.</p>	
<p>Measure description Provide a description of the measure, and identify its objective</p>	<p>Physical control</p> <p>Physical control, chemical control, grazing, and prescribed burning have all been proposed as control measures for <i>Ehrharta calycina</i>. Grazing is possibly the only control measure that will not work for rapid eradications due to the intensity and duration of grazing required for the elimination of <i>E. calycina</i>. However, there are no specific measures recommended in the literature for rapid eradication of new introductions of <i>E. calycina</i>. Only physical control is likely to be practical and cost-effective for small invasions.</p> <p>With small invasions, plants can be cut out (or pulled) from the ground while ensuring that the crown is removed (Western Australian Herbarium, 1998; Ray et al, 2018.), but this can also stimulate seed germination (DiTomaso et al., 2013). Plant densities can be reduced if seedlings are repeatedly removed over a number of years, but the length of seed viability for <i>E. calycina</i> is uncertain, with reports of 5 years viability (Western Australian Herbarium, 1998), but possibly greater than 45 years (Smith et al., 1999).</p>
<p>Scale of application At what scale is the measure applied? What is the largest scale at which it has been successfully used? Please provide examples, with areas (km² or ha) if possible.</p>	<p>There are no specific guidelines for the scale at which this measure can be used, but because manual removal is a labour-intensive method of control, this measure is only cost-effective for small invasions (Western Australian Herbarium, 1998). For the related <i>E. erecta</i>, Ray et al (2018) tested hand pulling in experimental treatment plots of 4 m² at 12 sites in Santa Cruz, California.</p>

<p>Effectiveness of the measure Is it effective in relation to its objective? Has the measure previously worked, failed?</p> <p>Please select one of the categories of effectiveness (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<table border="1"> <tr> <td>Effectiveness of measures</td> <td><i>Effective</i></td> <td></td> <td><i>Neutral</i></td> <td>X</td> <td><i>Ineffective</i></td> <td></td> </tr> </table>	Effectiveness of measures	<i>Effective</i>		<i>Neutral</i>	X	<i>Ineffective</i>		<p><i>Rationale:</i> Physical control can be effective for managing small invasions if the crown is removed when plants are dug out, but this can stimulate seed germination and also requires numerous follow-up treatments. For the related <i>E. erecta</i>, Ray et al (2018) conclude that hand pulling is an effective measure to reduce species cover (by 76%) for up to 2 years, but that multiple treatments would be needed to eradicate the species at local scales.</p>														
Effectiveness of measures	<i>Effective</i>		<i>Neutral</i>	X	<i>Ineffective</i>																		
<p>Effort required e.g. period of time over which measure need to be applied to have results</p>	<p>There are no documented successful eradications of <i>E. calycina</i>, therefore the period of time over which physical control needs to be used is uncertain. <i>E. calycina</i> is a prolific seed producer (Smith et al., 1999) and seed viability of <i>E. calycina</i> possibly exceeds 45 years (Smith et al., 1999), but recommendations for this species' control suggest that populations of <i>E. calycina</i> can be reduced to very low numbers after just two years of treatment (DiTomaso et al., 2013).</p> <p>For the related <i>E. erecta</i>, Ray et al (2018) found that within a two-hour window, 21 volunteers could remove the species from 32 m². They also estimated that each volunteer pulled at a rate of approximately 0.75 to 1 m²·h⁻¹ (32 m²/21 people/2 h, including transit time between plots).</p>																						
<p>Resources required¹ e.g. cost, staff, equipment etc.</p>	<p>Physical control only requires trained staff or volunteers (for accurate species identification) and tools (spades, trowels or clippers). Ray et al. (2018) used volunteers, but estimated that if they had to pay labour costs it would be more expensive than herbicide application (see herbicide application table below).</p>																						
<p>Side effects (incl. potential) – both positive and negative i.e. positive or negative side effects of the measure on public health, environment including non-targeted species, etc.</p> <p>For each of the side effect types please select one of the impact categories (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<table border="1"> <tr> <td>Environmental effects</td> <td><i>Positive</i></td> <td></td> <td><i>Neutral or mixed</i></td> <td>X</td> <td><i>Negative</i></td> <td></td> </tr> <tr> <td>Social effects</td> <td><i>Positive</i></td> <td></td> <td><i>Neutral or mixed</i></td> <td>X</td> <td><i>Negative</i></td> <td></td> </tr> <tr> <td>Economic effects</td> <td><i>Positive</i></td> <td></td> <td><i>Neutral or mixed</i></td> <td>X</td> <td><i>Negative</i></td> <td></td> </tr> </table>	Environmental effects	<i>Positive</i>		<i>Neutral or mixed</i>	X	<i>Negative</i>		Social effects	<i>Positive</i>		<i>Neutral or mixed</i>	X	<i>Negative</i>		Economic effects	<i>Positive</i>		<i>Neutral or mixed</i>	X	<i>Negative</i>		<p><i>Rationale:</i> Hand pulling will have lower non-target impacts than other methods of control, such as herbicide application.</p>
Environmental effects	<i>Positive</i>		<i>Neutral or mixed</i>	X	<i>Negative</i>																		
Social effects	<i>Positive</i>		<i>Neutral or mixed</i>	X	<i>Negative</i>																		
Economic effects	<i>Positive</i>		<i>Neutral or mixed</i>	X	<i>Negative</i>																		
<p>Acceptability to stakeholders e.g. impacted economic activities, animal welfare considerations, public perception, etc.</p>	<table border="1"> <tr> <td>Acceptability to stakeholders</td> <td><i>Acceptable</i></td> <td></td> <td><i>Neutral or mixed</i></td> <td>X</td> <td><i>Unacceptable</i></td> <td></td> </tr> </table>	Acceptability to stakeholders	<i>Acceptable</i>		<i>Neutral or mixed</i>	X	<i>Unacceptable</i>																
Acceptability to stakeholders	<i>Acceptable</i>		<i>Neutral or mixed</i>	X	<i>Unacceptable</i>																		

<p>Please select one of the categories of acceptability (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<p><i>Rationale:</i> The low effectiveness and labour intensiveness of this measure is likely to reduce its acceptability to invasive species managers as a control method. However, it may be seen by the public as a more acceptable measure than others, such as herbicide application.</p>							
<p>Additional cost information ¹ When not already included above, or in the species Risk Assessment.</p> <ul style="list-style-type: none"> - implementation cost for Member States - the cost of inaction - the cost-effectiveness - the socio-economic aspects <p>Include quantitative &/or qualitative data, and case studies (incl. from countries outside the EU).</p>	<p>Implementation costs are unknown.</p> <p>Costs of inaction associated with this measure are likely to be similar to those detailed in the <i>Prevention of intentional introductions and spread</i> table.</p> <p>The cost-effectiveness of physical control for rapid eradication of this species is unknown, but likely low because of the high labour costs.</p> <p>There are no known socio-economic aspects.</p>							
<p>Level of confidence on the information provided ²</p> <p>Please select one of the confidence categories along with a statement to support the category chosen. See <i>Notes</i> section at the bottom of this document.</p> <p>NOTE – this is not related to the effectiveness of the measure</p>	<p><i>Inconclusive</i></p>	<p>X</p>	<p><i>Unresolved</i></p>		<p><i>Established but incomplete</i></p>		<p><i>Well established</i></p>	
<p><i>Rationale:</i> There is insufficient evidence to support the use of physical control for rapid eradication of this species. There is also no readily available information on the costs of using such an approach, and on the scales at which it is practical and cost-effective.</p>								

Management - Measures to achieve management of the species once it has become widely spread within a Member State, or part of a Member State's territory. (cf. Article 19), i.e. **not** at an early stage of invasion (see Rapid eradication table above). These measures can be aimed at eradication, population control or containment of a population of the species. **This table is repeated for each of the management measures identified.**

<p>Measure description Provide a description of the measure, and identify its objective</p>	<p>Chemical control</p> <p>A number of herbicides have been recommended to control <i>Ehrharta calycina</i>. Fluazifop, glyphosate and imazapyr have all been recommended for broadcast foliar or spot treatments (DiTomaso et al., 2013; Frey, 2005; Pickart, 2000; Western Australian Herbarium, 1998), but there do not appear to be any published experimental trials for the use of these chemicals on <i>E. calycina</i>. Arrow® 2EC and Poast® have recently been trialed for aerial spraying of <i>E. calycina</i>, but no results of this study seem to be available (USFWS, 2014). Glyphosate has been suggested as the most effective chemical to use for <i>E. calycina</i> control (DiTomaso et al., 2013).</p> <p>Chemical control has also been recommended in conjunction with mowing and burning. Frey (2005) recommends mowing this species prior to the application of fluazifop, but only for plants that are not seeding. Chemical control has been recommended for use on plants 4 to 6 weeks post germination after unplanned fires (Western Australian Herbarium, 1998)</p> <p>The objective of chemical control is to mitigate the impacts and control populations of this species. Chemical control could be used to help contain invasions of this species, but it is uncertain if this method will be effective for eradication in the long term. It is important to note that EU/national/local legislation on the use of plant protection products and biocides needs to be respected.</p>					
<p>Scale of application At what scale is the measure applied? What is the largest scale at which it has been successfully used? Please provide examples, with areas (km² or ha) if possible.</p>	<p>There are no specific guidelines for the scale at which chemical control can be used. Aerial spraying of Arrow® 2EC and Poast® has been trialed on experimental plots covering 40 acres (USFWS, 2014). Presumably large areas can be treated using chemical control as recommended herbicide concentrations are provided by DiTomaso et al. (2013) in pints per acre.</p>					
<p>Effectiveness of the measure Is it effective in relation to its objective? Has the measure previously worked, failed?</p> <p>Please select one of the categories of effectiveness (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<table border="1" data-bbox="611 954 1854 1018"> <tr> <td data-bbox="611 954 891 1018">Effectiveness of measures</td> <td data-bbox="891 954 1120 1018"><i>Effective</i></td> <td data-bbox="1120 954 1478 1018"><i>Neutral</i></td> <td data-bbox="1478 954 1536 1018">X</td> <td data-bbox="1536 954 1803 1018"><i>Ineffective</i></td> </tr> </table> <p><i>Rationale:</i> Glyphosate is likely the most effective control method (DiTomaso et al., 2013) and has been quoted as being able to kill (all) plants after just one application (Pickart 2000). However, as there are no published experimental chemical control trials, it is difficult to assess effectiveness of this measure. For the related <i>E. erecta</i>, Ray et al (2018) found that spraying 4m² test plots with 2.5% glyphosate solution, with a follow-up treatment using a 3% to 4% glyphosate solution, was effective to substantially reduce its cover (by 59%) for up to 2 years.</p>	Effectiveness of measures	<i>Effective</i>	<i>Neutral</i>	X	<i>Ineffective</i>
Effectiveness of measures	<i>Effective</i>	<i>Neutral</i>	X	<i>Ineffective</i>		
<p>Effort required e.g. period of time over which measure need to be applied to have results</p>	<p>It has been recommended that chemical control be applied on actively growing plants, typically in spring just after germination until just before the plants seed (DiTomaso et al., 2013; Western Australian Herbarium, 1998).</p> <p>Because <i>E. calycina</i> has such long seed viability (possibly greater than 45 years; Smith et al., 1999), follow-up treatments may be needed for many years.</p>					

<p>Resources required¹ e.g. cost, staff, equipment etc.</p>	<p>Herbicide application requires trained staff, equipment (e.g. backpack sprayers, ropewicks, spray wands, or in the case of aerial application, a helicopter fitted with a suitable spray device), herbicides and surfactants.</p> <p>The only cost estimates of chemical control available suggest the cost of hand pulling and herbicide backpack-spraying of <i>E. calycina</i> to be about USD 300 (ca. EUR 263) per acre, while aerial application of herbicides is estimated to cost USD 30 (ca. EUR 26) per acre (Kinkade, 2015).</p> <p>For herbicide control of the related <i>E. erecta</i>, Ray et al (2018) detail labour costs at approximately USD 39 h⁻¹ for a total of USD 156 (ca. EUR 137) for 4 h (2 h per application). Approximately 7.5 L of glyphosate solution were needed to spray twelve 4 m² plots. Averaging the percentage of glyphosate used in the two applications to 3%, a total of 0.47 L of glyphosate was used, for a cost of USD 23.52 (ca. EUR 21).</p>						
<p>Side effects (incl. potential) – both positive and negative i.e. positive or negative side effects of the measure on public health, environment including non-targeted species, etc.</p> <p>For each of the side effect types please select one of the impact categories (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<p>Environmental effects</p>	<p>Positive</p>	<p>Neutral or mixed</p>	<p>X</p>	<p>Negative</p>	<p>X</p>	
<p>Acceptability to stakeholders e.g. impacted economic activities, animal welfare considerations, public perception, etc.</p> <p>Please select one of the categories of acceptability (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<p>Acceptability to stakeholders</p>	<p>Acceptable</p>	<p>Neutral or mixed</p>	<p>X</p>	<p>Unacceptable</p>	<p></p>	
<p>Additional cost information¹ When not already included above, or in the species Risk Assessment.</p>	<p>The best estimates of implementation costs can be found in the <i>Resources required</i> section above.</p> <p>Costs of inaction associated with this measure are likely to be similar to those detailed in the <i>Prevention of intentional introductions and spread</i> table.</p>						

<ul style="list-style-type: none"> - implementation cost for Member States - the cost of inaction - the cost-effectiveness - the socio-economic aspects <p>Include quantitative &/or qualitative data, and case studies (incl. from countries outside the EU).</p>	<p>The cost-effectiveness of chemical control for management of this species is unknown, but will probably be higher than for other methods of control.</p> <p>There are no known socio-economic aspects.</p>							
<p>Level of confidence on the information provided ²</p> <p>Please select one of the confidence categories along with a statement to support the category chosen. See <i>Notes</i> section at the bottom of this document.</p> <p>NOTE – this is not related to the effectiveness of the measure</p>	<i>Inconclusive</i>	<input checked="" type="checkbox"/>	<i>Unresolved</i>	<input type="checkbox"/>	<i>Established but incomplete</i>	<input type="checkbox"/>	<i>Well established</i>	<input type="checkbox"/>
<p><i>Rationale:</i> Chemical control is recommended for controlling <i>E. calycina</i>, but there are no published experimental trials for the use of herbicides on <i>E. calycina</i>. The available guidelines for chemical control of <i>E. calycina</i> suggest a diversity of chemicals, application procedures and timing making it difficult to determine how best to implement chemical control. There are also almost no figures available on the costs of chemical control of this species, and no studies on its cost effectiveness.</p>								

<p>Management - Measures to achieve management of the species once it has become widely spread within a Member State, or part of a Member State's territory. (cf. Article 19), i.e. not at an early stage of invasion (see Rapid eradication table above). These measures can be aimed at eradication, population control or containment of a population of the species. This table is repeated for each of the management measures identified.</p>	
<p>Measure description</p> <p>Provide a description of the measure, and identify its objective</p>	<p>Grazing</p> <p><i>Ehrharta calycina</i> appears unable to survive heavy grazing, particularly when it is flowering (DiTomaso et al., 2013; Rossiter, 1947). However, <i>E. calycina</i> was found to be able to survive the removal of up to 80% of its aboveground biomass (Van der Westhuizen and Joubert, 1983), suggesting it can tolerate a certain degree of grazing, a finding similar to other studies (DiTomaso et al., 2013; Rossiter, 1947).</p> <p>The objective of this measure is to mitigate the impacts and control populations of this species.</p>
<p>Scale of application</p> <p>At what scale is the measure applied? What is the largest scale at which it</p>	<p>There are no specific recommendations for the scale at which grazing can be used.</p>

has been successfully used? Please provide examples, with areas (km ² or ha) if possible.							
<p>Effectiveness of the measure Is it effective in relation to its objective? Has the measure previously worked, failed?</p> <p>Please select one of the categories of effectiveness (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	Effectiveness of measures	<i>Effective</i>		<i>Neutral</i>		<i>Ineffective</i>	X
<p><i>Rationale:</i> Grazing is only infrequently mentioned as a control method for <i>E. calycina</i>, and it has been suggested that this species is somewhat tolerant of grazing (DiTomaso et al., 2013; Van der Westhuizen and Joubert, 1983). Seeds of this species may also be dispersed in animal dung (Newsome et al., 2008).</p>							
<p>Effort required e.g. period of time over which measure need to be applied to have results</p>	<p>This species appears more sensitive to grazing during its flowering period (Rossiter, 1947), which is during spring to mid-summer (University of California, Berkeley, 2018; Western Australian Herbarium, 1998).</p>						
<p>Resources required¹ e.g. cost, staff, equipment etc.</p>	<p>This measure would require domestic livestock (sheep have been suggested) and staff to manage the animals. The cost of implementation is unknown.</p>						
<p>Side effects (incl. potential) – both positive and negative i.e. positive or negative side effects of the measure on public health, environment including non-targeted species, etc.</p> <p>For each of the side effect types please select one of the impact categories (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	Environmental effects	<i>Positive</i>		<i>Neutral or mixed</i>	X	<i>Negative</i>	
		<i>Positive</i>		<i>Neutral or mixed</i>	X	<i>Negative</i>	
		<i>Positive</i>		<i>Neutral or mixed</i>	X	<i>Negative</i>	
<p><i>Rationale:</i> Grazing may not be suitable or desirable in all locations where <i>E. calycina</i> occurs (e.g. certain conservation areas) and could have negative environmental effects.</p>							
<p>Acceptability to stakeholders e.g. impacted economic activities, animal welfare considerations, public perception, etc.</p>	Acceptability to stakeholders	<i>Acceptable</i>		<i>Neutral or mixed</i>	X	<i>Unacceptable</i>	

<p>Please select one of the categories of acceptability (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>					
<p>Additional cost information ¹ When not already included above, or in the species Risk Assessment.</p> <ul style="list-style-type: none"> - implementation cost for Member States - the cost of inaction - the cost-effectiveness - the socio-economic aspects <p>Include quantitative &/or qualitative data, and case studies (incl. from countries outside the EU).</p>	<p>Implementation costs are unknown.</p> <p>Costs of inaction associated with this measure are likely similar to those detailed in the <i>Prevention of intentional introductions and spread</i> table.</p> <p>The cost-effectiveness of grazing for management of this species is unknown, but is likely low because of the low effectiveness of this measure.</p> <p>There are no known socio-economic aspects.</p>				
<p>Level of confidence on the information provided ²</p> <p>Please select one of the confidence categories along with a statement to support the category chosen. See <i>Notes</i> section at the bottom of this document.</p> <p>NOTE – this is not related to the effectiveness of the measure</p>	<i>Inconclusive</i>	X	<i>Unresolved</i>	<i>Established but incomplete</i>	<i>Well established</i>
<p><i>Rationale:</i> Since there is little mention of the use of grazing for <i>E. calycina</i> control, it is difficult to determine whether there is agreement in the effectiveness of this measure. More studies are needed to determine its cost-effectiveness and implementation.</p>					

Management - Measures to achieve management of the species once it has become widely spread within a Member State, or part of a Member State's territory. (cf. Article 19), i.e. **not** at an early stage of invasion (see Rapid eradication table above). These measures can be aimed at eradication, population control or containment of a population of the species. **This table is repeated for each of the management measures identified.**

<p>Measure description Provide a description of the measure, and identify its objective</p>	<p>Prescribed burning</p>
---	----------------------------------

	<p>Fires are generally not recommended as a control method for <i>Ehrharta calycina</i> because (1) this species is able to resprout after fires (DiTomaso et al., 2013), (2) fires can cause damage to non-target species (DiTomaso et al., 2013), and (3) <i>E. calycina</i> tends to increase in abundance with increased fire frequency (Milberg & Lamont, 1995). However, intense fires can also kill off a large portion of this species' seedbank because most of its seeds accumulate in the topsoil (Smith et al., 1999). Although fire does not stimulate germination of this species (Smith et al., 1999), seedlings that emerge after fires can be controlled with herbicides (Western Australian Herbarium, 1998).</p> <p>The objective of this measure is to mitigate the impacts and control populations of this species.</p>																					
<p>Scale of application At what scale is the measure applied? What is the largest scale at which it has been successfully used? Please provide examples, with areas (km² or ha) if possible.</p>	<p>It is unknown at what scales burning could be applied.</p>																					
<p>Effectiveness of the measure Is it effective in relation to its objective? Has the measure previously worked, failed? Please select one of the categories of effectiveness (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<table border="1" data-bbox="618 689 1854 751"> <tr> <td><i>Effectiveness of measures</i></td> <td><i>Effective</i></td> <td></td> <td><i>Neutral</i></td> <td>X</td> <td><i>Ineffective</i></td> <td></td> </tr> </table> <p><i>Rationale:</i> There are no studies to assess the effectiveness of this measure.</p>	<i>Effectiveness of measures</i>	<i>Effective</i>		<i>Neutral</i>	X	<i>Ineffective</i>															
<i>Effectiveness of measures</i>	<i>Effective</i>		<i>Neutral</i>	X	<i>Ineffective</i>																	
<p>Effort required e.g. period of time over which measure need to be applied to have results</p>	<p>The effort required is unknown.</p>																					
<p>Resources required¹ e.g. cost, staff, equipment etc.</p>	<p>Staff trained in managing prescribed burns, firefighting equipment, fuel to ignite fires. Costs are unknown.</p>																					
<p>Side effects (incl. potential) – both positive and negative i.e. positive or negative side effects of the measure on public health, environment including non-targeted species, etc.</p>	<table border="1" data-bbox="618 1193 1854 1294"> <tr> <td><i>Environmental effects</i></td> <td><i>Positive</i></td> <td></td> <td><i>Neutral or mixed</i></td> <td>X</td> <td><i>Negative</i></td> <td></td> </tr> <tr> <td><i>Social effects</i></td> <td><i>Positive</i></td> <td></td> <td><i>Neutral or mixed</i></td> <td>X</td> <td><i>Negative</i></td> <td></td> </tr> <tr> <td><i>Economic effects</i></td> <td><i>Positive</i></td> <td></td> <td><i>Neutral or mixed</i></td> <td>X</td> <td><i>Negative</i></td> <td></td> </tr> </table> <p><i>Rationale:</i> There may be negative environmental effects since non-target species could be killed during prescribed burns. There could also be negative social and economic effects if fires burn out of control.</p>	<i>Environmental effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>	X	<i>Negative</i>		<i>Social effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>	X	<i>Negative</i>		<i>Economic effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>	X	<i>Negative</i>	
<i>Environmental effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>	X	<i>Negative</i>																	
<i>Social effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>	X	<i>Negative</i>																	
<i>Economic effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>	X	<i>Negative</i>																	

<p>For each of the side effect types please select one of the impact categories (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>									
<p>Acceptability to stakeholders e.g. impacted economic activities, animal welfare considerations, public perception, etc.</p> <p>Please select one of the categories of acceptability (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<p>Acceptability to stakeholders</p>	<p><i>Acceptable</i></p>		<p><i>Neutral or mixed</i></p>		<p><i>Unacceptable</i></p>	<p>X</p>	<p><i>Rationale:</i> At present this measure is unlikely to be acceptable to invasive species managers because of a lack of evidence for its effectiveness. The general public is also likely to be wary of fires, especially if these are near to where people live or work.</p>	
<p>Additional cost information¹ When not already included above, or in the species Risk Assessment.</p> <ul style="list-style-type: none"> - implementation cost for Member States - the cost of inaction - the cost-effectiveness - the socio-economic aspects <p>Include quantitative &/or qualitative data, and case studies (incl. from countries outside the EU).</p>	<p>Implementation costs are unknown.</p> <p>Costs of inaction associated with this measure are likely similar to those detailed in the <i>Prevention of intentional introductions and spread</i> table.</p> <p>The cost-effectiveness of prescribed burns for management of this species is unknown.</p> <p>There are no known socio-economic aspects.</p>								
<p>Level of confidence on the information provided²</p> <p>Please select one of the confidence categories along with a statement to support the category chosen. See <i>Notes</i> section at the bottom of this document.</p> <p>NOTE – this is not related to the effectiveness of the measure</p>	<p><i>Inconclusive</i></p>	<p>X</p>	<p><i>Unresolved</i></p>		<p><i>Established but incomplete</i></p>		<p><i>Well established</i></p>	<p><i>Rationale:</i> There are a very limited number of studies that have investigated the use of fire for controlling <i>E. calycina</i>. It is not known how effective this measure is, what the costs are likely to be, and how this measure should be implemented.</p>	

Bibliography³

See guidance section

- Bacelar, H., Silva-Pando, F.J. & Pinto da Silva, A. R. (1989). O género *Ehrharta* na Europa. A propósito da ocorrência de *E. calycina* Sm. em Portugal e na Espanha. In Consellería de Agricultura, Xunta de Galicia (Ed.) *Sobre flora y vegetación de Galicia* (pp. 95-99). Santiago.
- Charpin, A. & Zarco, C. R. (1982). Presencia de *Ehrharta calycina* Smith en España. *Saussurea*, 13, 187-188.
- Chimera, C. (2015). Hawai'i-Pacific Weed Risk Assessment for *Ehrharta calycina*. Retrieved from <https://www.dropbox.com/sh/eafyh6cvmao4tnb/AABJ6ifqIXEUKPUFag52o8oNa/Ehrharta%20calycina.pdf?dl=0>
- Conn, J. S., Stockdale, C. A., Werdin-Pfisterer, N. R., & Morgan, J. C. (2010). Characterizing pathways of invasive plant spread to Alaska: II. Propagules from imported hay and straw. *Invasive Plant Science and Management*, 3(3), 276-285. <https://doi.org/10.1614/IPSM-D-09-00041.1>
- DiTomaso, J. M., Kyser, G. B., Oneto, et al. (2013). Weed Control in Natural Areas in the Western United States. University of California, Davis, CA: Weed Research and Information Center. Retrieved from https://wric.ucdavis.edu/information/natural%20areas/wr_E/Ehrharta.pdf
- EPPO (2018). *Pest risk analysis for Ehrharta calycina*. Paris: EPPO
- Fish, L., Mashau, A. C., Moeaha, M. J., Nembudani, M. T. (2015). Identification guide to southern African grasses. *Strelitzia*, 36, 271–276. Pretoria: South African National Biodiversity Institute.
- Fisher, J. L., Veneklaas, E. J., Lambers, H., & Loneragan, W. A. (2006). Enhanced soil and leaf nutrient status of a Western Australian Banksia woodland community invaded by *Ehrharta calycina* and *Pelargonium capitatum*. *Plant and Soil*, 284(1-2), 253-264. <https://doi.org/10.1007/s11104-006-0042-z>
- Fisher, J. L., Loneragan, W. A., Dixon, K., Delaney, J., & Veneklaas, E. J. (2009). Altered vegetation structure and composition linked to fire frequency and plant invasion in a biodiverse woodland. *Biological Conservation*, 142(10), 2270-2281. <https://doi.org/10.1016/j.biocon.2009.05.001>
- Fraga-Arguimbau, P. (2014). Notes i contribucions al coneixement de la flora de Menorca (X). Notes florístiques. *Bolletí de la Societat d'Història Natural de les Balears*, 57(1), pp.161-189.
- Frey M. (2005). *Element Stewardship Abstract for Ehrharta spp. Thunb.* Arlington, Virginia, USA: The Nature Conservancy. Retrieved from <http://www.imapinvasives.org/GIST/ESA/esapages/ehrhcaly.html>
- Gallo, T., & Waitt, D. (2011). Creating a successful citizen science model to detect and report invasive species. *BioScience*, 61(6), 459-465. <https://doi.org/10.1525/bio.2011.61.6.8>
- Genovesi, P., Scalera, R., Brunel, S., Roy, D., & Solarz, W. (2010). *Towards an early warning and information system for invasive alien species (IAS) threatening biodiversity in Europe* (EEA Technical Report No 5/2010). Copenhagen: European Environment Agency. Retrieved from <https://www.eea.europa.eu/publications/information-system-invasive-alien-species>
- Greuter, W. & Raus, T. (1998). Med-Checklist notulae, 17. *Willdenowia*, 28, 163-174.
- Grice, A. C., Clarkson, J. D., Murphy, H. T., Fletcher, C. S., & Westcott, D. A. (2013). Containment as a strategic option for managing plant invasion. *Plant Protection Quarterly*, 28(3), 62.

- IPPC (2017). ISMP 41 International movement of used vehicles, machinery and equipment. 12 pp. Rome: FAO. Retrieved from https://www.ippc.int/static/media/files/publication/en/2017/05/ISPM_41_2017_En_2017-05-15.pdf
- Kinkade, J. (2015). Grass attack: Efforts to manage veldt grass in the Guadalupe dunes go aerial. *New Times*. Retrieved from <https://www.newtimeslo.com/sanluisobispo/grass-attack-efforts-to-manage-velde-grass-in-the-guadalupe-dunes-go-aerial/Content?oid=2936908>
- Kettunen, M., Heikkilä, J., Underwood, E., & Vyliaudaite, I. (2014). *Instruments for financing action on invasive alien species (IAS): review and assessment of selected examples and their applicability in Finland*. London & Brussels: Institute for European Environmental Policy. Retrieved from https://ieep.eu/uploads/articles/attachments/d1b7abe7-75be-4aeb-8d82-b75194bf1272/Kettunen_at_al_2014_review_of_instruments_financing_IAS_action_FINAL_June_2014.pdf?v=63664509852
- Maistrello, L., Dioli, P., Bariselli, M., Mazzoli, G. L., & Giacalone-Forini, I. (2016). Citizen science and early detection of invasive species: phenology of first occurrences of *Halyomorpha halys* in Southern Europe. *Biological Invasions*, 18(11), 3109-3116. <https://doi.org/10.1007/s10530-016-1217-z>
- Mashau, A. C. (2008). *Ehrharta calycina* Sm. *PlantZAfrica species description*. Retrieved from <http://www.plantzafrica.com/plantefg/ehrhartacaly.htm>
- Milberg, P., & Lamont, B. B. (1995). Fire enhances weed invasion of roadside vegetation in southwestern Australia. *Biological Conservation*, 73(1), 45-49. [https://doi.org/10.1016/0006-3207\(95\)90061-6](https://doi.org/10.1016/0006-3207(95)90061-6)
- Moore, J. L., Runge, M. C., Webber, B. L., & Wilson, J. R. (2011). Contain or eradicate? Optimizing the management goal for Australian acacia invasions in the face of uncertainty. *Diversity and Distributions*, 17(5), 1047-1059. <https://doi.org/10.1111/j.1472-4642.2011.00809.x>
- Newsome, D., Smith, A., & Moore, S. A. (2008). Horse riding in protected areas: a critical review and implications for research and management. *Current Issues in Tourism*, 11(2), 144-166. <https://doi.org/10.2167/cit336.0>
- Pickart, A. (2000). *Ehrharta* spp. California Invasive Plant Council (CAL-IPC). Retrieved from <http://www.cal-ipc.org/ip/management/ipcw/pages/detailreport.cfm@usernumber=44&surveynumber=182.php>
- Quattrocchi, U., (2006). *CRC World Dictionary of Grasses: Common Names, Scientific Names, Eponyms, Synonyms, and Etymology - 3 Volume Set*. Boca Raton, Florida (USA): CRC Press Taylor & Francis.
- Ray, C.A., Sherman, J.J., Godhino, A.L., Hanson, N. & Parker, I.M. (2018). Impacts and best management practices for erect veldtgrass (*Ehrharta erecta*). *Invasive Plant Science and Management*, 11(1), 40-48 <https://doi.org/10.1017/inp.2018.2>
- Rossiter, R. C. (1947). Studies on perennial veldt grass (*Ehrharta calycina*). *Bulletin of the Council for Scientific and Industrial Research of Australia*, 227, pp.35+4.
- Shindler, B., Gordon, R., Brunson, M. W., & Olsen, C. (2011). Public perceptions of sagebrush ecosystem management in the Great Basin. *Rangeland Ecology & Management*, 64(4), 335-343. <https://doi.org/10.2111/REM-D-10-00012.1>
- Smith, M. A., Bell, D. T., & Loneragan, W. A. (1999). Comparative seed germination ecology of *Austrostipa compressa* and *Ehrharta calycina* (Poaceae) in a Western Australian Banksia woodland. *Australian Journal of Ecology*, 24(1), 35-42. <https://doi.org/10.1046/j.1442-9993.1999.00944.x>
- Trunzo, J., (2015). *Native small mammal use of an invasive grass: Heermann's kangaroo rats (Dipodomys heermanni) and Veldt grass (Ehrharta calycina) in coastal California* (M.Sc. thesis, California Polytechnic State University (USA)).
- Tulloch, A. I., Possingham, H. P., Joseph, L. N., Szabo, J., & Martin, T. G. (2013). Realising the full potential of citizen science monitoring programs. *Biological Conservation*, 165, 128-138. <https://doi.org/10.1016/j.biocon.2013.05.025>
- University of California, Berkeley (2018). *Jepson eFlora; Taxon page: Ehrharta calycina*. Retrieved from http://ucjeps.berkeley.edu/eflora/eflora_display.php?tid=23852

USFWS (2014). *Draft Environmental Assessment for the Aerial Herbicide Application Research Study Guadalupe-Nipomo Dunes National Wildlife Refuge San Luis Obispo County, California*. Retrieved from https://www.fws.gov/uploadedFiles/Region_8/NWRS/Zone_1/Hopper_Mountain_Complex/Guadalupe-Nipomo_Dunes/PDFs/Aerial_Spraying_Draft_EA_final.pdf

Valdés, B., Talavera, S., & Fernandez-Galiano, E., (1987). *Flora vascular de Andalucía Occidental: 3*. Barcelona (Spain): Ketres Editora.

Valdés, B. (2015). Xenophytes in the Doñana territory (SW Spain). *Flora Mediterranea*, 25, 55-64.

Van der Westhuizen, F. G. J. & Joubert, J. G. V., (1983). Die invloed van snoei tydens antese op koolsuurgasabsorpsie en koolhidraatinhoud van *Ehrharta calycina* en *Osteospermum sinuatum*. *Proceedings of the Annual Congresses of the Grassland Society of Southern Africa*, 18, 106-112.


Western Australian Herbarium (1998). *FloraBase—the Western Australian Flora: Ehrharta calycina Sm.* Retrieved from <https://florabase.dpaw.wa.gov.au/browse/profile/347>

Wittkuhn, R. S., (2010). Wind-aided seed dispersal of Perennial Veld Grass (*Ehrharta calycina*): implications for restoration in weedy urban bushland remnants. *Ecological Management & Restoration*, 11(2):148-150. <https://doi.org/10.1111/j.1442-8903.2010.00536.x>

Notes

1. Costs information. The assessment of the potential costs shall describe those costs quantitatively and/or qualitatively depending on what information is available. This can include case studies from across the Union or third countries.

2. Level of confidence¹: based on the quantity, quality and level of agreement in the evidence.



- **Well established:** comprehensive meta-analysis² or other synthesis or multiple independent studies that agree.
- **Established but incomplete:** general agreement although only a limited number of studies exist but no comprehensive synthesis and, or the studies that exist imprecisely address the question.
- **Unresolved:** multiple independent studies exist but conclusions do not agree.
- **Inconclusive:** limited evidence, recognising major knowledge gaps

3. Citations and bibliography. The APA formatting style for citing references in the text and in the bibliography is used.

e.g. Peer review papers will be written as follows:

In text citation: (Author & Author, Year)

¹ 2016. Guide on the production and integration of assessments from and across all scales (IPBES-4-INF-9), which is adapted from Moss and Schneider (2000). Assessment of confidence methodology is taken from IPBES.

² A statistical method for combining results from different studies which aims to identify patterns among study results, sources of disagreement among those results, or other relationships that may come to light in the context of multiple studies.

In bibliography: Author, A. A., & Author, B. B. (Publication Year). Article title. *Periodical Title*, Volume(Issue), pp.-pp.
(see <http://www.waikato.ac.nz/library/study/referencing/styles/apa>)