National Summary for Article 12

1. General information

1.1 Number and area of SPAs

The table below provides the total number and total area of sites designated under the Birds Directive (Special Protection Areas, SPAs), terrestrial area of sites and number and area of marine sites (i.e. any site with a marine component).

Empty cells in tables mean that the component requested was not applicable.

| All | | Terrestrial | | Marine | |
|-----------------------------------|------------|-------------|-----|------------|--|
| No. | Area (km²) | Area (km²) | No. | Area (km²) | |
| 56 | 13746 | 13746 | 0 | 0 | |
| Date of database used: 31-10-2012 | | | | | |

1.2 Number of SPAs with comprehensive management plans

Number of SPAs for which comprehensive management plans have been adopted: 13

Percentage of the network area covered by comprehensive management plans: 30%

Number of sites for which management plans are under preparation (optional field): not reported

1.3 Research and other work on bird populations

This section provides an indication of whether any of the activities listed in the section 6 of the General report have been carried out during the reporting period (for more details and references see the General report - the link to the report is given after the section 7 of this national summary).

National bird atlas: yes

National bird monitoring overview(s): yes

National bird red list: no

Other publication(s) of EU-wide interest: yes

2. Number of bird species/populations

This section provides a summary of the number of bird taxa (species and subspecific populations) for which a species-based report was completed, including a breakdown by season, and by subsets (e.g. Annex I, SPA trigger and non-native species).

| Season | All native taxa | Annex I | SPA trigger | Non-native |
|-----------|-----------------|---------|-------------|------------|
| Breeding | 217 | 71 | 83 | 1 |
| Wintering | 13 | 6 | 11 | 0 |
| Passage | 17 | 8 | 15 | 0 |
| Total | 247 | 85 | 109 | 1 |

Note: These statistics are based on the revised checklists. The harmonisation of the codes used for 'presence status' was needed and the summary of changes in comparison to the reported information by the Member State can be consulted through this link: http://bd.eionet.europa.eu/activities/Reporting Tool/Documents/Art 12 checklist changes.

Occasional or vagrant species, and species that went extinct nationally prior to 1980 (i.e. around the time the Birds Directive came into force), if indicated are excluded.

Number of taxa that went extinct nationally after 1980: 6

Number of newly arriving taxa: none

Number of taxa on checklist for which no reports received: none

3. Information on trends


This section provides information about trends of national bird populations.

Note: Article 12 reporting covers only a subset of Wintering taxa occurring in the national territory.

3.1 Population trends

The graphs show the percentages of taxa reported as having decreasing, stable, fluctuating, increasing or unknown population trends. Both short- and long-term population trends are included. The percentages are shown separately for breeding and wintering taxa.

Note: The trend category 'unknown' may include also taxa on the checklist for which no trend information was provided.




The table shows the numbers of taxa reported as having decreasing, stable, fluctuating, increasing or unknown population trends.

| Population trend | Breedi | ng taxa | Wintering taxa | | |
|------------------|------------|----------------------|----------------|-----------|--|
| | Short-term | Short-term Long-term | | Long-term | |
| Decreasing | 48 | 42 | 2 | 2 | |
| Stable | 42 | 3 | | 1 | |
| Fluctuating | 33 | 20 | 9 | 3 | |
| Increasing | 41 | 39 | 2 | 4 | |
| Unknown | 54 | 114 | | 3 | |


3.2 Comparison of population trends for subsets of taxa

The graphs show the percentages of taxa (all, Annex I and non-Annex I) within the different trend categories (see section 3.1). Both short- and long-term population trends are included. The graphs show results separately for breeding and wintering taxa.

Breeding taxa



Wintering taxa



The tables show the numbers of taxa (all, Annex I and non-Annex I) within the different trend categories.

Breeding taxa

| Population trend | Short-term | | | Long-term | | | |
|------------------|------------|---------|-------------|-----------|---------|-------------|--|
| | All taxa | Annex I | Non-Annex I | All taxa | Annex I | Non-Annex I | |
| Decreasing | 48 | 14 | 34 | 42 | 18 | 24 | |
| Stable | 42 | 13 | 29 | 3 | 1 | 2 | |
| Fluctuating | 33 | 20 | 13 | 20 | 10 | 10 | |
| Increasing | 41 | 15 | 26 | 39 | 23 | 16 | |
| Unknown | 54 | 9 | 45 | 114 | 19 | 95 | |

Wintering taxa

| Population trend | Short-term | | | | Long-term | |
|------------------|------------|---------|-------------|----------|-----------|-------------|
| | All taxa | Annex I | Non-Annex I | All taxa | Annex I | Non-Annex I |
| Decreasing | 2 | 1 | 1 | 2 | | 2 |
| Stable | | | | 1 | | 1 |
| Fluctuating | 9 | 4 | 5 | 3 | 2 | 1 |
| Increasing | 2 | 1 | 1 | 4 | 2 | 2 |
| Unknown | · | | | 3 | 2 | 1 |

3.3 Comparison of short- and long-term population trends

This section provides a comparison of short- and long-term population trends for taxa, highlighting combinations that represent potential improvements (in green) and deteriorations (in red) in their national status. The tables in this section show the numbers of taxa for each combination of short- and long-term trends.

Breeding taxa

| Long-term | Short-term population trend | | | | | |
|------------------|-----------------------------|--------|-------------|------------|---------|-------|
| population trend | Decreasing | Stable | Fluctuating | Increasing | Unknown | Total |
| Decreasing | 27 | 3 | 5 | | 7 | 42 |
| Stable | | 3 | | | | 3 |
| Fluctuating | 1 | | 19 | | | 20 |
| Increasing | | 6 | 5 | 19 | 9 | 39 |
| Unknown | 20 | 30 | 4 | 22 | 38 | 114 |
| Total | 48 | 42 | 33 | 41 | 54 | 218 |

Wintering taxa


| Long-term | Short-term population trend | | | | | |
|------------------|-----------------------------|--------|-------------|------------|---------|-------|
| population trend | Decreasing | Stable | Fluctuating | Increasing | Unknown | Total |
| Decreasing | 1 | | 1 | | | 2 |
| Stable | | | 1 | | | 1 |
| Fluctuating | | | 3 | | | 3 |
| Increasing | | | 2 | 2 | | 4 |
| Unknown | 1 | | 2 | | | 3 |
| Total | 2 | | 9 | 2 | | 13 |

3.4 Breeding range trends

Summary of the direction of short- and long-term range trends for breeding taxa.

The graphs show the percentages of taxa reported as having decreasing, stable, fluctuating, increasing or unknown breeding range trends. Both short- and long-term trends are included.

Note: The trend category 'unknown' may include also taxa on the checklist for which no trend information was provided.




The table shows the numbers of taxa reported as having decreasing, stable, fluctuating, increasing or unknown range trends.

| Breeding range trend | Breeding taxa | | | |
|----------------------|---------------|-----------|--|--|
| | Short-term | Long-term | | |
| Decreasing | 20 | 37 | | |
| Stable | 105 | 29 | | |
| Fluctuating | 28 | 22 | | |
| Increasing | 21 | 34 | | |
| Unknown | 44 | 96 | | |

3.5 Comparison of breeding range trends for subsets of taxa

The graphs show the percentages of bird taxa (all, Annex I and non-Annex I) within the different trend categories (see section 3.4). Both short- and long-term population trends are included.



The table shows the numbers of bird taxa (all, Annex I and non-Annex I) within the different trend categories.

| Population trend | Short-term | | | | Long-term | |
|------------------|------------|---------|-------------|----------|-----------|-------------|
| | All taxa | Annex I | Non-Annex I | All taxa | Annex I | Non-Annex I |
| Decreasing | 20 | 8 | 12 | 37 | 17 | 20 |
| Stable | 105 | 27 | 78 | 29 | 8 | 21 |
| Fluctuating | 28 | 15 | 13 | 22 | 10 | 12 |
| Increasing | 21 | 13 | 8 | 34 | 19 | 15 |
| Unknown | 44 | 8 | 36 | 96 | 17 | 79 |

3.6 Comparison of short- and long-term range trends

This section provides a comparison of short- and long-term range trends for taxa, highlighting combinations that represent potential improvements (in green) and deteriorations (in red) in national status. The table in this section shows the numbers of taxa for each combination of short- and long-term trends.

| Long-term range | Short-term range trend | | | | | |
|-----------------|------------------------|--------|-------------|------------|---------|-------|
| trend | Decreasing | Stable | Fluctuating | Increasing | Unknown | Total |
| Decreasing | 16 | 7 | 3 | 2 | 9 | 37 |
| Stable | | 28 | | 1 | | 29 |
| Fluctuating | | | 22 | | | 22 |
| Increasing | | 13 | 2 | 16 | 3 | 34 |
| Unknown | 4 | 57 | 1 | 2 | 32 | 96 |
| Total | 20 | 105 | 28 | 21 | 44 | 218 |

4. Implementation of international species plans


This section provides a summary of national implementation of international Species Action Plans (SAPs), Management Plans (MPs) and Brief Management Statements (BMSs) containing proposed actions in the Member State. The table shows the number of taxa with international plans and the number with national plans adopted.

| Type of plan | No. of taxa with international SAP, MP and BMS | No. of taxa with national plan adopted |
|----------------------------------|--|--|
| Species Action Plan (SAP) | 19 | 3 |
| Management Plan (MP) | 13 | |
| Brief Management Statement (BMS) | | |

5. Frequency of main pressures and threats

This section provides a summary of the main pressures/threats reported for taxa triggering SPA classification nationally. Only pressures/threats reported as having 'high' impact are considered in this section (one or more pressures/threats under each of the level 1 categories). For these high-impact pressures/threats a distinction is made in the bar-chart of those pressures/threats reported by the MS as primarily operating inside the Member State, or elsewhere.

Note: The figures under section 5 cover only taxa triggering SPA classifications nationally, i.e. those listed in Annex I, plus a selection of key migratory taxa for which SPAs have been classified, as identified in the species checklist.



% of taxa suffering one or more 'high' impact pressure/threat

Note: Threat/pressure categories not reported are omitted.

Total number of taxa considered in the calculation: 109

Number of taxa with no high ranking pressure/threat within country (or no pressure/threat reported): 19

Number of taxa with no high ranking pressure/threat in any location (or no pressure/threat reported): 18

| Pressure and threat categories | Number of taxa for which this threat/pressure was reported as having a 'high' impact |
|---|--|
| A - Agriculture | 44 |
| B - Sylviculture, forestry | 7 |
| C - Mining, extraction of materials and energy production | 3 |
| D - Transportation and service corridors | 7 |
| F - Biological resource use other than agriculture & forestry | 23 |
| G - Human intrusions and disturbances | 3 |
| H - Pollution | 2 |
| I - Invasive, other problematic species and genes | 1 |
| J - Natural System modifications* | 58 |
| K - Natural biotic and abiotic processes (without catastrophes) | 6 |
| L - Geological events, natural catastrophes | 2 |
| M - Climate change | 55 |
| XE - Threats and pressures from outside the EU territory | 1 |

^{*}e.g. fire and fire suppression, dredging, water abstractions from surface waters


6. SPA coverage and conservation measures

Note: The figures under section 6 cover only taxa listed in Annex I, plus a selection of key migratory taxa for which SPAs have been classified nationally, as identified in the species checklist.

6.1 Coverage of SPA trigger species populations by SPA network

This section provides a summary of the proportions of national populations of SPA trigger taxa occurring within the national SPA network. These graphs (separate graphs for wintering and breeding taxa) show the percentages of reported SPA trigger taxa in three classes based on their coverage by SPAs.

The geometric mean is used if Member States have reported minimum and maximum values. The table below shows the figures on which the calculations are based.



% of national population within the SPA network


% of national population within the SPA network

This table shows the number of reported SPA trigger taxa in three classes based on their coverage by SPA sites.

| _ | | T-4-1 | | | |
|----------------|-------|--------|---------|-------------------------|-------|
| Таха | 0-24% | 25-74% | 75-100% | unknown or not relevant | Total |
| Breeding taxa | 12 | 26 | 44 | 1 | 83 |
| Wintering taxa | | 4 | 7 | | 11 |

6.2 Main conservation measures

This section provides information on the relative importance of conservation measures at level 1 implemented during the reporting period for SPA trigger taxa. The graph shows the percentages of taxa for which one or more 'high' importance conservation measure was implemented.



% of taxa for which one or more 'high' impact measures were reported


Note: Numbers in brackets correspond to the numbers of reports where measure 1, 2, etc. is noted as being of high importance. Measures not reported are omitted.

Total number of assessments considered in the calculation: 109

Number of assessments with no high ranking conservation measures or no conservation measures at all reported: 2

6.3 Impact of conservation measures

This section provides information on effects of implemented conservation measures for each level 1 measure category. The figure shows, for each level 1 measure category, the frequency of reported effects. The table below shows the figures on which the calculations are based (full names of the measures are shown in the table).



% of bird taxa for which a particular effect of a 'high' impact measure was reported

■ maintain
■ enhance
■ longterm
■ no effect
■ unknown or not evaluated

Note: The numbers in brackets correspond to the total number of reported effects for all 'high' importance measures.

| | Number of reports | | | | | |
|--|-------------------|---------|----------|--|--------------------------------|--|
| Measure | | enhance | longterm | | unknown or not evaluated | |
| 2 - Measures related to agriculture and open habitats | 63 | 2 | | | | |
| 3 - Measures related to forests and wooded habitats | 23 | 10 | | | | |
| 4 - Measures related to wetland, freshwater and coastal habitats | 112 | 7 | | | | |
| 6 - Measures related to spatial planning | 147 | | | | | |
| 7 - Measures related to hunting, taking and fishing and species management | 49 | 14 | | | | |
| 8 - Measures related to urban areas, industry, energy and transport | 15 | | | | | |

The following categories were used by the Member States to show effects of implemented conservation measures:

- <u>a) Maintain</u> when the conservation measure is required to maintain the population size on the present level and/or to prevent any declining trend.
- b) Enhance when the conservation measure is required to increase the population size from a currently low level and/or to prevent a further declining trend alone or in conjunction with other measures.
- c) Long-term measure without short-term effect one reporting cycle or less but long-term positive effect in terms of increase of population size and/or turning a declining trend is expected.
- <u>d) No effect</u> measure without effect or that needs adaptation and that is not delivering any conservation benefit; measure failed in achieving its objectives or had adverse effects.
- e) Unknown effect.
- f) Not evaluated if the effect of the measure has not been evaluated.

7. Data quality and completeness

7.1 Mandatory information missing or reported as unknown (%)

The aim of this section is to provide an overview of the data gaps in the report; most of these gaps are due to insufficient knowledge. This section does not refer to potential errors or technical problems in the Member State's report and concentrates on what is relevant for evaluating data completeness.

The tables give the percentages of bird taxa with unknown or missing information for components of bird status.

Note: The statistics on missing and unknown information may also include missing and unknown information for recent coloniser, species which are on verge of extinction or species with marginal population in the national territory for which certain fields in the reporting format may not be relevant and therefore corresponding information was not reported.

7.1 a) Mandatory information missing (%)

| | Size | 0 |
|-----------------------|---------------|---|
| Population (breeding) | Trend (short) | 0 |
| | Trend (long) | 0 |
| | Size | 0 |
| Population (winter) | Trend (short) | 0 |
| | Trend (long) | 0 |
| | Area | 0 |
| Range (breeding) | Trend (short) | 0 |
| | Trend (long) | 0 |
| Pressures | s & threats | 0 |
| SPA network | Coverage | 0 |
| SFA HELWORK | Measures | 0 |
| Ма | 0 | |

7.1. b) Mandatory information reported as unknown (%)

| | Size | 0 |
|-----------------------|---------------|-----|
| Population (breeding) | Trend (short) | 25 |
| | Trend (long) | 54 |
| | Size | 0 |
| Population (winter) | Trend (short) | 0 |
| | Trend (long) | 23 |
| | Area | 0.5 |
| Range (breeding) | Trend (short) | 20 |
| | Trend (long) | 45 |
| Pressures | s & threats | 0 |
| SPA network | Coverage | 0 |
| SFA HELWORK | Measures | 0 |
| Ma | aps | 0.5 |

7.2 Data quality reported for key population and range parameters (%)

This section presents statistics on the data quality reported by Member States for key parameters of bird status.

| | Breeding population | | | Breeding range | | | Wintering population | | |
|--------------|---------------------|------------------|-----------------|----------------|------------------|-----------------|----------------------|------------------|-----------------|
| Data quality | Size | Trend (short) | Trend (long) | Area | Trend (short) | Trend (long) | Size | Trend (short) | Trend (long) |
| Good (%) | 17 | 16 | 11 | 61 | 11 | 11 | 38 | 23 | 38 |
| Moderate (%) | 56 | 56 | 21 | 37 | 17 | 17 | 46 | 77 | 15 |
| Poor (%) | 27 | 28 | 67 | 2 | 71 | 72 | 15 | 0 | 38 |
| No data (%) | 0 | 0 | 2 | 0 | 0 | 0 | 0 | 0 | 8 |

Source of information:

Link to the national general report on CDR
Link to the national report for birds on CDR

8. Bird species/subspecific populations reported

This section provides the list of bird taxa reported by the Member State, and the population size and short-term population trend direction ('+' increasing, '-' decreasing, '0' stable, 'F' fluctuating, 'x' unknown) for breeding and wintering taxa (the order of species follows the alphabetical order). For SPA trigger taxa occurring on passage an indication of presence or the size of the population is also provided.

For breeding taxa, population size is reported as number of breeding pairs, with just a few exceptions (which are indicated in the table), whereas population sizes for all wintering and passage taxa are in individuals.

Taxa listed on Annex I of the Directive are identified with a 'Y' in the 'Annex I' column. If the Member State reported on non-native taxa (other than for the three taxa listed in Annex II of the Birds Directive) the summary on these taxa is given in a separate table.

| Code | Species/subspecific population | Annex I | Breeding | Wintering | Passage |
|--------|--|----------------|-------------------|-------------------|---------|
| A402 | Accipiter brevipes | Υ | 0-1 (F) | | |
| A619 | Accipiter gentilis gentilis | N | 800-1300 (-) | | |
| A633 | Accipiter nisus nisus | N | 3300-4300 (x) | | |
| A298 | Acrocephalus arundinaceus | N | 200000-280000 (0) | | |
| A293 | Acrocephalus melanopogon | Υ | 10000-20000 (x) | | |
| A294 | Acrocephalus paludicola | Υ | 0 cmales (-) | | |
| A296 | Acrocephalus palustris | N | 56000-91000 (-) | | |
| A295 | Acrocephalus schoenobaenus | N | 224000-356000 (-) | | |
| A297 | Acrocephalus scirpaceus | N | 70000-137000 (-) | | |
| A168 | Actitis hypoleucos | N | 100- (F) | | |
| A324 | Aegithalos caudatus | N | 137000-229000 (+) | | |
| A223 | Aegolius funereus | Υ | 0-1 (F) | | |
| A247 | Alauda arvensis | N | 730000-900000 (-) | | |
| A229 | Alcedo atthis | Υ | 600- (x) | | |
| A054 | Anas acuta | N | 0-40 (-) | | |
| A056 | Anas clypeata | N | 150-320 (-) | | Р |
| A704 | Anas crecca crecca | N | 0-15 (x) | | Р |
| A050 | Anas penelope | N | | | Р |
| A705 | Anas platyrhynchos platyrhynchos | N | 50000- (-) | | Р |
| A055 | Anas querquedula [Western Siberia & Europe/West Africa] | N | 800-1500 (-) | | |
| A703 | Anas strepera strepera | N | 160-400 (+) | | Р |
| A394 | Anser albifrons albifrons | N | | 120000-200000 (+) | |
| A043 | Anser anser | N | 2100-3300 (+) | 20000-45000 (F) | |
| A042 | Anser erythropus | Υ | | | Р |
| A702 | Anser fabalis rossicus [West & Central Siberia/NE & SW Europe] | N | | 5000-13000 (-) | |
| A255 | Anthus campestris | Υ | 2400-4800 (x) | | |
| A256 | Anthus trivialis | N | 120000-190000 (0) | | |
| A226 | Apus apus | N | 3000-5000 (x) | | |
| A091 | Aquila chrysaetos | Υ | 4-5 (0) | | |
| A404 | Aquila heliaca | Υ | 125-160 (+) | | |
| A089 | Aquila pomarina | Υ | 27-40 (-) | | |
| A699 | Ardea cinerea cinerea | N | 3000-4000 (0) | | |
| A634-B | Ardea purpurea purpurea [East Europe, Black Sea & Mediterranean/Sub- | Y ^a | 800-1400 (0) | | |

| Code | Species/subspecific population | Annex I | Breeding | Wintering | Passage |
|--------|---|-----------------|-------------------------|-----------------|---------|
| | Saharan Africa] | | | | |
| A635 | Ardeola ralloides ralloides | Y ^a | 250-450 (F) | | |
| A222 | Asio flammeus | Υ | 10-80 (F) | 50-500 (F) | |
| A221 | Asio otus | N | 6500-12000 (x) | | |
| A218 | Athene noctua | N | 1500-2500 (x) | | |
| A059 | Aythya ferina | N | 2000-3000 (-) | 10000-15000 (F) | |
| A061 | Aythya fuligula | N | 70-100 (0) | 8000-12000 (F) | |
| A060-B | Aythya nyroca [Eastern Europe/E Mediterranean & Sahelian Africa] | Υ | 800-1500 (+) | | |
| A104 | Bonasa bonasia | Υ | 60-100 (-) | | |
| A688-B | Botaurus stellaris stellaris [C & E Europe, Black Sea & E Mediterranean (bre)] | Y ^a | 700-1000 cmales (x) | | |
| A396 | Branta ruficollis [Northern Siberia/Black Sea & Caspian] | Υ | | 150-650 (+) | |
| A215 | Bubo bubo | Υ | 50-60 (+) | | |
| A067 | Bucephala clangula | N | 0-1 (F) | 12000-18000 (F) | |
| A133 | Burhinus oedicnemus | Υ | 50-120 (-) | ` , | |
| A087 | Buteo buteo | N | 15000-30000 (0) | | |
| A088 | Buteo lagopus | N | , , | 300-500 (F) | |
| A403 | Buteo rufinus | Υ | 5-15 (+) | , | |
| A243 | Calandrella brachydactyla | Υ | 6-20 (-) | | |
| A149 | Calidris alpina [all non-breeding populations] | N | | | Р |
| A224 | Caprimulgus europaeus | Υ | 1000- cmales (-) | | |
| A366 | Carduelis cannabina | N | 71000-107000 (0) | | |
| A364 | Carduelis carduelis | N | 622000-763000 (0) | | |
| A745 | Carduelis chloris | N | 525000-639000 (+) | | |
| A365 | Carduelis spinus | N | 200-300 (x) | | |
| A698 | Casmerodius albus albus [W, C & SE Europe/Black Sea & Mediterranean] | Y ^{ba} | 3600-5500 (+) | | |
| A637 | Certhia brachydactyla all others | N | 25300-53400 (x) | | |
| A334 | Certhia familiaris | N | 5000-23000 (x) | | |
| A288 | Cettia cetti | N | 0-2 (F) | | |
| A682-B | Charadrius alexandrinus alexandrinus [Black Sea & East Mediterranean/Eastern Sahel] | Y ^a | 0-22 (-) | | |
| A726 | Charadrius dubius curonicus [Europe & North-west Africa/West Africa] | N | 800-1500 (x) | | |
| A734 | Chlidonias hybrida | Υ | 1000-6000 (F) | | |
| A198 | Chlidonias leucopterus | N | 30-4500 (F) | | |
| A197 | Chlidonias niger | Υ | 100-1400 (F) | | |
| A667-B | Ciconia ciconia ciconia [Central & Eastern Europe/Sub-Saharan Africa] | Y ^a | 5000-5500 (0) | | |
| A030-B | Ciconia nigra [Central & Eastern Europe/Sub-Saharan Africa] | Υ | 380-420 (+) | | |
| A264 | Cinclus cinclus | N | 5-10 (x) | | |
| A080 | Circaetus gallicus | Υ | 24-51 (F) | | |
| A081 | Circus aeruginosus | Υ | 5000-10000 bfemales (x) | | |
| A082 | Circus cyaneus | Υ | | 600-1200 (F) | |
| A084 | Circus pygargus | Υ | 50-200 bfemales (-) | , , | |
| A373 | Coccothraustes coccothraustes | N | 112000-163000 (+) | | |
| A206 | Columba livia [livia and domestica] | N | 146000-379000 (x) | | |
| A207 | Columba oenas | N | 5000-10000 (x) | | |
| A687 | Columba palumbus palumbus | N | 49000-116000 (+) | | |

| Code | Species/subspecific population | Annex I | Breeding | Wintering | Passage |
|--------------|---|---------------------|------------------------|-------------|---------|
| A231 | Coracias garrulus | Υ | 750-1050 (+) | | |
| A350 | Corvus corax | N | 4900-6000 (x) | | |
| A742 | Corvus corone cornix | N | 37000-45000 (+) | | |
| A743 | Corvus corone corone | N | 20-50 (x) | | |
| A348 | Corvus frugilegus | N | 18500-23500 (0) | | |
| A347 | Corvus monedula | N | 3000-6800 (-) | | |
| A113 | Coturnix coturnix | N | 29000-37000 cmales (-) | | |
| A122 | Crex crex [Europe & Western Asia/Sub-Saharan Africa] | Υ | 500-2000 cmales (F) | | |
| A212 | Cuculus canorus | N | 13400-14100 cmales (-) | | |
| A038-A | Cygnus cygnus [North-west Mainland Europe] | Υ | 1 (0) | | |
| A036 | Cygnus olor | N | 300-450 (+) | | |
| A738 | Delichon urbicum | N | 27000-36000 (-) | | |
| A239 | Dendrocopos leucotos | Υ | 260-760 (0) | | |
| A658 | Dendrocopos major all others | N | 274000-349000 (+) | | |
| A238 | Dendrocopos medius | Υ | 7000-16000 (0) | | |
| A240 | Dendrocopos minor | N | 5000-10000 (x) | | |
| A429 | Dendrocopos syriacus | Y | 21200-41900 (x) | | |
| A236 | Dryocopus martius | Ý | 9400-13100 (x) | | |
| A697 | Egretta garzetta | Y ^a | 750-1400 (F) | | |
| A378 | Emberiza cia | N | 400-700 (0) | | |
| A377 | Emberiza cirlus | N | 0 (-) | | |
| A376 | Emberiza citrinella | N | 605000-756000 (0) | | |
| A379 | Emberiza hortulana | Y | 0 (-) | | |
| A381 | Emberiza schoeniclus | N | 63000-104000 (+) | | |
| A269 | Erithacus rubecula | N | 306000-409000 (+) | | |
| A727 | Eudromias morinellus [Europe/North-west Africa] | Y | 300000-403000 (1) | | Р |
| A511 | Falco cherrug | Ý | 220-245 (+) | | ' |
| A098 | Falco columbarius | Ý | 220-243 (1) | 100-150 (F) | |
| A708 | Falco peregrinus peregrinus | Y ^a | 20-32 (+) | 100-130 (1) | |
| A099 | Falco subbuteo | N | 1900-2900 (x) | | |
| A099 A096 | Falco tinnunculus | N | 6200-7400 (x) | | + |
| A090 A097 | Falco vespertinus | Y | 700-1200 (-) | | - |
| A321 | Ficedula albicollis | Y | 7000-1200 (-) | | |
| A321 | Ficedula hypoleuca | N | 30-60 (x) | | + |
| A322 | Ficedula nypoleuca Ficedula parva | Y | 60-150 (-) | | + |
| A657 | Fringilla coelebs all others | N N | 1130000-1360000 (+) | | - |
| A723 | Fullica atra atra | N | 25000-130000 (+) | | |
| A723 A244 | Galerida cristata | N | 50000-100000 (x) | | |
| A244 A153 | Gallinago gallinago | N | 300-600 (F) | | |
| A753 A721 | Gallinula chloropus chloropus [Europe & North Africa] | | | | - |
| | | N N | 6000-12000 (x) | | |
| A342 | Garrulus glandarius | N | 29000-72000 (0) | | - |
| A515 | Glareola nordmanni [SE Europe & Western Asia/Southern Africa] | N Y ^a | 0-1 (F) | | - |
| A625-B | Glareola pratincola pratincola [Black Sea & E Mediterranean/Eastern Sahel zone] | , | 20-40 (F) | | |
| A217 | Glaucidium passerinum | Υ | 0-2 (F) | | |

| Code | Species/subspecific population | Annex I | Breeding | Wintering | Passage |
|--------|---|----------------|-------------------|--------------|---------|
| A639-B | Grus grus grus [other populations] | Y ^a | | | Р |
| A075 | Haliaeetus albicilla | Υ | 226-271 (+) | 500-800 (F) | |
| A092 | Hieraaetus pennatus | Υ | 0-2 (x) | | |
| A131 | Himantopus himantopus | Υ | 200-1000 (F) | | |
| A299 | Hippolais icterina | N | 6500-15000 (x) | | |
| A740 | Hippolais pallida | N | 200-500 (x) | | |
| A251 | Hirundo rustica | N | 89000-102000 (-) | | |
| A617-B | Ixobrychus minutus minutus [C & E Europe, Black Sea & E Mediterranean/Sub-Saharan Africa] | Y ^a | 4000-6000 (0) | | |
| A233 | Jynx torquilla | N | 11400-15300 (+) | | |
| A338 | Lanius collurio | Υ | 56000-65000 (-) | | |
| A653 | Lanius excubitor excubitor | N | 5-40 (+) | | |
| A339 | Lanius minor | Υ | 3300-4500 (-) | | |
| A459 | Larus cachinnans | N | 1-27 (x) | | |
| A182 | Larus canus | N | 0-2 (F) | | |
| A176 | Larus melanocephalus | Υ | 230-590 (+) | | |
| A604 | Larus michahellis | N | 0-1 (F) | | |
| A179 | Larus ridibundus | N | 6000- (F) | | |
| A614-B | Limosa limosa [Eastern Europe/Central & Eastern Africa] | N | 120-600 (-) | | Р |
| A291 | Locustella fluviatilis | N | 9500-14200 (-) | | |
| A292 | Locustella luscinioides | N | 58700-98300 (x) | | |
| A290 | Locustella naevia | N | 4700-7800 (-) | | |
| A369 | Loxia curvirostra | N | 200-400 (x) | | |
| A246 | Lullula arborea | Υ | 1000- (-) | | |
| A270 | Luscinia luscinia | N | 0 (x) | | |
| A271 | Luscinia megarhynchos | N | 225000-281000 (0) | | |
| A612 | Luscinia svecica cyanecula | Y ^a | 1200-1800 (+) | | |
| A767-B | Mergellus albellus [North-west & Central Europe (win)] | Υ | () | 600-1500 (-) | |
| A654-B | Mergus merganser merganser [other populations] | N | 1 (0) | () | |
| A230 | Merops apiaster | N | 17000-24000 (0) | | |
| A746 | Miliaria calandra | N | 62000-74000 (-) | | |
| A073 | Milvus migrans | Υ | 123-163 (0) | | |
| A074 | Milvus milvus | Y | 4-10 (0) | | |
| A280 | Monticola saxatilis | N | 0 (-) | | |
| A262 | Motacilla alba | N | 157000-210000 (0) | | |
| A261 | Motacilla cinerea | N | 220-320 (0) | | |
| A260 | Motacilla flava | N | 150000-225000 (0) | | |
| A319 | Muscicapa striata | N | 46000-74000 (-) | | |
| A058-A | Netta rufina [South-west & Central Europe/West Mediterranean] | N | 80-100 (+) | | |
| A768 | Numenius arquata arquata [Europe/Europe, North & West Africa] | N | 20-60 (x) | | Р |
| A158 | Numenius phaeopus | N | | | P |
| A159 | Numenius tenuirostris [Central Siberia/Mediterranean & SW Asia] | Ϋ́ | | | Р |
| A610-A | Nycticorax nycticorax [C & E Europe/Black Sea & E Mediterranean (bre)] | Y ^a | 2200-3600 (0) | | · |
| A277 | Oenanthe oenanthe | N | 20000-30000 (x) | | |

| Code | Species/subspecific population | Annex I | Breeding | Wintering | Passage |
|--------|--|----------------|----------------------|-----------|---------|
| A337 | Oriolus oriolus | N | 83000-96000 (0) | | |
| A129 | Otis tarda | Υ | 500-609 males (+) | | |
| A214 | Otus scops | N | 200-400 (0) | | |
| A094 | Pandion haliaetus | Υ | | | Р |
| A323 | Panurus biarmicus | N | 12000-15000 (0) | | |
| A656 | Parus ater all others | N | 10300-26000 (x) | | |
| A329 | Parus caeruleus | N | 220000-295000 (F) | | |
| A327 | Parus cristatus | N | 3500-5000 (x) | | |
| A330 | Parus major | N | 1330000-1610000 (+) | | |
| A326 | Parus montanus | N | 1500-2500 (x) | | |
| A325 | Parus palustris | N | 88600-136500 (+) | | |
| A620 | Passer domesticus | N | 970000-1250000 (-) | | |
| A356 | Passer montanus | N | 1100000-1370000 (0) | | |
| A644 | Perdix perdix all others | N | 11388-12419 (-) | | |
| A072 | Pernis apivorus | Υ | 500-750 (0) | | |
| A391 | Phalacrocorax carbo sinensis | N | 2500-2600 (-) | | |
| A393 | Phalacrocorax pygmeus [Black Sea & Mediterranean] | Υ | 600-1500 (+) | | |
| A115-X | Phasianus colchicus | N | 306403-339419 (-) | | |
| A151 | Philomachus pugnax | Υ | , | | Р |
| A273 | Phoenicurus ochruros | N | 214000-289000 (+) | | |
| A274 | Phoenicurus phoenicurus | N | 1000- (x) | | |
| A315 | Phylloscopus collybita | N | 529000-663000 (+) | | |
| A314 | Phylloscopus sibilatrix | N | 92400-152700 (0) | | |
| A316 | Phylloscopus trochilus | N | 15000-20000 (0) | | |
| A343 | Pica pica | N | 34000-86000 (0) | | |
| A234 | Picus canus | Y | 1500-2400 (0) | | |
| A235 | Picus viridis | N | 8300-11400 (x) | | |
| A607-B | Platalea leucorodia leucorodia [Cent. & SE Europe/Mediterranean & Tropical Africa] | Y ^a | 850-1400 (F) | | |
| A700 | Plegadis falcinellus falcinellus [Black Sea & Mediterranean/West Africa] | Y ^a | 2-20 (F) | | |
| A140 | Pluvialis apricaria | Υ | , , | | Р |
| A691 | Podiceps cristatus cristatus | N | 3000-4000 (x) | | |
| A665-B | Podiceps grisegena grisegena [Black Sea & Mediterranean (win)] | N | 40-80 (F) | | |
| A692 | Podiceps nigricollis nigricollis [Europe/South & West Europe & North Africa] | N | 300-1100 (F) | | |
| A719 | Porzana parva parva [Western Eurasia/Africa] | Y ^a | 1800-3000 cmales (x) | | |
| A119 | Porzana porzana [Europe/Africa] | Υ | 600-1500 cmales (F) | | |
| A720 | Porzana pusilla intermedia [Europe (bre)] | Y ^a | 0-15 cmales (F) | | |
| A266 | Prunella modularis | N | 1200-1500 (x) | | |
| A372 | Pyrrhula pyrrhula | N | 0-10 (-) | | |
| A718 | Rallus aquaticus aquaticus [Europe & North Africa] | N | 5000-10000 (x) | | |
| A132-B | Recurvirostra avosetta [South-east Europe, Black Sea & Turkey (bre)] | Y | 100-850 (F) | | |
| A318 | Regulus ignicapillus | N | 400-500 (x) | | |
| A317 | Regulus regulus | N | 1500-12000 (+) | | |
| A336 | Remiz pendulinus | N | 4500-13000 (x) | | |
| A249 | Riparia riparia | N | 15000-50000 (-) | | |

| Code | Species/subspecific population | Annex I | Breeding | Wintering | Passage |
|--------|--|----------------|---------------------|-----------|---------|
| A275 | Saxicola rubetra | N | 113000-183000 (-) | | |
| A276 | Saxicola torquatus | N | 360000-434000 (-) | | |
| A155 | Scolopax rusticola [Europe/South & West Europe & North Africa] | N | 10- cmales (x) | | |
| A361 | Serinus serinus | N | 142000-200000 (-) | | |
| A332 | Sitta europaea | N | 176000-247000 (+) | | |
| A631-A | Sterna albifrons albifrons [Europe north of Mediterranean (bre)] | Y ^a | 2-19 (F) | | |
| A193 | Sterna hirundo | Υ | 400-1500 (F) | | |
| A209 | Streptopelia decaocto | N | 291000-377000 (+) | | |
| A210 | Streptopelia turtur | N | 64000-150000 (0) | | |
| A219 | Strix aluco | N | 5000-8000 (0) | | |
| A220 | Strix uralensis | Υ | 160-260 (F) | | |
| A351 | Sturnus vulgaris | N | 710000-990000 (0) | | |
| A311 | Sylvia atricapilla | N | 929000-1121000 (+) | | |
| A310 | Sylvia borin | N | 500- (-) | | |
| A309 | Sylvia communis | N | 196000-267000 (-) | | |
| A308 | Sylvia curruca | N | 90000-136000 (0) | | |
| A307 | Sylvia nisoria | Υ | 38900-70800 (0) | | |
| A690 | Tachybaptus ruficollis ruficollis [Europe & North-west Africa] | N | 3300-5000 (x) | | |
| A048 | Tadorna tadorna | N | 5-10 (+) | | |
| A166 | Tringa glareola | Υ | | | Р |
| A162 | Tringa totanus | N | 400-1000 (F) | | |
| A676 | Troglodytes troglodytes all others | N | 27000-53000 (x) | | |
| A283 | Turdus merula | N | 1310000-1610000 (+) | | |
| A285 | Turdus philomelos | N | 365000-468000 (+) | | |
| A284 | Turdus pilaris | N | 30-100 (x) | | |
| A287 | Turdus viscivorus | N | 7500-19000 (x) | | |
| A213 | Tyto alba | N | 800-1000 (x) | | |
| A232 | Upupa epops | N | 5700-6800 (x) | | |
| A142 | Vanellus vanellus [Europe, W Asia/Europe, N Africa & SW Asia] | N | 29000-38000 (0) | | |

Note: The abbreviation Y^a is used for taxa (typically subspecies) listed in the Annex I at higher taxonomical level. The code Y^b indicates that the Annex I contains a synonym of the name used in the checklist.