

Information on measures and related costs in relation to species considered for inclusion on the Union list: *Cortaderia jubata*

This note has been drafted by IUCN within the framework of the contract No 07.0202/2017/763436/SER/ENV.D2 “Technical and Scientific support in relation to the Implementation of Regulation 1143/2014 on Invasive Alien Species”. The information and views set out in this note do not necessarily reflect the official opinion of the Commission, or IUCN. The Commission does not guarantee the accuracy of the data included in this note. Neither the Commission nor IUCN or any person acting on the Commission’s behalf, including any authors or contributors of the notes themselves, may be held responsible for the use which may be made of the information contained therein. Reproduction is authorised provided the source is acknowledged. This document shall be cited as:

Visser, V. 2018. Information on measures and related costs in relation to species considered for inclusion on the Union list: *Cortaderia jubata*. Technical note prepared by IUCN for the European Commission.

Date of completion: 20/08/2018

Comments which could support improvement of this document are welcome. Please send your comments by e-mail to ENV-IAS@ec.europa.eu.

Species (scientific name)	<i>Cortaderia jubata</i> (Lemoine ex Carrière) Stapf Bot. Mag. 124: t. 7607. 1898
Species (common name)	Andean pampas grass, Andes grass, Jubatagrass, jubata grass, pampas grass, pink pampas grass, purple pampas grass, pampasgras (Afrikaans), cortadera, sacuara (Spanish)
Author(s)	Vernon Visser, African Climate & Development Institute
Date Completed	20/08/2018
Reviewer	Gary J Houliston, Landcare Research

Summary

Highlight of measures that provide the most cost-effective options to prevent the introduction, achieve early detection, rapidly eradicate and manage the species, including significant gaps in information or knowledge to identify cost-effective measures.

Jubata grass, *Cortaderia jubata* (Lemoine ex Carrière) (Poaceae), is a perennial grass species native to Argentina, Chile, Bolivia, Ecuador, Peru and Colombia (Testoni & Villamil, 2014). It can grow up to 4 m in height and has large, serrated leaves and a tall, fluffy inflorescence (sometimes referred to as a plume) (Clayton, Vorontsova, Harman & Williamson, 2006 onwards). This species is currently not known to occur in the EU, but was trialled as an ornamental in the late 1800s in France, Ireland, and Switzerland (Carrière, 1878; Hooker, 1898) and more recently at two nurseries in the UK (Royal Horticultural Society, 2009). In the regions where this species has become invasive (Australia; New Zealand; South Africa; USA: California, Hawaii, Oregon), this species invades a wide range of habitats, but is most common in disturbed environments such as roadsides, disturbed vegetation, forestry plantations and recently burnt vegetation (Edgar & Connor, 2000; Parsons & Cuthbertson, 2004; Robinson, 1984; Starr, Starr & Loope, 2003).

Jubata grass outcompetes native plants in areas in which it invades, and can form monotypic stands (CABI, 2018; Daehler, 2006; Lambrinos, 2000; Queensland Government, 2018; Peterson & Russo, 1988). These invasions can cause reductions in the abundance and diversity of arthropods and rodents (Lambrinos, 2000). When jubata grass invades plantations, forestry operations are hampered and become much more expensive (Gadgil, Knowles & Zabkiewicz, 1984).

A common problem across all of the measures proposed in this note is that jubata grass and the closely related *C. selloana* are morphologically extremely similar (Testoni & Linder, 2017) and are easily confused with one another (Lambrinos, 2000; Robinson, 1984). Even taxonomic descriptions of these species are very variable (e.g. Edgar & Connor, 2000; Robinson, 1984; Testoni & Linder, 2017), probably because the two species represent a species complex: *C. selloana* is octoploid ($2n=72$) and *C. jubata* is duodecaploid ($2n=108$), and *C. selloana* is gynodioecious (the species has female and hermaphroditic plants) while *C. jubata* is apomictic (flowers do not require fertilisation to form seeds). These two aspects of the ecology of these two species suggest that *C. jubata* is derived from *C. selloana* (Testoni & Linder, 2017). Indeed, Testoni & Linder (2017) assign *C. jubata* to subspecies status of *C. selloana*. However, these species appear to be genetically distinct and are easily distinguished using genetic barcoding (Houliston & Goeke, 2017). *C. selloana* is already common across much of the EU (DAISIE European Invasive Alien Species Gateway, 2008) and it is possible that populations of jubata grass have been misidentified as *C. selloana*. This uncertainty makes it difficult to determine whether jubata grass has already established within the EU using taxonomic characters alone. Both species also have similar impacts (DiTomaso, 2010). If both species are listed as IAS of Union concern, many of the measures proposed here could be applied to both species and, to some degree, without much concern for the accurate identification of the species.

Prevention: The most appropriate measure for preventing entry of jubata grass into a Member State is *a ban on keeping, importing, selling, breeding and growing* of this species. Seeds are the most likely life stage to be introduced and should be banned, but live plants also have the potential to be introduced for horticultural purposes and also should be banned. *Phytosanitary measures* are likely to be ineffective for preventing entry via the principal pathways through which this species could be moved.

A survey of known introduction sites and a sample of C. selloana populations will help a great deal for determining whether jubata grass is already present in the EU. This measure is particularly important though as a **surveillance measure** and to support **early detection**. Surveys are recommended for the four known introduction sites in France, Switzerland and the UK, and a number of *C. selloana* populations in the EU, with a search radius of at least 50 km due to the possibility of seed dispersal (New Zealand Plant Conservation Network, 2018). A reassessment of the identity of any *Cortaderia* plants within the survey area will be required.

The use of *citizen science and resource managers' data* is another important **surveillance measure for early detection**. This is a low-cost option for early detection because these types of IT infrastructure and monitoring programs already exist. Jubata grass is easy to detect because of its distinctive appearance, but data collected through these networks will need to be carefully inspected for the correct species identification.

Measures for **rapid eradication for new introductions** and **management** of widespread invasions are the same for jubata grass, with physical control more feasible for smaller invasions and younger plants, and chemical control more cost effective and practical for larger invasions and difficult to reach plants (DiTomaso et al., 2010; Gosling, Shaw & Beadel, 2000; Popay, Timmins & McCluggage, 2003), but often a combination of both methods has been used in eradication and management programs (Gosling et al., 2000; Penniman, Buchanan & Loope, 2011). The effectiveness of these measures is largely dependent on locating and destroying jubata grass plants before they flower, because this species' seeds are able to disperse such large distances. However, the short viability of these seeds in the soil seedbank makes eradication and management easier if reproductive plants can be controlled. Biological control is currently not an option, with very little research having been done on control agents. Grazing has only been suggested as an effective control method in New Zealand (Gadgil et al., 1984; Gosling et al., 2000) and Australia (NSW Government, 2018), but is probably limited in its effectiveness because of the difficulties of managing grazing (Gadgil et al., 1984; Gosling et al., 2000), and because *C. jubata* still successfully invades

even under high grazing pressure (Lambrinos, 2006). Oversowing of pasture species, especially nitrogen-fixing plants, in timber plantations has also only been suggested as an effective control method in New Zealand, and only in combination with grazing (Gadgil et al., 1984; Gosling et al., 2000).

Prevention of intentional introductions and spread – measures for preventing the species being introduced intentionally. **This table is repeated for each of the prevention measures identified.**

<p>Measure description Provide a description of the measure, and identify its objective</p>	<p>A ban on keeping, importing, selling, breeding and growing (as would be required under Article 7 of the IAS Regulation).</p> <p>The principal pathway for intentional introductions of this species is via horticulture and horticultural plantings (CABI, 2018), although this species has also been (rarely) used as a forage plant (see references below). This measure therefore will seek to prevent the introduction and spread of <i>C. jubata</i> via these pathways.</p> <p>Although there is no evidence that <i>C. jubata</i> is promoted as an ornamental plant within the EU, this species has been historically planted as an ornamental in the late 1800s in France, Ireland, and Switzerland (Carrière, 1878; Hooker, 1898) and more recently was trialed as an ornamental at two nurseries in the UK (Royal Horticultural Society, 2009). In Australia (Queensland Government, 2017), California (Costas Lippmann, 1977; Peterson & Russo, 1988), Hawai'i (Starr et al., 2003), New Zealand (Houlston & Goeke, 2017) and South Africa (Robinson, 1984) this species is currently, or was recently, planted as an ornamental.</p> <p>Seeds of this species can also be purchased from online suppliers from outside of the EU (e.g., from https://www.amazon.com/PAMPAS-GRASS-Cortaderia-jubata-seeds/dp/B00480KMME). The large inflorescences of this species (and more so of <i>C. selloana</i>) were historically used for decorative purposes, principally in California (Costas Lippmann, 1977), and it seems that the inflorescences of at least <i>C. selloana</i> are available for floral bouquets (https://www.etsy.com/uk/market/pampas_grass), although the use of <i>C. jubata</i> for these purposes cannot be excluded.</p> <p><i>C. jubata</i> has been planted as a forage plant in California (Peterson & Russo, 1988) and New Zealand (Gadgil et al., 1984). There is no evidence that the species is promoted as a forage plant within the EU.</p> <p>This measure would need to be applied across the EU, but with a focus on areas at high risk of invasion (for details see EPPO, 2018).</p>				
<p>Effectiveness of the measure Is it effective in relation to its objective? Has the measure previously worked, failed?</p>	<p><i>Effectiveness of measures</i></p>	<p><i>Effective</i></p>	<p><i>Neutral</i></p>	<p><i>Ineffective</i></p>	<p>X</p>
<p><i>Rationale:</i> No specific information is available on the effectiveness of preventing intentional introductions through banning the keeping, importing, selling, breeding and growing of <i>C. jubata</i>. In addition, there is little evidence to suggest the species is currently found in the ornamental trade, though this could change in the future. However, there is good scientific support for producing positive net economic benefits from banning the import and introduction of potentially invasive ornamental plant species (e.g., Keller, Lodge & Finnoff, 2007).</p> <p>It is however possible that <i>C. jubata</i> is already established in the EU. <i>C. selloana</i> (commonly found in trade within the EU; DAISIE European Invasive Alien Species Gateway, 2008) and <i>C. jubata</i> can be easily confused and therefore one species may be misidentified for another, even by experts (Lambrinos, 2001; Robinson, 1984). Moreover, <i>C. jubata</i> is recorded from historical and</p>					

	<p>recent horticultural trials within the EU (Hooker, 1898; Royal Horticultural Society, 2009). Therefore, this measure is likely only to be effective if the same measures are applied for <i>C. selloana</i>, and even then this measure could possibly only be effective in limiting future introductions and subsequent re-invasion of <i>C. jubata</i> within the EU.</p> <p>This measure would require high administrative effort to ensure compliance and would need to be applied indefinitely due to the risk of introductions from elsewhere in the world. This measure would require a large budget to finance many well-trained staff to monitor and ensure compliance (Kettunen et al., 2014).</p>																					
<p>Side effects (incl. potential) – both positive and negative i.e. positive or negative side effects of the measure on public health, environment including non-targeted species, etc.</p> <p>For each of the side effect types please select one of the impact categories (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<table border="1" data-bbox="613 421 1854 528"> <tr> <td><i>Environmental effects</i></td> <td>Positive</td> <td></td> <td>Neutral or mixed</td> <td>X</td> <td>Negative</td> <td></td> </tr> <tr> <td><i>Social effects</i></td> <td>Positive</td> <td></td> <td>Neutral or mixed</td> <td>X</td> <td>Negative</td> <td></td> </tr> <tr> <td><i>Economic effects</i></td> <td>Positive</td> <td></td> <td>Neutral or mixed</td> <td>X</td> <td>Negative</td> <td></td> </tr> </table> <p><i>Rationale:</i> Although this species has been planted as an ornamental in places such as Australia, California, Hawai'i, New Zealand and South Africa (but only trialed as an ornamental within the EU), it is currently not widely sold as a horticultural species anywhere in the world, with <i>C. selloana</i> seeming to be the favoured species in this regard (Starr et al., 2003). Therefore, this measure is unlikely to have a negative economic side effect on the horticultural industry if only <i>C. jubata</i> (and not <i>C. selloana</i>) were to be banned.</p> <p>There will be no positive or negative environmental or social side effects to this measure.</p>	<i>Environmental effects</i>	Positive		Neutral or mixed	X	Negative		<i>Social effects</i>	Positive		Neutral or mixed	X	Negative		<i>Economic effects</i>	Positive		Neutral or mixed	X	Negative	
<i>Environmental effects</i>	Positive		Neutral or mixed	X	Negative																	
<i>Social effects</i>	Positive		Neutral or mixed	X	Negative																	
<i>Economic effects</i>	Positive		Neutral or mixed	X	Negative																	
<p>Acceptability to stakeholders e.g. impacted economic activities, animal welfare considerations, public perception, etc.</p> <p>Please select one of the categories of acceptability (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<table border="1" data-bbox="613 828 1854 895"> <tr> <td><i>Acceptability to stakeholders</i></td> <td>Acceptable</td> <td></td> <td>Neutral or mixed</td> <td>X</td> <td>Unacceptable</td> <td></td> </tr> </table> <p><i>Rationale:</i> It seems unlikely that this measure will be unacceptable to the horticultural industry or to the public. The only potentially significant use of <i>C. jubata</i> is for horticultural purposes, and even for this purpose this species seems to have little popularity: in recent trials conducted in the UK, <i>C. jubata</i> was grown at two nurseries, but neither nursery selected this species for further commercialisation (the species is not listed as available on their websites: http://www.dinglenurseryandgarden.co.uk, http://www.knollgardens.co.uk; the species was not given an Award of Garden Merit; Royal Horticultural Society, 2009).</p> <p>However, due to the difficulty of distinguishing between <i>C. jubata</i> and <i>C. selloana</i>, this measure is only likely to be effective if both species are banned. Due to the popularity of <i>C. selloana</i> as an ornamental species, a ban of this species is likely not to be acceptable to the horticultural industry.</p>	<i>Acceptability to stakeholders</i>	Acceptable		Neutral or mixed	X	Unacceptable															
<i>Acceptability to stakeholders</i>	Acceptable		Neutral or mixed	X	Unacceptable																	
<p>Additional cost information¹ When not already included above, or in the species Risk Assessment.</p>	<p>Implementation costs for Member States will be dependent on the cost of enforcing such a ban, but figures are not readily available in the public domain. Kettunen et al. (2014) suggest that costs for this type of measure will be relatively high.</p> <p>The cost of inaction of preventing intentional introductions of this species can be estimated based on costs of controlling invasions in other regions around the world. In California, the most cost-effective control method is glyphosate application (see</p>																					

- the cost of inaction
- the cost-effectiveness
- the socio-economic aspects

Include quantitative &/or qualitative data, and case studies (incl. from countries outside the EU).

Management tables for details) which costs USD 2,800 (ca. EUR 2,441) per hectare (DiTomaso, Drewitz & Kyser, 2008), and in New Zealand aerial application of herbicide can cost NZD 3,885 (ca. USD 2,500 / EUR 2,219) per hectare (Popay et al., 2003).

Another approach to estimating the **costs of inaction** of implementing this measure can be derived from estimates based on management and control of *C. selloana*, which is already widespread and invasive in the EU. In Spain, *C. selloana* was found to be the 6th most widespread invasive species and had the 13th most amount of money spent on its control in the last decade (EUR 8,600 in 8 of Spain’s 17 autonomous communities; this estimate is likely to be a gross underestimate of the true amount spent on this species’ management; Andreu, Vila & Hulme, 2009).

It is unknown whether this measure would be **cost-effective**. It has been suggested that because of the high costs of implementation and the high administrative burden, bans such as those suggested by this measure, are highly unlikely to be cost-effective (Kettunen et al., 2014). However, theoretical models suggest that there are major net positive economic benefits to preventing the entry of invasive species (Keller et al., 2007). There are, however, no known cost-benefit studies specific to *C. jubata*.

There are no known **socio-economic aspects**.

Level of confidence on the information provided ²

Please select one of the confidence categories along with a statement to support the category chosen. See *Notes* section at the bottom of this document.

NOTE – this is not related to the effectiveness of the measure

<i>Inconclusive</i>		<i>Unresolved</i>		<i>Established but incomplete</i>	X	<i>Well established</i>	
---------------------	--	-------------------	--	-----------------------------------	---	-------------------------	--

Rationale: There is a large body of literature (not specific to *C. jubata*) that supports a ban on keeping, importing, selling, breeding and growing an alien species that is found in trade. However, there is no information specific to *C. jubata* to support this measure, either in the EU or in third countries.

Prevention of un-intentional introductions and spread – measures for preventing the species being introduced un-intentionally (cf. Article 13 of the IAS Regulation). **This table is repeated for each of the prevention measures identified.**

<p>Measure description Provide a description of the measure, and identify its objective</p>	<p>Phytosanitary inspections, in particular related to the movement of garden waste, animals and soil, including soil on vehicles and machinery.</p> <p>A number of aspects of the biology of <i>C. jubata</i> are likely to influence the specific details of phytosanitary inspections for this species (see <i>Surveillance measures to support early detection</i> for further details on this species’ biology). This species usually flowers from mid-summer to early autumn (CABI, 2018; DiTomaso et al., 2010; Edgar & Connor, 2000; Robinson, 1984). Much of the invasive potential of pampas grass arises from its ability to produce thousands to millions of wind-dispersed seeds per year (up to 338,000 germinable seeds per year; Drewitz & DiTomaso, 2004), but most seeds only disperse within a small radius of the parent plants (Saura-Mas & Lloret, 2005) and are viable for only up to four months under winter (wet) field conditions (Drewitz & DiTomaso, 2004). While seeds are the primary mode of dispersal, this species is also able to resprout or re-establish from the upper rootstock (Drewitz & DiTomaso, 2004).</p> <p><i>C. jubata</i> has the potential to be introduced as a contaminant of garden waste, animals and soil, including soil on vehicles and machinery (CABI, 2018; University of Queensland, 2018). However, there are no quantitative measures of the extent and probability of such introductions occurring. Moreover, <i>C. jubata</i> may not even occur within the region (unless there are populations that have been misidentified as <i>C. selloana</i>; see <i>Summary</i> for details). Therefore, the likelihood of this species being unintentionally introduced via the movement of garden waste, animals and soil, including soil on vehicles and machinery within the EU is very low. Unintentional introductions of <i>C. jubata</i> with the importation of these materials/objects from areas outside of the EU where this species already occurs (see <i>Summary</i> for details) is possible, but has a low probability given that there are no documented cases of introductions into new countries via these pathways, this species has very short-lived seed viability, and previous introductions of this species into new countries were seemingly intentional (for horticulture).</p> <p>It is only recently that an ISPM Standard, no. 41 (IPPC, 2017), has been drafted and adopted on ‘International movement of used vehicles, machinery and equipment’. This focuses on reducing the risks of transporting contaminants (soil, seeds, plant debris, pests) associated with the international movement (either traded or for operational relocation) of vehicles, machinery and equipment (VME) that may have been used in agriculture and forestry, as well as for construction, industrial, mining waste management, and military purposes.</p> <p>For those VMEs that represent a contaminant risk, the phytosanitary measures recommended are detailed in the ISPM, and cover cleaning, prevention and disposal requirements. These include cleaning using pressure washing or compressed air cleaning, chemical or temperature treatments, storing and handling VMEs that prevent contact with soil, and keeping vegetation short around storage areas or ports.</p> <p>Phytosanitary inspections of these different materials/objects would vary and need to be developed or aligned with current phytosanitary measures.</p>
--	---

	<p>The objective of such a measure would be to prevent the entry of seeds or root materials of this species into the EU and into Member States, where this species has not established, to prevent its further unintentional spread within the EU. This measure is, however, dependent to some degree on whether <i>C. jubata</i> populations are found within the EU (see <i>Surveillance measures to support early detection – Survey of known introduction sites</i>).</p>			
<p>Scale of application At what scale is the measure applied? What is the largest scale at which it has been successfully used? Please provide examples, with areas (km² or ha) if possible.</p>	<p>This measure would need to be applied across the EU, but with a focus on movement of these materials/objects from areas of known introductions of <i>C. jubata</i>, both within and outside of the EU.</p> <p>No phytosanitary measures currently exist for this species, or for the similar <i>C. selloana</i> in the EU.</p>			
<p>Effectiveness of the measure Is it effective in relation to its objective? Has the measure previously worked, failed?</p> <p>Please select one of the categories of effectiveness (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<p>Effectiveness of measures</p>	<p>Effective</p>	<p>Neutral</p>	<p>Ineffective X</p>
<p><i>Rationale:</i> Identification of <i>C. jubata</i> seeds and root material is impossible without genetic barcoding, or when found together with the large inflorescences. Distinguishing between <i>C. jubata</i> and <i>C. selloana</i> is, therefore, difficult and, as such, this measure would require the banning of both species to have any measure of effectiveness.</p> <p>It is difficult to assess whether VMEs present a risk, and therefore when to apply the relevant phytosanitary measure (IPPC, 2017). The ISPM provides a number of elements to consider when assessing risk; distance of movement (shorter distances are a lower risk), complexity of VME structure (more complex is a higher risk), origin and prior use (VMEs in close proximity to vegetation are a higher risk), storage (VMEs stored outside, near vegetation are a higher risk), and intended location or use (VMEs for use in agriculture, forestry, or close proximity to vegetation are a higher risk).</p> <p>In addition, the inspection, cleaning and treatment will normally take place in the exporting country to meet import requirements. However, there are no EU regulations on phytosanitary requirements for imports of VMEs. Therefore, for the measure to be effective, either regulations need to be developed to regulate VME imports, or inspections and phytosanitary measures would need to be applied at EU ports and also at EU/non-EU border facilities.</p>				
<p>Effort required e.g. period of time over which measure need to be applied to have results</p>	<p>This measure would have to be applied indefinitely due to the possibility of viable rootstock being imported, and because this species has a long flowering period (summer to autumn) that together with its occurrence in both Northern and Southern Hemispheres means there would be an all-year round possibility of viable seed being introduced.</p>			
<p>Resources required¹ e.g. cost, staff, equipment etc.</p>	<p>Phytosanitary inspections require trained staff and identification material for accurate identification of <i>C. jubata</i>. Visual identification would require inflorescences to be present (Testoni & Linder, 2017), but other plant material could potentially be identified by barcoding (it is genetically distinguishable from <i>C. selloana</i>; Houlston & Goeke, 2017).</p>			

	<p>Facilities required for the inspection, cleaning, and treatment of VME may include: surfaces that prevent contact with soil, including soil traps and wastewater management systems, temperature treatment facilities, and fumigation or chemical treatment facilities (IPPC, 2017). In addition, trained staff are needed to undertake the inspections and phytosanitary measures, and suitable disposal facilities are required, especially if implemented within the EU.</p>						
<p>Side effects (incl. potential) – both positive and negative i.e. positive or negative side effects of the measure on public health, environment including non-targeted species, etc.</p> <p>For each of the side effect types please select one of the impact categories (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<p>Environmental effects</p>	<p>Positive</p>	<input type="checkbox"/>	<p>Neutral or mixed</p>	<input checked="" type="checkbox"/>	<p>Negative</p>	<input type="checkbox"/>
	<p>Social effects</p>	<p>Positive</p>	<input type="checkbox"/>	<p>Neutral or mixed</p>	<input checked="" type="checkbox"/>	<p>Negative</p>	<input type="checkbox"/>
	<p>Economic effects</p>	<p>Positive</p>	<input checked="" type="checkbox"/>	<p>Neutral or mixed</p>	<input type="checkbox"/>	<p>Negative</p>	<input type="checkbox"/>
	<p><i>Rationale:</i> Inspections would have an economic cost to those undertaking it, which may include both government and the private sector. There would also be economic costs associated with cleaning/treating infected materials, and with any delays in the transport of high risk materials due to inspections.</p>						
	<p>There will be no positive or negative environmental or social side effects to this measure, apart from the cleaning and treatment of high risk VMEs would address additional invasive alien species.</p>						
<p>Acceptability to stakeholders e.g. impacted economic activities, animal welfare considerations, public perception, etc.</p> <p>Please select one of the categories of acceptability (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<p>Acceptability to stakeholders</p>	<p>Acceptable</p>	<input type="checkbox"/>	<p>Neutral or mixed</p>	<input type="checkbox"/>	<p>Unacceptable</p>	<input checked="" type="checkbox"/>
	<p><i>Rationale:</i> Due to the measure being ineffective, it is likely this measure would be seen as unacceptable, especially by those sectors involved in the transport of high risk materials.</p>						
<p>Additional cost information¹ When not already included above, or in the species Risk Assessment.</p> <ul style="list-style-type: none"> - implementation cost for Member States - the cost of inaction - the cost-effectiveness - the socio-economic aspects 	<p>Implementation costs for Member States will be high because of the need for trained staff and long-term implementation of this measure, but figures are not readily available in the public domain.</p> <p>For costs of inaction, see above table, <i>Prevention of intentional introductions and spread</i>.</p> <p>This measure is unlikely to be cost-effective because of the high costs of implementation, but there are no studies specific to <i>C. jubata</i> to support this.</p> <p>There are no known socio-economic aspects.</p>						

Include quantitative &/or qualitative data, and case studies (incl. from countries outside the EU).					
Level of confidence on the information provided ² Please select one of the confidence categories along with a statement to support the category chosen. See <i>Notes</i> section at the bottom of this document. NOTE – this is not related to the effectiveness of the measure	<i>Inconclusive</i>	X	<i>Unresolved</i>	<i>Established but incomplete</i>	<i>Well established</i>
<i>Rationale:</i> There is limited evidence to support or reject the use of phytosanitary measures to prevent unintentional introductions of <i>C. jubata</i> . This is largely because of uncertainty as to whether the species is already present in the EU, whether seed or rootstock is actually transported in these materials/objects, and whether it would be cost effective to implement such a measure.					

Prevention of secondary spread of the species – measures for preventing the species spreading once they have been introduced (cf. Article 13 of the IAS Regulation).	
Measure description Provide a description of the measure, and identify its objective	Removal and destruction of any and all <i>C. jubata</i> plants in gardens and ornamental trials. <i>C. jubata</i> spreads naturally primarily by wind dispersal of its seeds as these are small and light and have long fine hairs that assist with long distance dispersal (Bellgard et al., 2010). Ornamental plantings of the related <i>C. selloana</i> have been found to be the primary sources of invasions in California (Okada, Ahmad & Jasieniuk, 2007). Dispersal distances as great as 50 km have been reported (New Zealand Plant Conservation Network, 2018). Inflorescences of this species can bear over 100,000 seeds (Drewitz & DiTomaso, 2004), making this species extremely fertile. However, on the positive side from a management perspective, seeds generally are viable for only a short time (up to four months under winter (wet) field conditions; Drewitz & DiTomaso, 2004). Seeds may also be transported by water or on animals (Queensland Government, 2018). The removal and destruction of all plants of this species in gardens and ornamental trials will prevent the secondary spread of this species, particularly via wind dispersal. This can only be achieved through preventing their intentional introduction (see <i>Prevention of intentional introductions and spread</i> table), and rapid eradication of established populations (see <i>Rapid eradication for new introductions</i> tables).
Scale of application At what scale is the measure applied? What is the largest scale at which it has been successfully used? Please	See <i>Prevention of intentional introductions and spread</i> , and <i>Rapid eradication for new introductions</i> tables.

provide examples, with areas (km ² or ha) if possible.								
<p>Effectiveness of the measure Is it effective in relation to its objective? Has the measure previously worked, failed?</p> <p>Please select one of the categories of effectiveness (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	Effectiveness of measures	Effective		Neutral		Ineffective		
<p>Effort required e.g. period of time over which measure need to be applied to have results</p>		See <i>Prevention of intentional introductions and spread</i> , and <i>Rapid eradication for new introductions</i> tables.						
<p>Resources required¹ e.g. cost, staff, equipment etc.</p>		See <i>Prevention of intentional introductions and spread</i> , and <i>Rapid eradication for new introductions</i> tables.						
<p>Side effects (incl. potential) – both positive and negative i.e. positive or negative side effects of the measure on public health, environment including non-targeted species, etc.</p>	Environmental effects	Positive		Neutral or mixed		Negative		
		Social effects	Positive		Neutral or mixed		Negative	
		Economic effects	Positive		Neutral or mixed		Negative	
<p>Acceptability to stakeholders e.g. impacted economic activities, animal welfare considerations, public perception, etc.</p> <p>Please select one of the categories of acceptability (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>		Acceptability to stakeholders	Acceptable		Neutral or mixed		Unacceptable	
<p>Additional cost information¹ When not already included above, or in the species Risk Assessment.</p>		See <i>Prevention of intentional introductions and spread</i> , and <i>Rapid eradication for new introductions</i> tables.						

<ul style="list-style-type: none"> - implementation cost for Member States - the cost of inaction - the cost-effectiveness - the socio-economic aspects <p>Include quantitative &/or qualitative data, and case studies (incl. from countries outside the EU).</p>								
<p>Level of confidence on the information provided ²</p> <p>Please select one of the confidence categories along with a statement to support the category chosen. See <i>Notes</i> section at the bottom of this document.</p> <p>NOTE – this is not related to the effectiveness of the measure</p>	<i>Inconclusive</i>		<i>Unresolved</i>		<i>Established but incomplete</i>		<i>Well established</i>	
<p><i>Rationale:</i> See <i>Prevention of intentional introductions and spread</i>, and <i>Rapid eradication for new introductions</i> tables.</p>								

<p>Surveillance measures to support early detection - Measures to run an effective surveillance system for achieving an early detection of a new occurrence (cf. Article 16). This section assumes that the species is not currently present in a Member State, or part of a Member State's territory. This table is repeated for each of the early detection measures identified.</p>	
<p>Measure description Provide a description of the measure, and identify its objective</p>	<p>Survey of known introduction sites and a sample of <i>C. selloana</i> populations</p> <p><i>C. jubata</i> has only been recorded as being introduced into a very limited number of sites in the EU, including very early introductions in the late 1800s at Belgrove, Cork (Ireland. However, these plants were killed by frost; Hooker, 1898), Zurich Botanical Gardens (Switzerland), Nancy (France) (Carrière, 1878), and later in the UK at Dingle Nurseries & Garden and Knoll Gardens (Royal Horticultural Society, 2009).</p> <p><i>C. jubata</i> is easily confused with the closely related <i>C. selloana</i> (Lambrinos, 2001; Robinson, 1984), which is already common across much of the EU (DAISIE European Invasive Alien Species Gateway, 2008). Many of the measures proposed in this note are dependent on knowing whether <i>C. jubata</i> is already present in the EU. Therefore the objective of this measure would be to determine if <i>C. jubata</i> has escaped cultivation and whether perhaps it has been mistaken for <i>C. selloana</i> in areas currently thought to only have <i>C. selloana</i> and not <i>C. jubata</i>.</p>

	<p>The following factors would need to be taken into consideration for designing a survey for the early detection of <i>C. jubata</i>:</p> <ul style="list-style-type: none"> • Seed dispersal of this species has been recorded up to a maximum of 50 km (New Zealand Plant Conservation Network, 2018), although it is likely that seeds of this species, like those of <i>C. selloana</i>, mostly only disperse up to within 40 m of the parent plant (Saura-Mas & Lloret, 2005). • <i>C. jubata</i> invades a wide variety of habitats. It is particularly known for invading disturbed/ruderal areas such as roadsides, logged forests/plantations and recently burnt vegetation (Edgar & Connor, 2000; Parsons & Cuthbertson, 2004; Robinson, 1984; Starr et al., 2003). <p>Therefore, it is advisable that a minimum of a 50 km search radius around the survey sites be included in any surveys (with particular attention being paid to disturbed areas), but extensive field surveys probably only need be close (within a few 100 m) to where the plants were grown. The U.S. Fish & Wildlife Service provides useful guidelines on survey design (https://www.fws.gov/invasives/staffTrainingModule/assessing/inventory.html).</p> <p>In Maui, aerial surveys have been suggested as an effective method of detecting flowering mature plants in difficult to reach locations (Starr et al., 2003), and this could be considered for conducting the broader survey (paying attention to the flowering time of this species). Alternatively, satellite remote sensing imagery has been successfully used to detect large <i>C. jubata</i> plants in California (Underwood, Ustin & DiPietro, 2003); although this may not be as useful for early detection, as detected plants would probably already have been reproducing for a number of years before their detection. Moreover, the difficulty in distinguishing between <i>C. jubata</i> and <i>C. selloana</i> might make this impractical.</p> <p>The nurseries in the UK at which this species were recently grown may have kept the plants in greenhouses. This would have limited the chances of wind dispersal of this species and it would therefore only be necessary to enquire whether plants were disposed of in a manner that would prevent their spread via rootstock. If spread via rootstock was possible, locations at which plant material was disposed of would need to be inspected, but if not, then no further surveys will be required.</p>
<p>Scale of application At what scale is the measure applied? What is the largest scale at which it has been successfully used? Please provide examples, with areas (km² or ha) if possible.</p>	<p>This measure would need to be applied at the four known introduction sites of this species, within a search radius of at least 50 km of these locations. In addition, a sample of <i>C. selloana</i> populations across the EU should similarly be surveyed. Published surveillance strategies are available and would require a more in-depth analysis to design an optimal surveillance strategy (Epanchin-Niell et al., 2012; Hauser & McCarthy, 2009).</p>

<p>Effectiveness of the measure Is it effective in relation to its objective? Has the measure previously worked, failed?</p> <p>Please select one of the categories of effectiveness (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<table border="1"> <tr> <td><i>Effectiveness of measures</i></td> <td><i>Effective</i></td> <td></td> <td><i>Neutral</i></td> <td>X</td> <td><i>Ineffective</i></td> <td></td> </tr> </table>	<i>Effectiveness of measures</i>	<i>Effective</i>		<i>Neutral</i>	X	<i>Ineffective</i>															
<i>Effectiveness of measures</i>	<i>Effective</i>		<i>Neutral</i>	X	<i>Ineffective</i>																	
<p>Effort required e.g. period of time over which measure need to be applied to have results</p>	<p><i>Rationale:</i> <i>C. jubata</i> has a very distinctive growth form and is also a large plant, both of which make it an easy to detect species. In Maui, roadside and aerial surveys have been successfully used for early detection of both <i>C. jubata</i> and <i>C. selloana</i> (Penniman et al., 2011; Starr et al., 2003).</p> <p>However, the difficulty in distinguishing between <i>C. jubata</i> and <i>C. selloana</i> reduces the effectiveness of this measure. Surveyors will be required to be very familiar with the morphological differences between these two species, or be able to use DNA barcoding, in order to ensure this measure is effective.</p> <p>The only figures available on the search effort for this species come from Maui where 24,425 acres were surveyed over a period of 3,063 hours in 2012, with 3,910 plants being detected and controlled (HISC Established Pests Working Group, 2013). However, one cannot calculate survey effort from these figures because the total time includes survey and control efforts.</p> <p>Surveys would need to be conducted during the flowering period of <i>C. jubata</i> (i.e., mid-summer to autumn) to ensure accurate identification and easy detectability of the species. Based on the phenology of <i>C. jubata</i> and <i>C. selloana</i> elsewhere in the world, particularly in California, <i>C. jubata</i> could be distinguished from <i>C. selloana</i> if flowers are already present in July. However, later in summer (August onwards) there is likely to be overlap in the flowering period of these two species (DiTomaso et al., 2010).</p>																					
<p>Resources required¹ e.g. cost, staff, equipment etc.</p>	<p>Surveying of these sites will require trained professionals with knowledge of the identification of <i>C. jubata</i> and <i>C. selloana</i>. For the identification of <i>C. jubata</i> and <i>C. selloana</i>, easy-to-use identification guides (e.g., Cal-IPC, 2018; DiTomaso et al., 2010; Testoni & Linder, 2017), or DNA barcoding (Houliston & Goeke, 2017) will be required. Surveyors will need access to vehicles to explore the survey area. If aerial surveys are to be conducted, the use of a helicopter or UAV (drone) will be needed. There are no published guidelines for the number of surveyors needed to effectively detect <i>C. jubata</i> plants within any given area.</p>																					
<p>Side effects (incl. potential) – both positive and negative i.e. positive or negative side effects of the measure on public health, environment including non-targeted species, etc.</p> <p>For each of the side effect types please select one of the impact categories (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<table border="1"> <tr> <td><i>Environmental effects</i></td> <td><i>Positive</i></td> <td></td> <td><i>Neutral or mixed</i></td> <td>X</td> <td><i>Negative</i></td> <td></td> </tr> <tr> <td><i>Social effects</i></td> <td><i>Positive</i></td> <td></td> <td><i>Neutral or mixed</i></td> <td></td> <td><i>Negative</i></td> <td>X</td> </tr> <tr> <td><i>Economic effects</i></td> <td><i>Positive</i></td> <td>X</td> <td><i>Neutral or mixed</i></td> <td></td> <td><i>Negative</i></td> <td></td> </tr> </table> <p><i>Rationale:</i> There could be negative social (incl. safety) side effects for surveyors if surveying needs to be conducted in difficult to reach locations (e.g., cliff faces).</p> <p>Surveys could provide positive side effects through the creation of employment opportunities.</p>	<i>Environmental effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>	X	<i>Negative</i>		<i>Social effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>		<i>Negative</i>	X	<i>Economic effects</i>	<i>Positive</i>	X	<i>Neutral or mixed</i>		<i>Negative</i>	
<i>Environmental effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>	X	<i>Negative</i>																	
<i>Social effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>		<i>Negative</i>	X																
<i>Economic effects</i>	<i>Positive</i>	X	<i>Neutral or mixed</i>		<i>Negative</i>																	

<p>Acceptability to stakeholders e.g. impacted economic activities, animal welfare considerations, public perception, etc.</p> <p>Please select one of the categories of acceptability (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<table border="1" data-bbox="613 153 1854 220"> <tr> <td><i>Acceptability to stakeholders</i></td> <td><i>Acceptable</i></td> <td></td> <td><i>Neutral or mixed</i></td> <td>X</td> <td><i>Unacceptable</i></td> <td></td> </tr> </table> <p><i>Rationale:</i> Surveys will almost certainly need to be conducted on privately owned land, and this may not be acceptable to the relevant landowners. Moreover, there may be resistance on the part of any landowners found to possess this species to have it removed. On Catalina Island in California, landowners appear more willing to have a species removed if a native species is offered in replacement (Parish, 2017).</p>	<i>Acceptability to stakeholders</i>	<i>Acceptable</i>		<i>Neutral or mixed</i>	X	<i>Unacceptable</i>		
<i>Acceptability to stakeholders</i>	<i>Acceptable</i>		<i>Neutral or mixed</i>	X	<i>Unacceptable</i>				
<p>Additional cost information¹ When not already included above, or in the species Risk Assessment.</p> <ul style="list-style-type: none"> - implementation cost for Member States - the cost of inaction - the cost-effectiveness - the socio-economic aspects <p>Include quantitative &/or qualitative data, and case studies (incl. from countries outside the EU).</p>	<p>Implementation costs for such a survey are not readily available in the scientific literature. The budget for early detection roadside surveys of all invasive species in Hawai'i was estimated at USD 100,000 per year per county (i.e. USD 500,000 for all counties; Hawai'i Department of Transportation, 2011). Given a total road length of ~15,500 km in Hawai'i (U.S. Department of Transportation Federal Highway Administration, 2013), this equates to ~USD 32.30 (ca. EUR 28.2) per km surveyed.</p> <p>Costs of inaction associated with this measure are likely to be similar to those detailed in the <i>Prevention of intentional introductions and spread</i> table above, if this species escapes from gardens or field trials.</p> <p>To determine the cost effectiveness of surveying for <i>C. jubata</i> around the known introduction sites, one would need to conduct a formal analysis using knowledge of this species, e.g., its detectability, probability of occurring in the landscape, potential benefits of detection and eradication, population establishment and growth rates, and costs of surveying and management (Epanchin-Niell et al., 2012; Hauser & McCarthy, 2009). However, based on general findings from studies of survey cost-benefits, it seems likely that given the high detectability of <i>C. jubata</i> and the high potential costs of invasions, surveys would be cost effective.</p> <p>Socio-economic aspects include the potential loss of revenue to the two nurseries that have been growing <i>C. jubata</i>, but this seems minor given that neither nursery selected this species for further commercialisation.</p>								
<p>Level of confidence on the information provided²</p> <p>Please select one of the confidence categories along with a statement to support the category chosen. See <i>Notes</i> section at the bottom of this document.</p> <p>NOTE – this is not related to the effectiveness of the measure</p>	<table border="1" data-bbox="613 1011 1895 1078"> <tr> <td><i>Inconclusive</i></td> <td></td> <td><i>Unresolved</i></td> <td></td> <td><i>Established but incomplete</i></td> <td>X</td> <td><i>Well established</i></td> <td></td> </tr> </table> <p><i>Rationale:</i> There is a substantial amount of evidence to suggest that <i>C. jubata</i> was only trialled at a limited number of locations in the EU. However, there is always the potential that there are more localities that are not reported in the scientific or grey literature and that <i>C. jubata</i> has been introduced under the false assumption that it is <i>C. selloana</i>.</p>	<i>Inconclusive</i>		<i>Unresolved</i>		<i>Established but incomplete</i>	X	<i>Well established</i>	
<i>Inconclusive</i>		<i>Unresolved</i>		<i>Established but incomplete</i>	X	<i>Well established</i>			

Surveillance measures to support early detection - Measures to run an effective surveillance system for achieving an early detection of a new occurrence (cf. Article 16). This section assumes that the species is not currently present in a Member State, or part of a Member State's territory. **This table is repeated for each of the early detection measures identified.**

<p>Measure description Provide a description of the measure, and identify its objective</p>	<p>Use citizen science and resource managers' data for early detection</p> <p>The objectives of this measure would be (1) to promote <i>C. jubata</i> as a target for identification to invasive species citizen science platforms, and (2) to provide citizen scientists and resource managers with the knowledge to identify this species and thereby support its early detection.</p> <p>Citizen science locality data has been shown to be very useful for the early detection of invasive species (Gallo & Waitt, 2011; Maistrello et al., 2016). Numerous such databases currently exist, including EASIN (https://easin.jrc.ec.europa.eu/), which is the official EU platform for reporting alien species occurrences and the accompanying smart phone application. However, <i>C. jubata</i> does not currently feature in EASIN. There are also a number of other European and national IAS awareness and citizen science IAS monitoring programs into which <i>C. jubata</i> could be incorporated, which are important resources for environmental agencies, resource managers and decision makers (Genovesi et al., 2010).</p>						
<p>Scale of application At what scale is the measure applied? What is the largest scale at which it has been successfully used? Please provide examples, with areas (km² or ha) if possible.</p>	<p>This measure would need to be applied across the EU, but countries/regions with high climatic suitability should be particularly prioritised (see EPPO, 2018 for details).</p>						
<p>Effectiveness of the measure Is it effective in relation to its objective? Has the measure previously worked, failed? Please select one of the categories of effectiveness (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<p><i>Effectiveness of measures</i></p>	<p><i>Effective</i></p>		<p><i>Neutral</i></p>		<p><i>Ineffective</i></p>	<p>X</p>
<p>Effort required e.g. period of time over which measure need to be applied to have results</p>	<p>Citizen science programs need good quality assessment of the data collected, well designed and standardised methods of data collection, an explicit goal or hypothesis (i.e., in this case, the early detection of <i>C. jubata</i>), feedback to participants on their contributions as a reward for their participation (Silvertown, 2009). One area that would require significant effort would be the peer review of data to ensure that any <i>C. jubata</i> records are not <i>C. selloana</i>.</p>						

<p>Resources required ¹ e.g. cost, staff, equipment etc.</p>	<p>This measure will require a well-designed and supported citizen science platform and ideally smart phone application. The use of EASIN and established national systems for this purpose is possible, but the promotion of recording <i>C. jubata</i> will be required.</p> <p>Accurate identification of <i>C. jubata</i> will need to be supported through easy-to-use identification guides (e.g., Cal-IPC, 2018; DiTomaso et al., 2010; Testoni & Linder, 2017), as <i>C. jubata</i> is easily confused with the closely related <i>C. selloana</i> (Lambrinos, 2001; Robinson, 1984), which is already common across much of the EU (DAISIE European Invasive Alien Species Gateway, 2008). New records of <i>C. jubata</i> identified by citizen scientists and invasive species managers will need to be verified by taxonomists or DNA barcoding, due to the difficulty of distinguishing <i>C. jubata</i> and <i>C. selloana</i>.</p>						
<p>Side effects (incl. potential) – both positive and negative i.e. positive or negative side effects of the measure on public health, environment including non-targeted species, etc.</p> <p>For each of the side effect types please select one of the impact categories (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<p>Environmental effects</p>	<p>Positive</p>	<p>X</p>	<p>Neutral or mixed</p>		<p>Negative</p>	
<p>Social effects</p>	<p>Positive</p>	<p>X</p>	<p>Neutral or mixed</p>		<p>Negative</p>		
<p>Economic effects</p>	<p>Positive</p>		<p>Neutral or mixed</p>	<p>X</p>	<p>Negative</p>		
<p>Acceptability to stakeholders e.g. impacted economic activities, animal welfare considerations, public perception, etc.</p> <p>Please select one of the categories of acceptability (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<p>Acceptability to stakeholders</p>	<p>Acceptable</p>	<p>X</p>	<p>Neutral or mixed</p>		<p>Unacceptable</p>	
<p>Additional cost information ¹ When not already included above, or in the species Risk Assessment. - implementation cost for Member States - the cost of inaction</p>	<p>Implementation costs of setting up and running citizen science databases and awareness raising programs are large (Genovesi et al., 2010), but databases and programs like these are already running and funded by the EU (e.g., EASIN) and individual Member States. Therefore additional costs for promoting the collection of <i>C. jubata</i> records, and raising awareness of this species, are likely to be minimal.</p> <p>Costs of inaction associated with this measure are likely to be similar to those detailed in the <i>Prevention of intentional introductions and spread</i> table above, if this species escapes.</p>						

<p>- the cost-effectiveness - the socio-economic aspects</p> <p>Include quantitative &/or qualitative data, and case studies (incl. from countries outside the EU).</p>	<p>Cost-effectiveness of citizen science programs is well established and justified elsewhere (e.g., Gallo et al., 2011; Genovesi et al., 2010; Maistrello et al., 2016; Tulloch et al., 2013).</p> <p>There are no known additional socio-economic aspects to consider.</p>							
<p>Level of confidence on the information provided²</p> <p>Please select one of the confidence categories along with a statement to support the category chosen. See <i>Notes</i> section at the bottom of this document.</p> <p>NOTE – this is not related to the effectiveness of the measure</p>	<p><i>Inconclusive</i></p>		<p><i>Unresolved</i></p>		<p><i>Established but incomplete</i></p>		<p><i>Well established</i></p>	<p>X</p>
<p><i>Rationale:</i> There is considerable evidence to support the use of citizen science for early detection of invasive species (see <i>Measure description</i>).</p>								

<p>Rapid eradication for new introductions - Measures to achieve eradication <u>at an early stage of invasion</u>, after an early detection of a new occurrence (cf. Article 17). This section assumes that the species is not currently present in a Member State, or part of a Member State's territory. This table is repeated for each of the eradication measures identified.</p>	
<p>Measure description Provide a description of the measure, and identify its objective</p>	<p>There are no recommended measures for rapid eradication of new introductions specific to <i>C. jubata</i>, or to <i>Cortaderia</i> species in general. Moreover, given that this species is thought not to have established within the EU, these measures will largely be the same as for management (section below).</p> <p>Physical control</p> <p>Small seedlings can be pulled by hand, but larger plants will require digging up by spade, axe, hoe, having a chain tied around them and pulled out by a vehicle, or even dug up with the use of a digger (DiTomaso et al., 2010; Gosling et al., 2000).</p>
<p>Scale of application At what scale is the measure applied? What is the largest scale at which it has been successfully used? Please provide examples, with areas (km² or ha) if possible.</p>	<p>There are no specific recommendations for the largest scale at which physical control is possible. However, numerous authors note that physical control is only practical and economical for small invasions of a few plants (DiTomaso et al., 2010; Gosling et al., 2000; Schmalzer & Hinkle, 1987). This suggests physical control is suitable for rapid eradication for new introductions.</p> <p>On Moloka'i in Hawai'i, <i>C. jubata</i> was successfully eradicated from an 11 ha area over a period of seven years of control and monitoring (Penniman et al., 2011), probably using only physical control as there were only six plants.</p>

<p>Effectiveness of the measure Is it effective in relation to its objective? Has the measure previously worked, failed?</p> <p>Please select one of the categories of effectiveness (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<p>Effectiveness of measures</p>	<p>Effective</p>	<p>X</p>	<p>Neutral</p>		<p>Ineffective</p>	
	<p><i>Rationale:</i> Physical control is an extremely effective method of controlling <i>C. jubata</i>, with a 98-100% success rate (DiTomaso et al., 2008), although it can disturb surrounding vegetation and encourage the re-establishment of <i>C. jubata</i> seedlings (Gosling et al., 2000). Other considerations to improve the effectiveness of the measure include removing all the inflorescences and destroying these to prevent seed dispersal, and pulling up the majority of roots to prevent resprouting from rootstock (DiTomaso et al., 2010).</p> <p>This measure can only be effective if <i>C. jubata</i> can be accurately distinguished from individuals of <i>C. selloana</i>. However, physical control of both species would ensure the effectiveness of this measure.</p>						
<p>Effort required e.g. period of time over which measure need to be applied to have results</p>	<p>It is not known exactly how long physical control would need to be implemented to ensure a successful eradication. Managers generally declare a species eradicated after five years of no detections, although there are more quantitative methods of determining the optimal amount of time to continue searching for a species before declaring a successful eradication (e.g., Regan et al., 2006). <i>C. jubata</i> has a short-lived seedbank under natural conditions (Drewitz & DiTomaso, 2004), and therefore eradication is possible within a time period as short as a year as long as all reproducing individuals of <i>C. jubata</i> are located and removed. However, an eradication program in Hawaii was only declared successful after seven years of monitoring (Penniman et al., 2011).</p>						
<p>Resources required¹ e.g. cost, staff, equipment etc.</p>	<p>Depending on the size of plants, different resources will be required. All plant sizes will require manual labour. Small seedlings can be removed by hand pulling, so labour is probably the only resource required for follow-up treatments of <i>C. jubata</i> invasions. Medium-sized plants will require tools (spades, axes or hoes) and even larger plants will require machinery (vehicles and chains or diggers).</p> <p>There are almost no cost estimates specific to <i>C. jubata</i> (or even for <i>C. selloana</i>) available for these methods. In New Zealand physical control was estimated to cost NZD 150 p.ha⁻¹ in 1983 (ca. USD 325 / EUR 283 p.ha⁻¹ at current prices) (Gadgil et al., 1984).</p>						
<p>Side effects (incl. potential) – both positive and negative i.e. positive or negative side effects of the measure on public health, environment including non-targeted species, etc.</p> <p>For each of the side effect types please select one of the impact</p>	<p>Environmental effects</p>	<p>Positive</p>	<p></p>	<p>Neutral or mixed</p>	<p></p>	<p>Negative</p>	<p>X</p>
	<p>Social effects</p>	<p>Positive</p>	<p>X</p>	<p>Neutral or mixed</p>	<p></p>	<p>Negative</p>	<p></p>
	<p>Economic effects</p>	<p>Positive</p>	<p>X</p>	<p>Neutral or mixed</p>	<p></p>	<p>Negative</p>	<p></p>
	<p><i>Rationale:</i> This measure can have negative environmental effects in that physical control using machinery can damage surrounding vegetation, and it can encourage germination of <i>C. jubata</i> seeds (Gosling et al., 2000).</p>						

<p>categories (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>						
<p>Acceptability to stakeholders e.g. impacted economic activities, animal welfare considerations, public perception, etc.</p> <p>Please select one of the categories of acceptability (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<p>Acceptability to stakeholders</p>	<p>Acceptable</p>	<p>Neutral or mixed</p>	<p>X</p>	<p>Unacceptable</p>	
<p>Additional cost information¹ When not already included above, or in the species Risk Assessment.</p> <ul style="list-style-type: none"> - implementation cost for Member States - the cost of inaction - the cost-effectiveness - the socio-economic aspects <p>Include quantitative &/or qualitative data, and case studies (incl. from countries outside the EU).</p>	<p>Implementation costs are uncertain, but see best estimates from New Zealand above.</p> <p>Costs of inaction associated with this measure are likely to be similar to those detailed in the <i>Prevention of intentional introductions and spread</i> table above, if this species escapes.</p> <p>The cost-effectiveness of physical control for rapid eradications of this species is unknown, but probably high due to the efficacy of the method and the high costs of inaction. For invasive plants in general, the economic effects are likely to be positive overall because the potential impacts of an invasive species outweigh the costs of its control at the early stages of an invasion (e.g., Leung et al., 2002).</p> <p>There are no known socio-economic aspects.</p>					
<p>Level of confidence on the information provided²</p> <p>Please select one of the confidence categories along with a statement to support the category chosen. See <i>Notes</i> section at the bottom of this document.</p> <p>NOTE – this is not related to the effectiveness of the measure</p>	<p>Inconclusive</p>	<p>Unresolved</p>	<p>Established but incomplete</p>	<p>X</p>	<p>Well established</p>	
	<p><i>Rationale:</i> There is adequate evidence to support the use of physical control to eradicate new introductions of this species. However, there is no readily available information on the costs of using such an approach.</p>					

Rapid eradication for new introductions - Measures to achieve eradication at an early stage of invasion, after an early detection of a new occurrence (cf. Article 17). This section assumes that the species is not currently present in a Member State, or part of a Member State's territory. **This table is repeated for each of the eradication measures identified.**

<p>Measure description Provide a description of the measure, and identify its objective</p>	<p>Chemical control</p> <p>The use of glyphosate, imazapyr, fluazifop-P-butyl, quizalofop, hexazinone, amitrole, dalapon, and sethoxydim have either been trialled or recommended for use on <i>C. jubata</i> (DiTomaso et al., 2008; Gosling et al., 2000; Schmalzer & Hinkle, 1987). However, only a few of these have been tested using experimental field trials (DiTomaso et al., 2008). Glyphosate is generally the recommended chemical of choice for controlling <i>C. jubata</i> (DiTomaso et al., 2008, 2010; Gosling et al., 2000).</p> <p>It is important to note that EU/national/local legislation on the use of plant protection products and biocides needs to be respected and authorities should check to ensure chemicals are licensed for use in their respective countries/regions.</p>						
<p>Scale of application At what scale is the measure applied? What is the largest scale at which it has been successfully used? Please provide examples, with areas (km² or ha) if possible.</p>	<p>In the case of rapid eradications, chemical control is the only practical control method for difficult to reach invasions of <i>C. jubata</i> (Popay et al., 2003), and perhaps also for larger plants for which physical removal is not practical (DiTomaso et al., 2008). Rapid eradications of <i>C. jubata</i> are probably only feasible for smaller invasions (lower numbers of plants within a relatively small area) using chemical control, but it is difficult to provide specific numbers. On Tawhiti Rahi Island and Aorangi Island in New Zealand, both <i>C. jubata</i> and <i>C. selloana</i> were largely eradicated using physical and chemical control from a 272 ha area (only one detection per year in the last two years of a seven year eradication and monitoring program) (Coulston, 2002).</p>						
<p>Effectiveness of the measure Is it effective in relation to its objective? Has the measure previously worked, failed?</p> <p>Please select one of the categories of effectiveness (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<p><i>Effectiveness of measures</i></p>	<p><i>Effective</i></p>		<p><i>Neutral</i></p>	<p>X</p>	<p><i>Ineffective</i></p>	
<p><i>Rationale:</i> Chemical control has been shown to be highly effective under certain circumstances. In a Californian trial, spraying of glyphosate at an 8% concentration in early summer was found to be the most cost-effective control method and had a control rate of 99% (however, ropewick application of glyphosate at a 16% concentration in early summer was better for not affecting non-target native species; DiTomaso et al., 2008). In general, glyphosate had a much higher successful control rate than imazapyr, fluazifop and sethoxydim (DiTomaso et al., 2008). The more commonly used spray-to-wet applications of glyphosate (at high concentrations) were as effective as ropewick applications of glyphosate, but even at low concentrations spray-to-wet applications of glyphosate were still more effective than the other herbicides tested (DiTomaso et al., 2008). Larger plants appear to be more likely to survive herbicide control than smaller plants (DiTomaso et al., 2008; Popay et al., 2003).</p> <p>Hexazinone, a root-absorbed herbicide, has been recommended in forestry plantations in California and New Zealand (DiTomaso et al., 2010; Gosling et al., 2000).</p> <p>Aerial spraying of haloxyfop from a beer keg from a helicopter has been shown to quite effective at reducing <i>C. jubata</i> growth, but not entirely effective at killing plants. A year after treatment there were numerous new shoots on treated plants and it appears that follow-up treatments are necessary (Popay et al., 2003).</p>							

	<p>This measure can only be effective if <i>C. jubata</i> can be accurately distinguished from individuals of <i>C. seloana</i>. However, chemical control of both species would ensure the effectiveness of this measure.</p>						
<p>Effort required e.g. period of time over which measure need to be applied to have results</p>	<p>If applied at the correct concentrations, and when the plants are not stressed, chemical control can be very effective, and follow-up treatments can be limited in number and duration, but there are no experimental trials to provide exact recommendations. However, providing high efficacy of herbicide treatments, follow-up treatments will likely only need to focus on the few surviving adult plants, and only on new recruits within the first year after initial treatment as <i>C. jubata</i> seeds do not persist for longer than four months in the soil seedbank under winter (wet) conditions (Drewitz & DiTomaso, 2004). In reality herbicide control is not 100% effective and an eradication program in Hawai'i was only deemed successful after seven years (Penniman et al., 2011).</p>						
<p>Resources required¹ e.g. cost, staff, equipment etc.</p>	<p>Herbicide application requires trained staff, equipment (e.g., backpack sprayers, ropewicks, spray wands, or in the case of aerial application, a helicopter fitted with a suitable spray device, e.g., beer keg), herbicides and surfactants.</p> <p>The following studies provided cost estimates (all inclusive) of controlling <i>C. jubata</i>:</p> <ul style="list-style-type: none"> • In California, the most cost effective control method for <i>C. jubata</i> was found to be spray-to-wet application of glyphosate, which cost USD 0.28 per 1 m² plant (USD 2,800 / ca. EUR 2441 per hectare; DiTomaso et al., 2008). • Ropewick application of glyphosate, a method less likely to affect non-target plants, was estimated to cost USD 0.80 per 1 m² plant (USD 8,000 / ca. EUR 6,975 per hectare; DiTomaso et al., 2008). • In New Zealand the cheapest herbicide (haloxyfop) control method using helicopters for inaccessible <i>C. jubata</i> infestations was found to cost NZD 3,885 (ca. USD 2,500 / EUR 2,219 per hectare). 						
<p>Side effects (incl. potential) – both positive and negative i.e. positive or negative side effects of the measure on public health, environment including non-targeted species, etc.</p> <p>For each of the side effect types please select one of the impact categories (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<p>Environmental effects</p>	<p>Positive</p>		<p>Neutral or mixed</p>		<p>Negative</p>	<p>X</p>
	<p>Social effects</p>	<p>Positive</p>		<p>Neutral or mixed</p>		<p>Negative</p>	<p>X</p>
	<p>Economic effects</p>	<p>Positive</p>		<p>Neutral or mixed</p>	<p>X</p>	<p>Negative</p>	
	<p><i>Rationale:</i> This measure can have negative environmental effects in that chemical control can also affect native species (DiTomaso et al., 2008; Gosling et al., 2000). Glyphosate is a commonly recommended broad-spectrum herbicide recommended for <i>C. jubata</i> control, which can also affect neighbouring native plants when applied via sprayers (DiTomaso et al., 2008; Gosling et al., 2000). For this reason, ropewick application of glyphosate has been recommended (DiTomaso et al., 2008) and is probably a very practical method for controlling a few plants in a new introduction. Hexazinone, which has been used to treat <i>C. jubata</i> in plantation forests, can affect native plants in light or sandy soils (Gosling et al., 2000). Haloxyfop is a grass-specific herbicide that has been used in aerial spraying of <i>C. jubata</i> in New Zealand and was found to damage the native <i>Austroderia splendens</i> (a close relative to <i>Cortaderia</i> species), but not other native plants, but was also not completely effective at killing <i>C. jubata</i> (Popay et al., 2003).</p> <p>Negative social side effects can include exposure to toxic substances if adequate precautions are not taken.</p>						

<p>Acceptability to stakeholders e.g. impacted economic activities, animal welfare considerations, public perception, etc.</p> <p>Please select one of the categories of acceptability (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<table border="1"> <tr> <td>Acceptability to stakeholders</td> <td><i>Acceptable</i></td> <td></td> <td><i>Neutral or mixed</i></td> <td>X</td> <td><i>Unacceptable</i></td> <td></td> </tr> </table>	Acceptability to stakeholders	<i>Acceptable</i>		<i>Neutral or mixed</i>	X	<i>Unacceptable</i>		<p><i>Rationale:</i> Invasive species managers are likely to favour chemical control because of its ease of use compared to physical control. However, public perceptions of chemical control are often negative (e.g., Shindler, Gordon, Brunson & Olsen, 2011). Moreover, chemical control is not always possible or permitted in conservation areas or in riparian areas and wetlands.</p>	
Acceptability to stakeholders	<i>Acceptable</i>		<i>Neutral or mixed</i>	X	<i>Unacceptable</i>					
<p>Additional cost information¹ When not already included above, or in the species Risk Assessment.</p> <ul style="list-style-type: none"> - implementation cost for Member States - the cost of inaction - the cost-effectiveness - the socio-economic aspects <p>Include quantitative &/or qualitative data, and case studies (incl. from countries outside the EU).</p>	<p>The best estimates of implementation costs in the <i>Resources required</i> section above.</p> <p>Costs of inaction associated with this measure are likely to be similar to those detailed in the <i>Prevention of intentional introductions and spread</i> table above, if this species escapes.</p> <p>The cost-effectiveness of physical control for rapid eradications of this species is unknown, but probably high due to the efficacy of the method and the high costs of inaction. For invasive plants in general, the economic effects are likely to be positive overall because the potential impacts of an invasive species outweigh the costs of its control at the early stages of an invasion (e.g., Leung et al., 2002).</p> <p>There are no known socio-economic aspects.</p>									
<p>Level of confidence on the information provided²</p> <p>Please select one of the confidence categories along with a statement to support the category chosen. See <i>Notes</i> section at the bottom of this document.</p> <p>NOTE – this is not related to the effectiveness of the measure</p>	<table border="1"> <tr> <td><i>Inconclusive</i></td> <td></td> <td><i>Unresolved</i></td> <td></td> <td><i>Established but incomplete</i></td> <td>X</td> <td><i>Well established</i></td> <td></td> </tr> </table>	<i>Inconclusive</i>		<i>Unresolved</i>		<i>Established but incomplete</i>	X	<i>Well established</i>		<p><i>Rationale:</i> There is adequate evidence to support the use of chemical control to eradicate new introductions of this species. However, there are insufficient studies providing guidelines on the duration and number of follow-up treatments required. Moreover, there are no cost-benefit analyses on the use of chemical control of <i>C. jubata</i>.</p>
<i>Inconclusive</i>		<i>Unresolved</i>		<i>Established but incomplete</i>	X	<i>Well established</i>				

Management - Measures to achieve management of the species once it has become widely spread within a Member State, or part of a Member State's territory. (cf. Article 19), i.e. **not** at an early stage of invasion (see Rapid eradication table above). These measures can be aimed at eradication, population control or containment of a population of the species. **This table is repeated for each of the management measures identified.**

<p>Measure description Provide a description of the measure, and identify its objective</p>	<p>Management measures overlap to a large extent with measures for rapid eradication for new introductions, as described in the section above. Probably the largest difference between measures in these two different sections is the scale at which control methods are applied. While both physical and chemical control are possible for both rapid eradication of new introductions and management, physical control is only practical and cost-effective for smaller invasions, while chemical control can be used for both small and large invasions (DiTomaso et al., 2008; Gosling et al., 2000; Schmalzer & Hinkle, 1987).</p> <p>Given that <i>C. jubata</i> is thought not to have established within the EU, management measures are probably not necessary and are provided here for completeness. One consideration though is the similarity in appearance of <i>C. jubata</i> to <i>C. selloana</i>, and the abundance and wide distribution of <i>C. selloana</i> in the EU. This may have prevented the accurate identification of <i>C. jubata</i> in the EU and it is plausible that the populations of <i>C. selloana</i> in the EU are in fact <i>C. jubata</i> and that this species is indeed much more widespread in the EU. This would therefore require the implementation of management measures.</p> <p>Physical control</p> <p>Small seedlings can be pulled by hand, but larger plants will require digging up by spade, axe, hoe, having a chain tied around them and pulled out by a vehicle, or even dug up with the use of a digger (DiTomaso et al., 2010; Gosling et al., 2000).</p>							
<p>Scale of application At what scale is the measure applied? What is the largest scale at which it has been successfully used? Please provide examples, with areas (km² or ha) if possible.</p>	<p>Numerous authors note that physical control is only practical and economical for small invasions of a few plants (DiTomaso et al., 2010; Gosling et al., 2000). This suggests physical control may not always be possible for management, and may require the additional use (integrated management) of chemical control as has been done in Hawaii (Penniman et al., 2011) and New Zealand (Gosling et al., 2000).</p>							
<p>Effectiveness of the measure Is it effective in relation to its objective? Has the measure previously worked, failed?</p> <p>Please select one of the categories of effectiveness (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<table border="1" data-bbox="613 951 1854 1011"> <tr> <td data-bbox="613 951 893 1011"><i>Effectiveness of measures</i></td> <td data-bbox="900 951 1120 1011"><i>Effective</i></td> <td data-bbox="1126 951 1176 1011"></td> <td data-bbox="1182 951 1476 1011"><i>Neutral</i></td> <td data-bbox="1482 951 1532 1011"></td> <td data-bbox="1538 951 1800 1011"><i>Ineffective</i></td> <td data-bbox="1807 951 1854 1011">X</td> </tr> </table> <p><i>Rationale:</i> Physical control is an extremely effective method of controlling <i>C. jubata</i>, with a 98-100% success rate (DiTomaso et al., 2008), but is only practical for smaller invasions (DiTomaso et al., 2008; Gosling et al. 2000; Schmalzer & Hinkle, 1987) and is therefore assessed as ineffective for managing large scale invasions.</p> <p>Other considerations to improve the effectiveness of the measure include removing all the inflorescences and destroying these to prevent seed dispersal, and pulling up the majority of roots to prevent resprouting from rootstock (DiTomaso et al., 2010).</p> <p>This measure can only be effective if <i>C. jubata</i> can be accurately distinguished from individuals of <i>C. selloana</i>. However, physical control of both species would ensure the effectiveness of this measure.</p>	<i>Effectiveness of measures</i>	<i>Effective</i>		<i>Neutral</i>		<i>Ineffective</i>	X
<i>Effectiveness of measures</i>	<i>Effective</i>		<i>Neutral</i>		<i>Ineffective</i>	X		

<p>Effort required e.g. period of time over which measure need to be applied to have results</p>	<p>As physical control is extremely labour intensive, considerable manpower will be needed to manage larger invasions of <i>C. jubata</i>, which may be prohibitively expensive or impractical. As a result, the amount of time required to manage invasions will also be considerably longer than using chemical control. However, the duration of monitoring post-treatment will be the same regardless of the size of invasion because physical control is extremely effective, with a 98-100% success rate (DiTomaso et al., 2008), and <i>C. jubata</i> has a very short-lived seedbank (up to four months in winter (wet) conditions; Drewitz & DiTomaso, 2004).</p> <p>Therefore eradication is possible within a time period as short as a year as long as all reproducing individuals of <i>C. jubata</i> are located and removed. However, an eradication program in Hawaii was only declared successful after seven years of monitoring (Penniman et al., 2011).</p>																										
<p>Resources required¹ e.g. cost, staff, equipment etc.</p>	<p>Depending on the size of plants, different resources will be required. All plant sizes will require manual labour. Small seedlings can be removed by hand pulling, so labour is probably the only resource required for follow-up treatments of <i>C. jubata</i> invasions. Medium-sized plants will require tools (spades, axes or hoes) and even larger plants will require machinery (vehicles and chains or diggers).</p> <p>There are almost no specific cost estimates for <i>C. jubata</i> available for these methods. In New Zealand physical control was estimated to cost NZD 150 p.ha⁻¹ in 1983 (USD 325 / ca. EUR 283 p.ha⁻¹ at current prices) (Gadgil et al., 1984).</p>																										
<p>Side effects (incl. potential) – both positive and negative i.e. positive or negative side effects of the measure on public health, environment including non-targeted species, etc.</p> <p>For each of the side effect types please select one of the impact categories (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<table border="1"> <tr> <td>Environmental effects</td> <td>Positive</td> <td>X</td> <td>Neutral or mixed</td> <td></td> <td>Negative</td> <td></td> </tr> <tr> <td>Social effects</td> <td>Positive</td> <td></td> <td>Neutral or mixed</td> <td>X</td> <td>Negative</td> <td></td> </tr> <tr> <td>Economic effects</td> <td>Positive</td> <td></td> <td>Neutral or mixed</td> <td></td> <td>Negative</td> <td>X</td> </tr> </table>	Environmental effects	Positive	X	Neutral or mixed		Negative		Social effects	Positive		Neutral or mixed	X	Negative		Economic effects	Positive		Neutral or mixed		Negative	X					
Environmental effects	Positive	X	Neutral or mixed		Negative																						
Social effects	Positive		Neutral or mixed	X	Negative																						
Economic effects	Positive		Neutral or mixed		Negative	X																					
<p><i>Rationale:</i> Physical control will have a positive economic side effect in that this species is known to negatively impact forestry operations in New Zealand by competing with forestry trees and making access to plantations more difficult (Gadgil et al., 1984). Costs of clearing <i>C. jubata</i> in forestry plantations was estimated to cost NZD 350 p.ha⁻¹ in 1983 (ca. USD 760 / EUR 662 p.ha⁻¹ at present value) and added 144% to the tending costs of plantations (Gadgil et al., 1984).</p> <p>This measure can have negative environmental effects in that physical control can damage surrounding vegetation, and it can encourage germination of <i>C. jubata</i> seeds (Gosling et al., 2000).</p>																											
<p>Acceptability to stakeholders e.g. impacted economic activities, animal welfare considerations, public perception, etc.</p> <p>Please select one of the categories of acceptability (with an 'X'), and</p>	<table border="1"> <tr> <td>Acceptability to stakeholders</td> <td>Acceptable</td> <td></td> <td>Neutral or mixed</td> <td>X</td> <td>Unacceptable</td> <td></td> </tr> </table>	Acceptability to stakeholders	Acceptable		Neutral or mixed	X	Unacceptable																				
Acceptability to stakeholders	Acceptable		Neutral or mixed	X	Unacceptable																						
<p><i>Rationale:</i> This measure is unlikely to be acceptable for management because of the high cost of physically controlling large invasions of <i>C. jubata</i>. Moreover, physical control has the potential to damage surrounding vegetation, and to provide opportunities for the re-establishment of <i>C. jubata</i>. The difficulty in distinguishing between <i>C. jubata</i> and <i>C. seloana</i> could also make this measure less acceptable. This measure would probably only be acceptable in addition to the use of chemical control.</p>																											

provide a rationale, with supporting evidence and examples if possible.								
<p>Additional cost information ¹</p> <p>When not already included above, or in the species Risk Assessment.</p> <ul style="list-style-type: none"> - implementation cost for Member States - the cost of inaction - the cost-effectiveness - the socio-economic aspects <p>Include quantitative &/or qualitative data, and case studies (incl. from countries outside the EU).</p>	<p>Implementation costs are uncertain, but the best estimates are from New Zealand (see above).</p> <p>Costs of inaction associated with this measure are likely to be similar to those detailed in the <i>Prevention of intentional introductions and spread</i> table above, if this species escapes.</p> <p>The cost-effectiveness of physical control for management of this species is unknown, but probably low because of the high cost of controlling large invasions of this species.</p> <p>There are no known socio-economic aspects.</p>							
<p>Level of confidence on the information provided ²</p> <p>Please select one of the confidence categories along with a statement to support the category chosen. See <i>Notes</i> section at the bottom of this document.</p> <p>NOTE – this is not related to the effectiveness of the measure</p>	<i>Inconclusive</i>	■	<i>Unresolved</i>	■	<i>Established but incomplete</i>	X	<i>Well established</i>	■
<p><i>Rationale:</i> There is adequate evidence to support the use of physical control for management of this species. However, there is no readily available information on the costs of using such an approach, and on the scales at which it is practical and cost-effective.</p>								

<p>Management - Measures to achieve management of the species once it has become widely spread within a Member State, or part of a Member State's territory. (cf. Article 19), i.e. not at an early stage of invasion (see Rapid eradication table above). These measures can be aimed at eradication, population control or containment of a population of the species. This table is repeated for each of the management measures identified.</p>	
<p>Measure description</p> <p>Provide a description of the measure, and identify its objective</p>	<p>As noted above for physical control, the use of chemical control for management of widespread <i>C. jubata</i> invasions is probably unnecessary due to this species probably not having established yet in the EU (unless <i>C. selloana</i> populations have been incorrectly identified).</p> <p>Chemical control</p>

	<p>The use of glyphosate, imazapyr, fluazifop-P-butyl, quizalofop, hexazinone, amitrole, dalapon, and sethoxydim have either been trialled or recommended for use on <i>C. jubata</i> (DiTomaso et al., 2008; Gosling et al., 2000; Schmalzer & Hinkle, 1987). However, only a few of these have been tested using experimental field trials (DiTomaso et al., 2008). Glyphosate is generally the recommended chemical of choice for controlling <i>C. jubata</i> (DiTomaso et al., 2008, 2010; Gosling et al., 2000).</p> <p>It is important to note that EU/national/local legislation on the use of plant protection products and biocides needs to be respected and authorities should check to ensure chemicals are licensed for use in their respective countries/regions</p>						
<p>Scale of application At what scale is the measure applied? What is the largest scale at which it has been successfully used? Please provide examples, with areas (km² or ha) if possible.</p>	<p>Chemical control is the only practical control method for large and difficult to reach invasions of <i>C. jubata</i> (DiTomaso et al., 2010; Gosling et al., 2000; Popay et al., 2003). Examples of the scale at which chemical control has been used, include:</p> <ul style="list-style-type: none"> • New Zealand (blocks around ~2,000 m² in size (Popay et al., 2003) • California. 600 m² in Vandenberg Air Force Base (DiTomaso et al., 2008) • Maui, Hawai'i. <i>C. jubata</i> was “established in numerous areas of rainforest as well as bogs on East and West Maui” and in Haleakala National Park, but has been greatly reduced in number due to an island-wide control campaign (Penniman et al., 2011). In 2012 the Maui and Moloka'i Invasive Species Committees controlled 3,910 acres using 3,063 hours of labour (HISC Established Pests Working Group, 2013). • On Tawhiti Rahi Island and Aorangi Island in New Zealand, both <i>C. jubata</i> and <i>C. selloana</i> were largely eradicated from a 272 ha area (only one detection per year in the last two years of a seven year eradication and monitoring program) (Coulston, 2002). <p>Chemical control therefore seems to be the control method of choice for larger invasions of <i>C. jubata</i>, but the dangers of affecting non-target plants should be kept in mind.</p>						
<p>Effectiveness of the measure Is it effective in relation to its objective? Has the measure previously worked, failed?</p> <p>Please select one of the categories of effectiveness (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<p>Effectiveness of measures</p>	<p>Effective</p>	<p>X</p>	<p>Neutral</p>		<p>Ineffective</p>	
<p><i>Rationale:</i> Chemical control has been shown to be highly effective under certain circumstances. In a Californian trial, spraying of glyphosate at an 8% concentration in early summer was found to be the most cost-effective control method and had a control rate of 99% (However, ropewick application of glyphosate at a 16% concentration in early summer was better for not affecting non-target native species; DiTomaso et al., 2008). In general, glyphosate had a much higher successful control rate than imazapyr, fluazifop and sethoxydim (DiTomaso et al., 2008). The more commonly used spray-to-wet applications of glyphosate (at high concentrations) were as effective as ropewick applications of glyphosate, but even at low concentrations spray-to-wet applications of glyphosate were still more effective than the other herbicides tested (DiTomaso et al., 2008). Larger plants appear to be more likely to survive herbicide control than smaller plants (DiTomaso et al., 2008; Popay et al., 2003).</p> <p>Hexazinone, a root-absorbed herbicide, has been recommended in forestry plantations in California and New Zealand (DiTomaso et al., 2010; Gosling et al., 2000).</p>							

	<p>Aerial spraying of haloxyfop from a beer keg from a helicopter has been shown to quite effective at reducing <i>C. jubata</i> growth, but not entirely effective at killing plants. A year after treatment there were numerous new shoots on treated plants and it appears that follow-up treatments are necessary (Popay et al., 2003).</p> <p>This measure can only be effective if <i>C. jubata</i> can be accurately distinguished from individuals of <i>C. selloana</i>. However, chemical control of both species would ensure the effectiveness of this measure.</p>																					
<p>Effort required e.g. period of time over which measure need to be applied to have results</p>	<p>If applied at the correct concentrations, and when the plants are not stressed, chemical control can be very effective, and follow-up treatments can be limited in number and duration, but there are no experimental trials to provide exact recommendations. However, providing high efficacy of herbicide treatments, follow-up treatments will likely only need to focus on the few surviving adult plants, and only new recruits within the first year as <i>C. jubata</i> seeds do not persist for longer than four months under winter (wet) field conditions (Drewitz & DiTomaso, 2004). In reality, herbicide control is not 100% effective and an eradication program in Hawai'i was only deemed successful after seven years (Penniman et al., 2011).</p> <p>As an example of the effort applied to manage widespread <i>C. jubata</i> and <i>C. selloana</i> invasions, in 2012 the Maui and Moloka'i Invasive Species Committees controlled 3,910 acres using 3,063 hours of labour (HISC Established Pests Working Group, 2013).</p>																					
<p>Resources required¹ e.g. cost, staff, equipment etc.</p>	<p>Herbicide application requires trained staff, equipment (e.g., backpack sprayers, ropewicks, spray wands, or in the case of aerial application, a helicopter fitted with a suitable spray device, e.g., beer keg), herbicides and surfactants.</p> <p>The following studies provided cost estimates (all inclusive) of controlling <i>C. jubata</i>:</p> <ul style="list-style-type: none"> • In California, the most cost effective control method for <i>C. jubata</i> was found to be spray-to-wet application of glyphosate, which cost USD 0.28 per 1 m² plant (USD 2,800 / ca. EUR 2,441 per hectare; DiTomaso et al., 2008). • Ropewick application of glyphosate, a method less likely to affect non-target plants, was estimated to cost USD 0.80 per 1 m² plant (USD 8,000 / ca. EUR 6,975 per hectare; DiTomaso et al., 2008). • In New Zealand, the cheapest herbicide (haloxyfop) control method using helicopters for inaccessible <i>C. jubata</i> infestations was found to cost NZD 3,885 (ca. USD 2500 / EUR 2,219) per hectare. 																					
<p>Side effects (incl. potential) – both positive and negative i.e. positive or negative side effects of the measure on public health, environment including non-targeted species, etc.</p> <p>For each of the side effect types please select one of the impact categories (with an 'X'), and provide a</p>	<table border="1" data-bbox="613 1086 1854 1185"> <tr> <td>Environmental effects</td> <td>Positive</td> <td></td> <td>Neutral or mixed</td> <td></td> <td>Negative</td> <td>X</td> </tr> <tr> <td>Social effects</td> <td>Positive</td> <td></td> <td>Neutral or mixed</td> <td>X</td> <td>Negative</td> <td></td> </tr> <tr> <td>Economic effects</td> <td>Positive</td> <td>X</td> <td>Neutral or mixed</td> <td></td> <td>Negative</td> <td></td> </tr> </table> <p><i>Rationale:</i> For the same reasons as <i>Management: physical control</i>, chemical control will have a positive economic side effect in relation to lower plantation forestry management costs.</p> <p>This measure can have negative environmental effects in that chemical control can also affect native species (DiTomaso et al., 2008; Gosling et al., 2000). Glyphosate is a commonly recommended broad-spectrum herbicide recommended for <i>C. jubata</i> control, but which can also affect neighbouring native plants when applied via sprayers (DiTomaso et al., 2008; Gosling et al.,</p>	Environmental effects	Positive		Neutral or mixed		Negative	X	Social effects	Positive		Neutral or mixed	X	Negative		Economic effects	Positive	X	Neutral or mixed		Negative	
Environmental effects	Positive		Neutral or mixed		Negative	X																
Social effects	Positive		Neutral or mixed	X	Negative																	
Economic effects	Positive	X	Neutral or mixed		Negative																	

<p>rationale, with supporting evidence and examples if possible.</p>	<p>2000). For this reason, ropewick application of glyphosate has been recommended (DiTomaso et al., 2008) and is probably a very practical method for controlling a few plants in a new introduction. Hexazinone, which has been used to treat <i>C. jubata</i> in plantation forests, can affect native plants in light or sandy soils (Gosling et al., 2000). Haloxyfop is a grass-specific herbicide that has been used in aerial spraying of <i>C. jubata</i> in New Zealand and was found to damage the native <i>Austroderia splendens</i> (a close relative of <i>Cortaderia</i> species), but not other native plants, but was also not completely effective at killing <i>C. jubata</i> (Popay et al., 2003).</p> <p>Negative social side effects can include exposure to toxic substances if adequate precautions are not taken.</p>						
<p>Acceptability to stakeholders e.g. impacted economic activities, animal welfare considerations, public perception, etc.</p> <p>Please select one of the categories of acceptability (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<p>Acceptability to stakeholders</p>	<p><i>Acceptable</i></p>		<p><i>Neutral or mixed</i></p>	<p>X</p>	<p><i>Unacceptable</i></p>	
<p><i>Rationale:</i> Invasive species managers are likely to favour chemical control because of its ease of use compared to physical control. However, public perceptions of chemical control are often negative (e.g., Shindler et al., 2011). Moreover, chemical control is not always possible or permitted in conservation areas or in riparian areas and wetlands.</p>							
<p>Additional cost information¹ When not already included above, or in the species Risk Assessment.</p> <ul style="list-style-type: none"> - implementation cost for Member States - the cost of inaction - the cost-effectiveness - the socio-economic aspects <p>Include quantitative &/or qualitative data, and case studies (incl. from countries outside the EU).</p>	<p>The best estimates of implementation costs can be found in the <i>Resources required</i> section above.</p> <p>Costs of inaction associated with this measure are likely to be similar to those detailed in the <i>Prevention of intentional introductions and spread</i> table above, if this species escapes.</p> <p>The cost-effectiveness of chemical control for management of this species is unknown, but probably higher than that of physical control for management.</p> <p>There are no known socio-economic aspects.</p>						
<p>Level of confidence on the information provided²</p> <p>Please select one of the confidence categories along with a statement to support the category chosen. See</p>	<p><i>Inconclusive</i></p>	<p><i>Unresolved</i></p>		<p><i>Established but incomplete</i></p>	<p>X</p>	<p><i>Well established</i></p>	
<p><i>Rationale:</i> There is adequate evidence to support the use of chemical control to manage this species. However, there are insufficient studies providing guidelines on the duration and number of follow-up treatments required. Moreover, there are no cost-benefit analyses on the use of chemical control of <i>C. jubata</i>.</p>							

Notes section at the bottom of this document. NOTE – this is not related to the effectiveness of the measure	
--	--

Management - Measures to achieve management of the species once it has become widely spread within a Member State, or part of a Member State’s territory. (cf. Article 19), i.e. not at an early stage of invasion (see Rapid eradication table above). These measures can be aimed at eradication, population control or containment of a population of the species. This table is repeated for each of the management measures identified.						
Measure description Provide a description of the measure, and identify its objective	<p>Grazing</p> <p>Using domestic livestock to graze <i>C. jubata</i> seedlings has been recommended as a control measure in plantation forests in New Zealand (Gadgil et al., 1984; Gosling et al., 2000) and as a temporary control measure in low-risk areas in Australia (NSW Government, 2018).</p> <p>Fencing, a supply of water for the livestock, and supplementary fodder with high protein are also required for this measure (Gadgil et al., 1984). It has also been suggested that areas invaded by <i>C. jubata</i> seedlings be grazed 3 to 4 times a year, and early in the rotation of plantation forestry (Gosling et al., 2000).</p>					
Scale of application At what scale is the measure applied? What is the largest scale at which it has been successfully used? Please provide examples, with areas (km ² or ha) if possible.	There are no specific recommendations for the scale of application of grazing. Its use in plantation forestry as a control measure suggests that it cannot be used at very broad scales. Moreover, the need for sufficiently high grazing pressure (Gosling et al., 2000; Lambrinos, 2006) also limits its application over large areas.					
Effectiveness of the measure Is it effective in relation to its objective? Has the measure previously worked, failed? Please select one of the categories of effectiveness (with an ‘X’), and provide a rationale, with supporting evidence and examples if possible.	<table border="1" style="width: 100%; text-align: center;"> <tr> <td style="background-color: #fce4d6;"><i>Effectiveness of measures</i></td> <td style="background-color: #fce4d6;"><i>Effective</i></td> <td style="background-color: #fce4d6;"><i>Neutral</i></td> <td style="background-color: #fce4d6;"><i>Ineffective</i></td> <td style="background-color: #fce4d6;">X</td> </tr> </table> <p><i>Rationale:</i> Grazing has only been suggested as an effective control method for seedlings of <i>C. jubata</i>, and only in New Zealand (Gadgil et al., 1984; Gosling et al., 2000) and Australia (NSW Government, 2018). In California, grazing is not recommended as an effective control method (e.g. DiTomaso et al., 2010), and in other regions grazing is not mentioned at all.</p> <p>The effectiveness of grazing is limited because of the difficulties of managing grazing (Gadgil et al., 1984; Gosling et al., 2000), because only seedlings are grazed, and because <i>C. jubata</i> still successfully invades even under high grazing pressure (Lambrinos,</p>	<i>Effectiveness of measures</i>	<i>Effective</i>	<i>Neutral</i>	<i>Ineffective</i>	X
<i>Effectiveness of measures</i>	<i>Effective</i>	<i>Neutral</i>	<i>Ineffective</i>	X		

	2006). Even with high herbivore pressure, <i>C. jubata</i> was still able to successfully invade in Californian chaparral, increasing in cover by 20% over 9 years (Lambrinos, 2006).						
Effort required e.g. period of time over which measure need to be applied to have results	There is little information available on the effort required. Gosling et al. (2000) recommend grazing 3 or 4 times a year, but provide no further details.						
Resources required ¹ e.g. cost, staff, equipment etc.	Suitable domestic livestock (cattle, goats or sheep), fencing, water supply, additional high-protein fodder, labour to manage the livestock.						
Side effects (incl. potential) – both positive and negative i.e. positive or negative side effects of the measure on public health, environment including non-targeted species, etc. For each of the side effect types please select one of the impact categories (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.	Environmental effects	Positive		Neutral or mixed	X	Negative	
	Social effects	Positive	X	Neutral or mixed		Negative	
	Economic effects	Positive		Neutral or mixed	X	Negative	
	<i>Rationale:</i> Grazing in environmentally sensitive areas could damage or kill native plant species, and influence nutrient flows. However, livestock would also eat <i>C. selloana</i> seedlings, providing control for both <i>Cortaderia</i> species.						
	Grazing could have positive social and economic side effects through the employment and income generated by allowing grazing in plantations or protected areas, for example. However, there is the potential for issues with animal health, particularly for cattle, if they are allowed to graze extensively on <i>Cortaderia</i> . This can lead to the development of “woody tongue” (Maas, 2009).						
Acceptability to stakeholders e.g. impacted economic activities, animal welfare considerations, public perception, etc. Please select one of the categories of acceptability (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.	Acceptability to stakeholders	Acceptable		Neutral or mixed		Unacceptable	X
	<i>Rationale:</i> The ineffectiveness of this measure is likely to make it unacceptable to invasive species managers. The possible negative side effects on native species will also make this measure unacceptable to conservation managers.						
Additional cost information ¹ When not already included above, or in the species Risk Assessment. - implementation cost for Member States - the cost of inaction	Implementation costs are unknown, but will be linked to the <i>Resources required</i> , as detailed in the relevant section above. Costs of inaction associated with this measure are likely to be similar to those detailed in the <i>Prevention of intentional introductions and spread</i> table above, if this species escapes. The cost-effectiveness of this measure is unknown.						

<p>- the cost-effectiveness - the socio-economic aspects</p> <p>Include quantitative &/or qualitative data, and case studies (incl. from countries outside the EU).</p>	<p>There are no known socio-economic aspects.</p>							
<p>Level of confidence on the information provided²</p> <p>Please select one of the confidence categories along with a statement to support the category chosen. See <i>Notes</i> section at the bottom of this document. NOTE – this is not related to the effectiveness of the measure</p>	<p><i>Inconclusive</i></p>	<p>X</p>	<p><i>Unresolved</i></p>		<p><i>Established but incomplete</i></p>		<p><i>Well established</i></p>	
<p>There do not appear to be any studies experimentally or quantitatively investigating the use of grazing for <i>C. jubata</i> (or <i>Cortaderia</i> spp. in general) management. Much of the evidence for or against the use of grazing for <i>C. jubata</i> control is anecdotal (e.g. DiTomaso et al., 2010; Gadgil et al., 1984; Gosling et al., 2000; NSW Government, 2018). The only study to scientifically investigate this measure was Lambrinos (2006), but this involved grazing by native herbivores, and also found that grazing was insufficient to prevent the spread of <i>C. jubata</i>.</p>								

<p>Management - Measures to achieve management of the species once it has become widely spread within a Member State, or part of a Member State's territory. (cf. Article 19), i.e. not at an early stage of invasion (see Rapid eradication table above). These measures can be aimed at eradication, population control or containment of a population of the species. This table is repeated for each of the management measures identified.</p>	
<p>Measure description Provide a description of the measure, and identify its objective</p>	<p>Oversowing</p> <p>In New Zealand, oversowing in timber plantations (or recently felled plantations or sites being prepared for plantations) has been recommended, in conjunction with grazing, as a way of controlling <i>C. jubata</i> invasions in timber plantations (Gadgil et al., 1984; Gosling et al., 2000). Nitrogen-fixing species (e.g. <i>Lotus pedunculatus</i>) were specifically recommended for oversowing in these studies.</p>
<p>Scale of application At what scale is the measure applied? What is the largest scale at which it has been successfully used? Please provide examples, with areas (km² or ha) if possible.</p>	<p>There are no specific recommendations for the scale of application of oversowing. However, its use in conjunction with grazing in plantation forestry as a control measure suggests that it cannot be used at very broad scales.</p>

<p>Effectiveness of the measure Is it effective in relation to its objective? Has the measure previously worked, failed?</p> <p>Please select one of the categories of effectiveness (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<table border="1"> <tr> <td>Effectiveness of measures</td> <td><i>Effective</i></td> <td></td> <td><i>Neutral</i></td> <td></td> <td><i>Ineffective</i></td> <td>X</td> </tr> </table>	Effectiveness of measures	<i>Effective</i>		<i>Neutral</i>		<i>Ineffective</i>	X														
Effectiveness of measures	<i>Effective</i>		<i>Neutral</i>		<i>Ineffective</i>	X																
<p>Effort required e.g. period of time over which measure need to be applied to have results</p>	<p>The effort required is unknown.</p>																					
<p>Resources required¹ e.g. cost, staff, equipment etc.</p>	<p>Resources will be the same as for <i>Management</i> using <i>Grazing</i>, except that the additional high-protein fodder will be provided by the species used for oversowing.</p>																					
<p>Side effects (incl. potential) – both positive and negative i.e. positive or negative side effects of the measure on public health, environment including non-targeted species, etc.</p> <p>For each of the side effect types please select one of the impact categories (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<table border="1"> <tr> <td>Environmental effects</td> <td><i>Positive</i></td> <td></td> <td><i>Neutral or mixed</i></td> <td>X</td> <td><i>Negative</i></td> <td></td> </tr> <tr> <td>Social effects</td> <td><i>Positive</i></td> <td></td> <td><i>Neutral or mixed</i></td> <td>X</td> <td><i>Negative</i></td> <td></td> </tr> <tr> <td>Economic effects</td> <td><i>Positive</i></td> <td></td> <td><i>Neutral or mixed</i></td> <td>X</td> <td><i>Negative</i></td> <td></td> </tr> </table> <p><i>Rationale:</i> Environmental side effects could be both negative and positive. A negative side effect might include the possibility of sowing other invasive species (recommended species include <i>Lotus pedunculatus</i>, an alien to New Zealand). A positive side effect might be erosion prevention.</p> <p>There are no known social or economic side effects.</p>	Environmental effects	<i>Positive</i>		<i>Neutral or mixed</i>	X	<i>Negative</i>		Social effects	<i>Positive</i>		<i>Neutral or mixed</i>	X	<i>Negative</i>		Economic effects	<i>Positive</i>		<i>Neutral or mixed</i>	X	<i>Negative</i>	
Environmental effects	<i>Positive</i>		<i>Neutral or mixed</i>	X	<i>Negative</i>																	
Social effects	<i>Positive</i>		<i>Neutral or mixed</i>	X	<i>Negative</i>																	
Economic effects	<i>Positive</i>		<i>Neutral or mixed</i>	X	<i>Negative</i>																	
<p>Acceptability to stakeholders e.g. impacted economic activities, animal welfare considerations, public perception, etc.</p> <p>Please select one of the categories of acceptability (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<table border="1"> <tr> <td>Acceptability to stakeholders</td> <td><i>Acceptable</i></td> <td></td> <td><i>Neutral or mixed</i></td> <td></td> <td><i>Unacceptable</i></td> <td>X</td> </tr> </table>	Acceptability to stakeholders	<i>Acceptable</i>		<i>Neutral or mixed</i>		<i>Unacceptable</i>	X														
Acceptability to stakeholders	<i>Acceptable</i>		<i>Neutral or mixed</i>		<i>Unacceptable</i>	X																
<p>Additional cost information¹</p>	<p>Implementation costs are unknown, but will be linked to the <i>Resources required</i>, as detailed in the relevant section above.</p>																					

<p>When not already included above, or in the species Risk Assessment.</p> <ul style="list-style-type: none"> - implementation cost for Member States - the cost of inaction - the cost-effectiveness - the socio-economic aspects <p>Include quantitative &/or qualitative data, and case studies (incl. from countries outside the EU).</p>	<p>Costs of inaction associated with this measure are likely to be similar to those detailed in the <i>Prevention of intentional introductions and spread</i> table above, if this species escapes.</p> <p>The cost-effectiveness of this measure is unknown.</p> <p>There are no known socio-economic aspects.</p>								
<p>Level of confidence on the information provided ²</p> <p>Please select one of the confidence categories along with a statement to support the category chosen. See <i>Notes</i> section at the bottom of this document.</p> <p>NOTE – this is not related to the effectiveness of the measure</p>	<p><i>Inconclusive</i></p>	<p>X</p>	<p><i>Unresolved</i></p>		<p><i>Established but incomplete</i></p>		<p><i>Well established</i></p>		<p>There do not appear to be any studies experimentally or quantitatively investigating the use of oversowing for <i>C. jubata</i> (or <i>Cortaderia</i> spp. in general) management. All the evidence for the use of grazing for <i>C. jubata</i> control is anecdotal and is very specific to plantation forestry in New Zealand (Gadgil et al., 1984; Gosling et al., 2000).</p>

<p>Management - Measures to achieve management of the species once it has become widely spread within a Member State, or part of a Member State's territory. (cf. Article 19), i.e. not at an early stage of invasion (see Rapid eradication table above). These measures can be aimed at eradication, population control or containment of a population of the species. This table is repeated for each of the management measures identified.</p>	
<p>Measure description</p> <p>Provide a description of the measure, and identify its objective</p>	<p>Biological control</p> <p>This measure has received relatively little attention for management of <i>C. jubata</i> with papers as recent as 2010 suggesting that biological control had not been investigated at all (e.g. DiTomaso et al., 2010). However, recently there appears to be more interest in this measure, particularly in New Zealand. Bellgard et al. (2010) conducted a survey of <i>C. jubata</i> and <i>C. selloana</i> in New Zealand for the presence of native invertebrate herbivores and fungal pathogens. They found the native flax notcher (<i>Tmetolophota steropastis</i>) to be the most damaging invertebrate herbivore, and identified a number of damaging fungal pathogens. In a follow-up study, one of these fungal pathogens (<i>Nigrospora oryzae</i>), used in conjunction with synthetic herbicides, was found to cause</p>

	<p>greater dieback of both <i>Cortaderia</i> species than when herbicides were applied without an inoculation of this fungus (Bellgard, Probst & Johnson, 2016).</p> <p>A recent study in Ecuador using a native fungal pathogen (<i>Ustilago quitensis</i>) found high infection rates in the inflorescences of <i>C. jubata</i> when using this fungus as a biological control agent (Torres et al., 2016).</p> <p>It is important to note that the release of macro-organisms as biological control agents is currently not regulated at EU level. Nevertheless national/regional laws are to be respected. Before any release of an alien species as a biological control agent an appropriate risk assessment should be made</p>																					
<p>Scale of application At what scale is the measure applied? What is the largest scale at which it has been successfully used? Please provide examples, with areas (km² or ha) if possible.</p>	<p>Unknown. No biological control agent has been released as yet.</p>																					
<p>Effectiveness of the measure Is it effective in relation to its objective? Has the measure previously worked, failed?</p> <p>Please select one of the categories of effectiveness (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<table border="1" data-bbox="613 724 1854 786"> <tr> <td><i>Effectiveness of measures</i></td> <td><i>Effective</i></td> <td></td> <td><i>Neutral</i></td> <td></td> <td><i>Ineffective</i></td> <td>X</td> </tr> </table> <p><i>Rationale:</i> Unknown. No biological control agent has been released or gone through experimental trials yet.</p>	<i>Effectiveness of measures</i>	<i>Effective</i>		<i>Neutral</i>		<i>Ineffective</i>	X														
<i>Effectiveness of measures</i>	<i>Effective</i>		<i>Neutral</i>		<i>Ineffective</i>	X																
<p>Effort required e.g. period of time over which measure need to be applied to have results</p>	<p>Unknown.</p>																					
<p>Resources required¹ e.g. cost, staff, equipment etc.</p>	<p>Unknown, although resources required are likely to be similar to most other biological control agents, e.g. rearing facilities, trained staff to release/apply biological control agent, etc. The cost of testing and monitoring of a biological control agent is unknown.</p>																					
<p>Side effects (incl. potential) – both positive and negative i.e. positive or negative side effects of the measure on public health,</p>	<table border="1" data-bbox="613 1259 1854 1358"> <tr> <td><i>Environmental effects</i></td> <td><i>Positive</i></td> <td></td> <td><i>Neutral or mixed</i></td> <td></td> <td><i>Negative</i></td> <td>X</td> </tr> <tr> <td><i>Social effects</i></td> <td><i>Positive</i></td> <td></td> <td><i>Neutral or mixed</i></td> <td></td> <td><i>Negative</i></td> <td></td> </tr> <tr> <td><i>Economic effects</i></td> <td><i>Positive</i></td> <td></td> <td><i>Neutral or mixed</i></td> <td></td> <td><i>Negative</i></td> <td></td> </tr> </table>	<i>Environmental effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>		<i>Negative</i>	X	<i>Social effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>		<i>Negative</i>		<i>Economic effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>		<i>Negative</i>	
<i>Environmental effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>		<i>Negative</i>	X																
<i>Social effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>		<i>Negative</i>																	
<i>Economic effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>		<i>Negative</i>																	

<p>environment including non-targeted species, etc.</p> <p>For each of the side effect types please select one of the impact categories (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<p><i>Rationale:</i> Although highly unlikely if proper testing protocols are followed, there is the potential for a biological control agent to affect a non-target species. There are no native <i>Cortaderia</i> species in the EU. No other side effects seem likely.</p>						
<p>Acceptability to stakeholders e.g. impacted economic activities, animal welfare considerations, public perception, etc.</p> <p>Please select one of the categories of acceptability (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<p>Acceptability to stakeholders</p>	<p>Acceptable</p>		<p>Neutral or mixed</p>	<p>X</p>	<p>Unacceptable</p>	
<p>Additional cost information¹ When not already included above, or in the species Risk Assessment.</p> <ul style="list-style-type: none"> - implementation cost for Member States - the cost of inaction - the cost-effectiveness - the socio-economic aspects <p>Include quantitative &/or qualitative data, and case studies (incl. from countries outside the EU).</p>	<p>Implementation costs are unknown.</p> <p>Costs of inaction associated with this measure are likely to be similar to those detailed in the <i>Prevention of intentional introductions and spread</i> table above, if this species escapes.</p> <p>The cost-effectiveness of this measure is unknown, but biological control is typically ranked as one of the most cost-effective control measures for widespread alien plant species (which <i>C. jubata</i> is thought not to be, in the EU) (Barratt, Moran, Bigler & Van Lenteren, 2018).</p> <p>There are no socio-economic aspects as <i>C. jubata</i> is not thought to be widespread in the EU, and there are currently no biological control agents available for this species.</p>						
<p>Level of confidence on the information provided²</p> <p>Please select one of the confidence categories along with a statement to support the category chosen. See <i>Notes</i> section at the bottom of this document.</p>	<p>Inconclusive</p>	<p>X</p>	<p>Unresolved</p>		<p>Established but incomplete</p>	<p>Well established</p>	
	<p>Very little research has been conducted on using biological control for <i>C. jubata</i> invasions.</p>						

NOTE – this is not related to the effectiveness of the measure

Bibliography³

See guidance section

- Andreu, J., Vila, M., & Hulme, P. E. (2009). An assessment of stakeholder perceptions and management of noxious alien plants in Spain. *Environmental Management*, 43(6), 1244. Retrieved from <https://doi.org/10.1007/s00267-009-9280-1>.
- Barratt, B. I. P., Moran, V. C., Bigler, F., & Van Lenteren, J. C. (2018). The status of biological control and recommendations for improving uptake for the future. *BioControl*, 63(1), 155-167.
- Bellgard, S. E., Winks, C. J., Than, D. J., & Aliaga, C. C. (2010). Natural enemies of the South American pampas grasses *Cortaderia* spp. in New Zealand. In S. M. Zydenbos (Ed.), *17th Australasian weeds conference proceedings: New frontiers in New Zealand, together we can beat the weeds: 26-30 September, 2010* (pp. 239-242). Christchurch, New Zealand: New Zealand Plant Protection Society.
- Bellgard, S. E., Probst, C. M., & Johnson, V. W. (2016). Synergism between synthetic herbicides and *Nigrospora oryzae* (Berk. & Broome) Petch for the inundative biocontrol of pampas in New Zealand. In *20th Australasian Weeds Conference, Perth, Western Australia, 11-15 September 2016* (pp. 274-278). Weeds Society of Western Australia.
- CABI. (2018). *Invasive Species Compendium Datasheet: Cortaderia jubata (purple pampas grass)*. Retrieved from <http://www.cabi.org/isc/datasheet/113484>.
- Cal-IPC. (2018) *Invasive Plants of California's Wildland Plant Report: Cortaderia jubata*. Retrieved from <http://www.cal-ipc.org/ip/management/ipcw/pages/detailreport.cfm@usernumber=33&surveynumber=182.php>.
- Carrière, E. A. (1878). *Gynerium jubatum*. *Revue Horticole*, 50, 449-450.
- Clayton, W. D., Vorontsova, M. S., Harman, K. T., & Williamson, H. (2006 onwards). *GrassBase - The Online World Grass Flora*. Retrieved from <https://www.kew.org/data/grasses-db/www/imp02543.htm>.
- Costas Lippmann, M. (1977). More on the weedy "pampas grass" in California. *Fremontia*, 4(4), 25-27.
- Coulston, G. J. (2002). Control of invasive plants on Poor Knights Islands, New Zealand. In C. R. Veitch, & M. N. Clout (Eds.), *Turning the tide: the eradication of invasive species* (pp.79-84). Gland, Switzerland and Cambridge, UK: IUCN SSC Invasive Species Specialist Group.
- Daehler, C. (2006). *Australian/New Zealand Weed Risk Assessment adapted for Hawai'i.: Cortaderia jubata*. Retrieved from http://www.hear.org/pier/wra/pacific/cortaderia_jubata_htmlwra.htm.
- DAISIE European Invasive Alien Species Gateway. (2008). *Cortaderia selloana*. Retrieved from <http://www.europe-aliens.org/speciesFactsheet.do?speciesId=3364>.
- DiTomaso, J. M., Drewitz, J. J., & Kyser, G. B. (2008). Jubatagrass (*Cortaderia jubata*) control using chemical and mechanical methods. *Invasive Plant Science and Management*, 1(1), 82-90. Retrieved from <https://doi.org/10.1614/IPSM-07-028>.
- DiTomaso, J. M., Healy, E., Bell, C. E., Drewitz, J., & Stanton, A. (2010). *Pampasgrass and jubatagrass threaten California coastal habitats*. Retrieved from <https://wric.ucdavis.edu/PDFs/pampasgrass%20and%20jubatagrass%20WRIC%20leaflet%2099-1.pdf>.

- Drewitz, J. J., & DiTomaso, J. M. (2004). Seed biology of jubatagrass (*Cortaderia jubata*). *Weed science*, 52(4), 525-530. Retrieved from <https://doi.org/10.1614/WS-03-081R>.
- Edgar, E., & Connor, H. E. (2000). *Flora of New Zealand Vol. V. Gramineae*. Lincoln, New Zealand: Manaaki Whenua Press. Retrieved from http://floraseries.landcareresearch.co.nz/pages/Taxon.aspx?id=_779f0059-bb6f-46ac-a5df-9ee70c7c87c3&fileName=Flora%205.xml.
- Epanchin-Niell, R. S., Haight, R. G., Berec, L., Kean, J. M., & Liebhold, A. M. (2012). Optimal surveillance and eradication of invasive species in heterogeneous landscapes. *Ecology letters*, 15(8), 803-812. Retrieved from <https://doi.org/10.1111/j.1461-0248.2012.01800.x>.
- EPPO. (2018). Pest risk analysis for *Cortaderia jubata*. EPPO, Paris.
- Gadgil, R. L., Knowles, A. L., & Zabkiewicz, J. A. (1984). Pampas-a new forest weed problem. In M. J. Hartley (Ed.), *Proceedings 37th New Zealand Weed and Pest Control Conference* (pp.187-190). New Zealand Weed and Pest Control Society.
- Gallo, T., & Waite, D. (2011). Creating a successful citizen science model to detect and report invasive species. *BioScience*, 61(6), 459-465. Retrieved from <https://doi.org/10.1525/bio.2011.61.6.8>.
- Genovesi, P., Scalera, R., Brunel, S., Roy, D., & Solarz, W. (2010). *Towards an early warning and information system for invasive alien species (IAS) threatening biodiversity in Europe* (EEA Technical Report No 5/2010). Copenhagen: European Environment Agency. Retrieved from <https://www.eea.europa.eu/publications/information-system-invasive-alien-species>.
- Gosling, D. S., Shaw, W. B., & Beadel, S. M. (2000). *Review of control methods for pampas grasses in New Zealand*. Science for Conservation, No. 165. Retrieved from <http://www.doc.govt.nz/documents/science-and-technical/Sfc165.pdf>.
- Hauser, C. E., & McCarthy, M. A. (2009). Streamlining 'search and destroy': cost-effective surveillance for invasive species management. *Ecology Letters*, 12(7), 683-692. Retrieved from <https://doi.org/10.1111/j.1461-0248.2009.01323.x>.
- Hawai'i Department of Transportation (HDOT). (2011). *2012-2022 Statewide Noxious Invasive Pest Program (SNIPP) Strategic Plan*. Retrieved from https://hidot.hawaii.gov/highways/files/2013/02/Landscape-SNIPP_Strategic_Plan.pdf.
- HISC Established Pests Working Group. (2013). Maui Invasive Species Committee and Moloka'i Invasive Species Committee 2012 Financial Year Report. Retrieved from <https://dlnr.hawaii.gov/hisc/files/2013/03/FY12-Report-MISC-MoMISC-EPWG.pdf>.
- Hooker, J. D. (1898). *Cortaderia jubata* Stapf. *Curtis's Botanical Magazine*, 54, Tab. 7607.
- Houliston, G. J., & Goeke, D. F. (2017). *Cortaderia* spp. in New Zealand: patterns of genetic variation in two widespread invasive species. *New Zealand Journal of Ecology*, 41(1), 107-112. Retrieved from <https://doi.org/10.20417/nzjecol.41.13>
- IPPC. (2017). ISMP 41 International movement of used vehicles, machinery and equipment. 12 pp. Rome: FAO. Retrieved from (https://www.ippc.int/static/media/files/publication/en/2017/05/ISPM_41_2017_En_2017-05-15.pdf)
- Keller, R. P., Lodge, D. M., & Finnoff, D. C. (2007). Risk assessment for invasive species produces net bioeconomic benefits. *Proceedings of the National Academy of Sciences*, 104(1), 203-207. Retrieved from <https://doi.org/10.1073/pnas.0605787104>.
- Kettunen, M., Heikkilä, J., Underwood, E., & Vyliaudaite, I. (2014). *Instruments for financing action on invasive alien species (IAS): review and assessment of selected examples and their applicability in Finland*. London & Brussels: Institute for European Environmental Policy. Retrieved from https://ieep.eu/uploads/articles/attachments/d1b7abe7-75be-4aeb-8d82-b75194bf1272/Kettunen_at_al_2014_review_of_instruments_financing_IAS_action_FINAL_June_2014.pdf?v=63664509852.


- Lambrinos, J.G. (2000). The impact of the invasive alien grass *Cortaderia jubata* (Lemoine) Stapf on an endangered mediterranean-type shrubland in California. *Diversity and Distributions*, 6(5), 217-231. Retrieved from <https://doi.org/10.1046/j.1472-4642.2000.00086.x>
- Lambrinos, J. G. (2006). Spatially variable propagule pressure and herbivory influence invasion of chaparral shrubland by an exotic grass. *Oecologia*, 147(2), 327-334. Retrieved from <https://doi.org/10.1007/s00442-005-0259-1>
- Leung, B., Lodge, D. M., Finnoff, D., Shogren, J. F., Lewis, M. A., & Lamberti, G. (2002). An ounce of prevention or a pound of cure: bioeconomic risk analysis of invasive species. *Proceedings of the Royal Society of London B: Biological Sciences*, 269(1508), 2407-2413. Retrieved from <https://doi.org/10.1098/rspb.2002.2179>.
- Maas, J. (2009). Woody tongue: recognition, treatment, and prevention. *UCD Vet Views: California Cattlemen's Magazine*, June 2009. Retrieved from https://www2.vetmed.ucdavis.edu/vetext/local_resources/pdfs/pdfs_beef/cca0906-woody-tongue.pdf
- Maistrello, L., Dioli, P., Bariselli, M., Mazzoli, G. L., & Giacalone-Forini, I. (2016). Citizen science and early detection of invasive species: phenology of first occurrences of *Halyomorpha halys* in Southern Europe. *Biological Invasions*, 18(11), 3109-3116. Retrieved from <https://doi.org/10.1007/s10530-016-1217-z>.
- Messing, R., & Brodeur, J. (2018). Current challenges to the implementation of classical biological control. *BioControl*, 63(1), 1-9. Retrieved from <https://doi.org/10.1007/s10526-017-9862-4>
- New Zealand Plant Conservation Network. (2018). *New Zealand's Flora: Cortaderia jubata*. Retrieved from http://www.nzpcn.org.nz/flora_details.aspx?ID=3752
- NSW Government. (2018). *NSW WeedWise: Pampas grass (Cortaderia species)*. Retrieved from <http://weeds.dpi.nsw.gov.au/Weeds/Details/100>.
- Okada, M., Ahmad, R., & Jasieniuk, M. (2007). Microsatellite variation points to local landscape plantings as sources of invasive pampas grass (*Cortaderia selloana*) in California. *Molecular ecology*, 16(23), 4956-4971. Retrieved from <https://doi.org/10.1111/j.1365-294X.2007.03568.x>.
- Parish, J. A. (2017). *Implementing an early detection program on Catalina Island: prioritizing landscaped grasses*. Retrieved from https://www.cal-ipc.org/wp-content/uploads/2018/02/2017_Symposium-early-detection-program-Catalina-Island-Parish.pdf.
- Parsons, W. T., & Cuthbertson, E. G. (2004). *Noxious weeds of Australia*. Second Edition. Collingwood, Australia: CSIRO publishing.
- Penniman, T. M., Buchanan, L., & Loope, L. L. (2011). Recent plant eradications on the islands of Maui County, Hawai'i. In C. R. Veitch, M. N. Clout, & D. R. Towns (Eds.), *Island invasives: eradication and management* (pp.325-331). Gland: IUCN.
- Peterson, D. L., & Russo, M. J. (1988). *Element Stewardship Abstract for Cortaderia jubata, pampas grass*. Arlington, Virginia, USA: The Nature Conservancy. Retrieved from <http://www.invasive.org/gist/esadocs/documnts/cortjub.pdf>.
- Popay, I., Timmins, S. M., & McCluggage, T. (2003). *Aerial spraying of pampas grass in difficult conservation sites*. Science for Conservation, 218. Wellington, New Zealand: Department of Conservation. Retrieved from <http://www.doc.govt.nz/documents/science-and-technical/sfc218.pdf>.
- Queensland Government. (2017). *Cortaderia jubata. Weeds of Australia; Biosecurity Queensland Edition*. Retrieve from https://keyserver.lucidcentral.org/weeds/data/media/Html/cortaderia_jubata.htm
- Robinson, E. R. (1984). Naturalized species of *Cortaderia* (Poaceae) in southern Africa. *South African Journal of Botany*, 3(5), 343-346. Retrieved from [https://doi.org/10.1016/S0022-4618\(16\)30023-7](https://doi.org/10.1016/S0022-4618(16)30023-7)
- Regan, T. J., McCarthy, M. A., Baxter, P. W., Dane Panetta, F., & Possingham, H. P. (2006). Optimal eradication: when to stop looking for an invasive plant. *Ecology letters*, 9(7), 759-766. Retrieved from <https://doi.org/10.1111/j.1461-0248.2006.00920.x>.
- Royal Horticultural Society. (2009). *Cortaderia: final trials report 2007-2009*. Wisley, UK: Royal Horticultural Society. Retrieved from <http://apps.rhs.org.uk/planttrials/TrialReports/Cortaderia%202009.pdf>

- Saura-Mas, S., & Lloret, F. (2005). Wind effects on dispersal patterns of the invasive alien *Cortaderia selloana* in Mediterranean wetlands. *Acta Oecologica*, 27(2), 129-133. Retrieved from <https://doi.org/10.1016/j.actao.2004.12.001>.
- Schmalzer, P. A., & Hinkle, C. R. (1987). *Species biology and potential for controlling four exotic plants (Ammophila arenaria, Carpobrotus edulis, Cortaderia jubata and Gasoul crystallinum) on Vandenberg Air Force Base, California* (NASA Technical Memorandum 100980). Kennedy Space Center, Florida: NASA. Retrieved from <https://ntrs.nasa.gov/archive/nasa/casi.ntrs.nasa.gov/19880008764.pdf>.
- Shindler, B., Gordon, R., Brunson, M. W., & Olsen, C. (2011). Public perceptions of sagebrush ecosystem management in the Great Basin. *Rangeland Ecology & Management*, 64(4), 335-343. Retrieved from <https://doi.org/10.2111/REM-D-10-00012.1>.
- Silvertown, J. (2009). A new dawn for citizen science. *Trends in Ecology & Evolution*, 24(9), 467-471. Retrieved from <https://doi.org/10.1016/j.tree.2009.03.017>.
- Starr, F., Starr, K. & Loope, L. (2003). *Cortaderia spp.* Hawaiian Ecosystems at Risk project (HEAR). Retrieved from http://www.hear.org/Pier/pdf/pohreports/cortaderia_spp.pdf
- Testoni, D., & Linder, H. P. (2017). Synoptic taxonomy of *Cortaderia* Stapf (Danthonioideae, Poaceae). *PhytoKeys*, 76, 39-69. Retrieved from <https://doi.org/10.3897/phytokeys.76.10808>.
- Testoni, D., & Villamil, C. B. (2014). Estudios en el género *Cortaderia* (Poaceae): I. Sistemática y nomenclatura de la sect. *Cortaderia*. *Darwiniana*, nueva serie, 2(2), 260-276.
- Tobin, P. C., Kean, J. M., Suckling, D. M., McCullough, D. G., Herms, D. A., & Stringer, L. D. (2014). Determinants of successful arthropod eradication programs. *Biological Invasions*, 16(2), 401-414. Retrieved from <https://doi.org/10.1007/s10530-013-0529-5>.
- Toomey, A. H., & Domroese, M. C. (2013). Can citizen science lead to positive conservation attitudes and behaviors? *Human Ecology Review*, 20(1), 50-62. Retrieved from <https://www.jstor.org/stable/24707571>.
- Torres, G., Jackeline, L., Salazar, A., & Andrea, C. (2016). *Evaluación del efecto biológico del hongo Ustilago quitensis sobre la meleza Cortaderia jubata* (Bachelor's thesis, Universidad de las Américas, Quito, Ecuador). Retrieved from <http://200.24.220.94/bitstream/33000/5465/1/UDLA-EC-TIB-2016-10.pdf>
- Underwood, E., Ustin, S., & DiPietro, D. (2003). Mapping nonnative plants using hyperspectral imagery. *Remote Sensing of Environment*, 86(2), 150-161. Retrieved from [https://doi.org/10.1016/S0034-4257\(03\)00096-8](https://doi.org/10.1016/S0034-4257(03)00096-8).
- University of Queensland. (2018). *Weeds of Australia: Cortaderia jubata*. Retrieved from https://keyserver.lucidcentral.org/weeds/data/media/Html/cortaderia_jubata.htm.
- U.S. Department of Transportation Federal Highway Administration. (2013). Functional system lane-length – 2013. Retrieved from <https://www.fhwa.dot.gov/policyinformation/statistics/2013/hm60.cfm>.

Notes

1. Costs information. The assessment of the potential costs shall describe those costs quantitatively and/or qualitatively depending on what information is available. This can include case studies from across the Union or third countries.

2. Level of confidence¹: based on the quantity, quality and level of agreement in the evidence.



- **Well established:** comprehensive meta-analysis² or other synthesis or multiple independent studies that agree.
- **Established but incomplete:** general agreement although only a limited number of studies exist but no comprehensive synthesis and, or the studies that exist imprecisely address the question.
- **Unresolved:** multiple independent studies exist but conclusions do not agree.
- **Inconclusive:** limited evidence, recognising major knowledge gaps

3. Citations and bibliography. The APA formatting style for citing references in the text and in the bibliography is used. e.g. Peer review papers will be written as follows:

¹ Assessment of confidence methodology is taken from IPBES. 2016. Guide on the production and integration of assessments from and across all scales (IPBES-4-INF-9), which is adapted from Moss and Schneider (2000).

² A statistical method for combining results from different studies which aims to identify patterns among study results, sources of disagreement among those results, or other relationships that may come to light in the context of multiple studies.

In text citation: (Author & Author, Year)

In bibliography: Author, A. A., & Author, B. B. (Publication Year). Article title. *Periodical Title*, Volume(Issue), pp.-pp.

(see <http://www.waikato.ac.nz/library/study/referencing/styles/apa>)