

Information on measures and related costs in relation to species considered for inclusion on the Union list: *Triadica sebifera*

This note has been drafted by IUCN within the framework of the contract No 07.0202/2017/763436/SER/ENV.D2 “Technical and Scientific support in relation to the Implementation of Regulation 1143/2014 on Invasive Alien Species”. The information and views set out in this note do not necessarily reflect the official opinion of the Commission, or IUCN. The Commission does not guarantee the accuracy of the data included in this note. Neither the Commission nor IUCN or any person acting on the Commission’s behalf, including any authors or contributors of the notes themselves, may be held responsible for the use which may be made of the information contained therein. Reproduction is authorised provided the source is acknowledged. This document shall be cited as:

Flory, S. L. 2018. Information on measures and related costs in relation to species considered for inclusion on the Union list: *Triadica sebifera*. Technical note prepared by IUCN for the European Commission.

Date of completion: 25/10/2018

Comments which could support improvement of this document are welcome. Please send your comments by e-mail to ENV-IAS@ec.europa.eu.

Species (scientific name)	<i>Triadica sebifera</i> (L.) Small (ITIS Standard Report 2018) [previously <i>Sapium sebiferum</i>]
Species (common name)	Chinese tallow tree, tallowtree, chicken tree, Florida aspen, popcorn tree, vegetable tallow, white wax berry
Author(s)	S. Luke Flory, University of Florida, USA
Date Completed	25/10/18
Reviewer	Geoff Wang, Clemson University, USA

Summary

Highlight of measures that provide the most cost-effective options to prevent the introduction, achieve early detection, rapidly eradicate and manage the species, including significant gaps in information or knowledge to identify cost-effective measures.

Triadica sebifera is a medium-sized deciduous tree that can grow up to 16 meters tall (CABI, 2018). The species is native to Asia, including China, Japan, and Vietnam. It has been introduced widely across the globe for its ornamental value and commercial applications in industry. The tree has most widely been introduced to the United States, but it also has been planted in Europe, South America, Africa, Asia and Australia (CABI, 2018; EDDMAPS, 2018; GBIF, 2018). Following introduction to the United States and Australia, it escaped cultivation and spread such that it is now widely naturalized in both countries (NSW Government Factsheet, 2017; USDA, 2008). *Triadica sebifera* currently does not occur in the natural environment in the European Union.

To develop effective measures to prevent the introduction and spread of the species in the EU and determine costs of such measures, the pathways and mechanisms for introduction and spread and how to detect the species must be considered. In addition, the biological characteristics of *T. sebifera* present some unique challenges for prevention of spread, eradication, and management. For example, it has been reported that more than 95% of *T. sebifera* seeds can initially be viable (Renne, Barrow,

Randall & Bridges, 2002) and adult trees can produce over 100,000 seeds per year (USDA, 2008; Jubinsky & Anderson, 1996; Bruce, Cameron, Harcombe & Jubinsky, 1997). Additionally, dispersal of *T. sebifera* plants can occur intentionally through ornamental or commercial plantings and additionally seeds can be spread by birds and water (Bruce, Cameron & Harcombe, 1997; Jubinsky & Anderson, 1996; Renne, Gauthreaux Jr & Gresham, 2000). However, the current lack of documented established trees in natural areas of the EU and the fact that most trees are not reproductive for 3-8 years (Duke, 1983) suggests it is possible to prevent invasions of this species in the EU.

Prevention of introductions. The primary pathway of *intentional* introduction is the transport of seeds or plants for planting. A ban on the import of plants and seeds and restrictions on growing the species across member states is critical to prevent intentional introduction to the EU. *Unintentional* introduction of *T. sebifera* is relatively unlikely due to the fairly large seed size of the species. There is a low chance that *T. sebifera* would be a contaminant of crop or horticultural seed mixes or in bird seed. Seeds are not likely to be hitchhikers on people or animals.

Prevention of secondary spread. *Triadica sebifera* is not currently known to occur in the natural environment of the EU. However, if the species were to establish in the EU, it could be spread by water along creeks, streams, or drainage canals, or overland during heavy rain and flooding events (Bruce et al., 1997).

Measures to support early detection. Identifying newly established *T. sebifera* individuals or populations in natural areas will require activities to educate natural resource professionals, people interested in gardening and horticulture, and citizen scientists. Identification of the species is relatively easy due to its unique leaf structure and bark. Individual trees are reproductively mature at approximately 3-8 years old (Duke, 1983) so 'early detection' could include any activity within that timeframe.

Rapid eradication of new introductions. Early detection is important for preventing establishment of *T. sebifera* in natural areas because it would allow for eradication prior to seed production and dispersal. Manual or mechanical removal is the safest and most direct and effective method to rapidly eradicate newly established individuals or populations. All belowground parts of the plant need to be removed to maximize effectiveness.

Management of established populations. Manual, mechanical, and chemical options are available for management of older, more established populations of *T. sebifera*. The amount of effort, cost, and effectiveness varies widely by method.

Prevention of intentional introductions and spread – measures for preventing the species being introduced intentionally. **This table is repeated for each of the prevention measures identified.** *If the species is listed as an invasive alien species of Union concern, this table is not needed, as the measure applies anyway.*

<p>Measure description Provide a description of the measure, and identify its objective</p>	<p>A ban on keeping, importing, selling, breeding and growing the species (as would be required under Article 7. of the EU IAS Regulation 1143/2014).</p> <p><i>Triadica sebifera</i> is not known to currently occur in natural areas in the European Union but the species was listed as one of nearly 100 species that are likely to “arrive, establish, spread and have an impact on biodiversity or related ecosystem services in the EU over the next decade” (Roy et al., 2015). In the United States and Australia, where <i>T. sebifera</i> has been repeatedly introduced and widely planted (Bruce et al., 1997; Siemann & Rogers, 2003a,b; Camarillo, Stovall & Sunda, 2015), it has escaped cultivation and spread into natural areas and disturbed habitats, in particular open habitats near waterways (McCormick, 2005; Gan et al., 2009; Pile et al., 2017). It is unknown if invasions would have occurred had the species not been intentionally introduced and planted widely.</p> <p>Intentional introduction as an ornamental plant, and subsequent escape, is the most likely mechanism by which the species could eventually occur in natural environments in the EU. The species is desirable as an ornamental in Asia but there is not currently a significant commercial value in the EU. Historically, the oil (from seed) and tallow (from the seed covering) derived from the plant was used for various purposes from candle making to machine oil to wood products, and it is still used in Chinese medicine (USDA, 2008; Gao et al.,2016).</p> <p>Banning the import of <i>T. sebifera</i> to the EU, cultivation of the species, and trade among Member States will be critical for preventing intentional introductions. For example, as of October 2018, the species was purportedly available for purchase at multiple online seed suppliers in the UK¹, although the volume of sales and the distribution are not known.</p> <p>The objective of this measure is to prevent the intentional introduction of the species through the ornamental trade.</p>						
<p>Effectiveness of the measure Is it effective in relation to its objective? Has the measure previously worked, failed? Please select one of the categories of effectiveness (with an ‘X’), and provide a rationale, with supporting evidence and examples if possible.</p>	<p><i>Effectiveness of measures</i></p>	<p><i>Effective</i></p>	<p>X</p>	<p><i>Neutral</i></p>		<p><i>Ineffective</i></p>	
<p>Banning the import, cultivation, sale, and transport of <i>T. sebifera</i> should be highly effective for preventing the introduction and spread of the species in the EU. As unintentional introductions of the species is relatively unlikely (see Un-intentional introduction table below), banning intentional introduction is the primary mechanism for keeping the species out of EU natural areas.</p> <p><i>Rationale:</i> The effectiveness of a ban on import, sale, and transport is dependent on sufficient enforcement, including education regarding the ban, monitoring of sales and movement of species, and penalties for noncompliance. Assuming such actions are taken, an import ban should be highly effective. Little evidence exists to support the effectiveness of bans for preventing the introduction of species, likely because data on lack of species occurrences is usually not often reported.</p>							

¹ E.g. <https://www.rhs.org.uk/Plants/Nurseries-Search-Result?query=239677> [accessed 25/10/2018]

<p>Side effects (incl. potential) – both positive and negative i.e. positive or negative side effects of the measure on public health, environment including non-targeted species, etc.</p> <p>For each of the side effect types please select one of the impact categories (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<table border="1" data-bbox="645 188 1888 292"> <tr> <td><i>Environmental effects</i></td> <td><i>Positive</i></td> <td></td> <td><i>Neutral or mixed</i></td> <td>X</td> <td><i>Negative</i></td> <td></td> </tr> <tr> <td><i>Social effects</i></td> <td><i>Positive</i></td> <td></td> <td><i>Neutral or mixed</i></td> <td>X</td> <td><i>Negative</i></td> <td></td> </tr> <tr> <td><i>Economic effects</i></td> <td><i>Positive</i></td> <td></td> <td><i>Neutral or mixed</i></td> <td>X</td> <td><i>Negative</i></td> <td></td> </tr> </table> <p><i>Rationale:</i> Preventing the intentional introduction and spread of <i>T. sebifera</i> should have neutral (no) environmental side effects.</p> <p>A ban on import and transport will have neutral or very minimal social effects. The only conceivable effect would be on a cultural group that relies on the species for a specific social purpose. For example, although products derived from the species are used in Chinese medicine (Gao et al., 2016), the products are available elsewhere and do not need to be produced in the EU from plants grown in the EU.</p> <p>Preventing the import, cultivation, and spread of <i>T. sebifera</i> will have neutral or very little economic effect. The species was once very valuable for production of oil and tallow products, woodcrafts and furniture, and other products (Lin et al., 1958; Scheld & Cowles, 1981; USDA, 2008), which is why it was widely planted in the United States (Bruce et al., 1997; Siemann & Rogers, 2003a,b; McCormick, 2005; Camarillo et al., 2015), but there is not currently an industry that relies on <i>T. sebifera</i>. There is also no evidence that the species is widely traded in the EU for horticultural purposes.</p>	<i>Environmental effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>	X	<i>Negative</i>		<i>Social effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>	X	<i>Negative</i>		<i>Economic effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>	X	<i>Negative</i>	
<i>Environmental effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>	X	<i>Negative</i>																	
<i>Social effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>	X	<i>Negative</i>																	
<i>Economic effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>	X	<i>Negative</i>																	
<p>Acceptability to stakeholders e.g. impacted economic activities, animal welfare considerations, public perception, etc.</p> <p>Please select one of the categories of acceptability (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<table border="1" data-bbox="645 786 1888 850"> <tr> <td><i>Acceptability to stakeholders</i></td> <td><i>Acceptable</i></td> <td></td> <td><i>Neutral or mixed</i></td> <td>X</td> <td><i>Unacceptable</i></td> <td></td> </tr> </table> <p><i>Rationale:</i> Given how few stakeholders appear to deal in <i>T. sebifera</i> seeds, plants, or products, it is likely that there would be little to no negative response from stakeholders regarding a ban on importing, keeping, or selling the species.</p>	<i>Acceptability to stakeholders</i>	<i>Acceptable</i>		<i>Neutral or mixed</i>	X	<i>Unacceptable</i>															
<i>Acceptability to stakeholders</i>	<i>Acceptable</i>		<i>Neutral or mixed</i>	X	<i>Unacceptable</i>																	
<p>Additional cost information¹ When not already included above, or in the species Risk Assessment.</p> <ul style="list-style-type: none"> - the cost of inaction - the cost-effectiveness - the socio-economic aspects 	<p>Invasions of <i>T. sebifera</i> can significantly affect native plant communities, and may affect ecosystem nutrient cycling (Cameron & Spencer, 1989; Bruce et al., 1995; Jubinsky & Anderson, 1996; Bruce et al., 1997; Neyland & Meyer, 1997; Pile et al., 2017) thus the cost of inaction could be quite high. The species tolerates a wide range of environmental conditions from wetlands to grasslands to forests in its invasive range in North America (Webster, Jenkins & Jose, 2006; Langeland & Enloe, 2015) so many habitats in the EU could be vulnerable to invasions.</p> <p>The cost-effectiveness of implementing a ban on the species should be very high. Enforcement would require some resources but they would be shared across other species that are also banned. The primary cost would be in educational materials to inform citizen scientists and the public. The benefits could be quite high if a ban prevents invasions, which would be very difficult to determine (Zenni and Nunez, 2013), and is not always true (Diaz et al., 2012).</p>																					

<p>Include quantitative &/or qualitative data, and case studies (incl. from countries outside the EU).</p>	<p>Apart from the species in the ornamental plant trade in the EU, which is quite minimal, there are no current socio-economic benefits of importing or keeping the species in the EU. There is little information available on the value of the species in horticulture but it appears to be minimal given the few sites that list it for sale in the EU.</p>								
<p>Level of confidence on the information provided²</p> <p>Please select one of the confidence categories along with a statement to support the category chosen. See <i>Notes</i> section at the bottom of this document.</p> <p>NOTE – this is not related to the effectiveness of the measure</p>	<p><i>Inconclusive</i></p>		<p><i>Unresolved</i></p>		<p><i>Established but incomplete</i></p>	<p>X</p>	<p><i>Well established</i></p>		<p><i>Rationale:</i> Information on the establishment and spread of the species in natural areas of the United States is well documented and thought to be reliable. The impacts of <i>T. sebifera</i> invasions also have been researched extensively. What is less known is how introductions might occur in the EU – currently, a couple of online sites sell the species, but less is known about the availability of plants for planting, e.g., at local nurseries who might cultivate and propagate the species. Historically there has been some interest in keeping and cultivating the tree for landscaping purposes because it grows quickly and has somewhat attractive foliage and fall colour.</p>

<p>Prevention of <u>un-intentional</u> introductions and spread – measures for preventing the species being introduced un-intentionally (cf. Article 13 of the IAS Regulation). This table is repeated for each of the prevention measures identified.</p>	
<p>Measure description</p> <p>Provide a description of the measure, and identify its objective</p>	<p>Adoption of ‘weed free’ certification for bird feed and crop seeds</p> <p>There is little information available on the unintentional introductions and spread of <i>T. sebifera</i>. Given its large seed size, it is unlikely to be accidentally introduced to the EU. However, introduction as a contaminant of bird seed or in a crop seed mix might be possible. Multiple species of birds in North America readily feed on <i>T. sebifera</i> seeds (McCormick, 2005) and seeds remain viable for many years (Cameron, Glumac & Eschelman, 2000) so it is conceivable that viable seeds could be introduced as a contaminant and spread. However, no evidence exists that such an introduction has happened or is likely to occur.</p> <p>To prevent the unintentional introduction of <i>T. sebifera</i> as a contaminant of bird seed or crop seed for planting, the primary measure is to implement a required certification of “weed-free” seed for any seed that is imported for bird food or crop seeds for planting.</p> <p>The objective of this measure is to prevent the unintentional introduction of <i>T. sebifera</i> as a contaminant of bird food or crop seeds for planting.</p>
<p>Scale of application</p>	<p>This measure would need to be applied to all Member States that import seed from the U.S., Asia, Australia, and other areas where <i>T. sebifera</i> is present.</p>

<p>At what scale is the measure applied? What is the largest scale at which it has been successfully used? Please provide examples, with areas (km² or ha) if possible.</p>	<p>No information is available on the largest scale at which this measure has been successfully applied.</p>						
<p>Effectiveness of the measure Is it effective in relation to its objective? Has the measure previously worked, failed? Please select one of the categories of effectiveness (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<p><i>Effectiveness of measures</i></p>	<p><i>Effective</i></p>	<input type="checkbox"/>	<p><i>Neutral</i></p>	<input checked="" type="checkbox"/>	<p><i>Ineffective</i></p>	<input type="checkbox"/>
<p>No information is available on the effectiveness of this measure for preventing unintentional introduction of non-native species.</p>							
<p>Effort required e.g. period of time over which measure need to be applied to have results</p>	<p>No quantitative information is available, but it is expected that substantial effort would be required to implement a requirement that all imported bird food and crop seeds are “weed-free.” Non weed-free products would need to be banned and imported products would need to be examined to ensure effectiveness of the weed-free certification.</p>						
<p>Resources required¹ e.g. cost, staff, equipment etc.</p>	<p>The primary resource required would be staff to educate sellers and buyers that all products are required to be weed-free. Staff also would be needed to enforce the ban on non weed-free products and to ensure that weed-free products are actually free from contaminants. It has been shown that bans may not always be effective (Diaz et al., 2012).</p>						
<p>Side effects (incl. potential) – both positive and negative i.e. positive or negative side effects of the measure on public health, environment including non-targeted species, etc. For each of the side effect types please select one of the impact categories (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<p><i>Environmental effects</i></p>	<p><i>Positive</i></p>	<input checked="" type="checkbox"/>	<p><i>Neutral or mixed</i></p>	<input type="checkbox"/>	<p><i>Negative</i></p>	<input type="checkbox"/>
<p><i>Social effects</i></p>							
<p><i>Economic effects</i></p>							
<p>Banning non weed-free bird food and crop seed products would also prevent additional potential invasive alien plants from being introduced. If the cost of bird food or crop seeds increases due to a requirement that all products are certified weed-free then there may be negative social and economic effects, but no information is available on whether or not such effects would occur.</p>							
<p>Acceptability to stakeholders e.g. impacted economic activities, animal welfare considerations, public perception, etc.</p>	<p><i>Acceptability to stakeholders</i></p>	<p><i>Acceptable</i></p>	<input type="checkbox"/>	<p><i>Neutral or mixed</i></p>	<input checked="" type="checkbox"/>	<p><i>Unacceptable</i></p>	<input type="checkbox"/>
<p>Stakeholders that are interested in preventing invasions would likely be receptive to a requirement for weed-free products. However, businesses that import/sell bird food and crop seed products may have a negative response if the margin of profit is</p>							

Please select one of the categories of acceptability (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.	less for weed-free products but if the profit margin is actually larger for weed-free products, then they may have a positive response to the restriction.							
Additional cost information ¹ When not already included above, or in the species Risk Assessment. - implementation cost for Member States - the cost of inaction - the cost-effectiveness - the socio-economic aspects Include quantitative &/or qualitative data, and case studies (incl. from countries outside the EU).	No information available.							
Level of confidence on the information provided ² Please select one of the confidence categories along with a statement to support the category chosen. See <i>Notes</i> section at the bottom of this document. NOTE – this is not related to the effectiveness of the measure	<i>Inconclusive</i>	X	<i>Unresolved</i>		<i>Established but incomplete</i>		<i>Well established</i>	
Very little information is available that is relevant to the possibility that <i>T. sebifera</i> might be unintentionally introduced via bird food or crop seed contamination, or otherwise.								

Prevention of secondary spread of the species – measures for preventing the species spreading once they have been introduced (cf. Article 13 of the IAS Regulation). This table is repeated for each of the prevention measures identified.	
Measure description Provide a description of the measure, and identify its objective	Removal of populations occurring along waterways. If <i>T. sebifera</i> were to be introduced to natural areas in the EU, the primary mechanism for secondary spread would be movement by water. The species often occurs along waterways, including streams, creeks, rivers, ponds, and in wetlands (Bruce et al., 1997; McCormick, 2005; Pile et al., 2017) and seed can be transported by water via waterways or during flood events. Dispersal of <i>T. sebifera</i> seeds can also occur biotically via animals, primarily birds. For example, in the southern part of the US, the introduced starling has been shown to eat the seed and spread them widely (Renne, Gauthreaux & Gresham, 2000; Renne, Spira & Bridges, 2001; Renne et al., 2002; Jubinsky & Anderson, 1996).

	<p>To prevent secondary spread by water, <i>T. sebifera</i> that occurs near water or in lowlands prone to flooding should be rapidly eradicated prior to reproductive maturity (see Rapid eradication table below).</p> <p>The objective of this measure is to prevent secondary spread of <i>T. sebifera</i> through waterways or during flood events.</p>																					
<p>Scale of application At what scale is the measure applied? What is the largest scale at which it has been successfully used? Please provide examples, with areas (km² or ha) if possible.</p>	<p>For this measure to be effective at preventing secondary spread, all <i>T. sebifera</i> trees along waterways or in lowlands prone to flooding would need to be removed.</p>																					
<p>Effectiveness of the measure Is it effective in relation to its objective? Has the measure previously worked, failed?</p> <p>Please select one of the categories of effectiveness (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<table border="1" data-bbox="645 553 1888 619"> <tr> <td><i>Effectiveness of measures</i></td> <td><i>Effective</i></td> <td>X</td> <td><i>Neutral</i></td> <td></td> <td><i>Ineffective</i></td> <td></td> </tr> </table> <p>If all reproductive trees are removed, then the measure should be very effective at preventing secondary spread of <i>T. sebifera</i>.</p>	<i>Effectiveness of measures</i>	<i>Effective</i>	X	<i>Neutral</i>		<i>Ineffective</i>															
<i>Effectiveness of measures</i>	<i>Effective</i>	X	<i>Neutral</i>		<i>Ineffective</i>																	
<p>Effort required e.g. period of time over which measure need to be applied to have results</p>	<p>The effort required to eradicate newly established individuals is determined by the size of plants and the number of plants in a given area. Larger plants or more individuals would take more time and effort. Herbicide treatment would require the least effort.</p>																					
<p>Resources required¹ e.g. cost, staff, equipment etc.</p>	<p>As with measures for rapid eradication and management, resources to prevent secondary spread include staff to locate and conduct the eradication measures, and equipment and/or chemicals to remove <i>T. sebifera</i>.</p>																					
<p>Side effects (incl. potential) – both positive and negative i.e. positive or negative side effects of the measure on public health, environment including non-targeted species, etc.</p> <p>For each of the side effect types please select one of the impact categories (with an 'X'), and provide a</p>	<table border="1" data-bbox="645 1086 1888 1190"> <tr> <td><i>Environmental effects</i></td> <td><i>Positive</i></td> <td></td> <td><i>Neutral or mixed</i></td> <td>X</td> <td><i>Negative</i></td> <td></td> </tr> <tr> <td><i>Social effects</i></td> <td><i>Positive</i></td> <td></td> <td><i>Neutral or mixed</i></td> <td>X</td> <td><i>Negative</i></td> <td></td> </tr> <tr> <td><i>Economic effects</i></td> <td><i>Positive</i></td> <td></td> <td><i>Neutral or mixed</i></td> <td>X</td> <td><i>Negative</i></td> <td></td> </tr> </table> <p>See tables below on Rapid eradication.</p>	<i>Environmental effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>	X	<i>Negative</i>		<i>Social effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>	X	<i>Negative</i>		<i>Economic effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>	X	<i>Negative</i>	
<i>Environmental effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>	X	<i>Negative</i>																	
<i>Social effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>	X	<i>Negative</i>																	
<i>Economic effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>	X	<i>Negative</i>																	

<p>rationale, with supporting evidence and examples if possible.</p>									
<p>Acceptability to stakeholders e.g. impacted economic activities, animal welfare considerations, public perception, etc.</p> <p>Please select one of the categories of acceptability (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<p>Acceptability to stakeholders</p>	<p>Acceptable</p>	<p>X</p>	<p>Neutral or mixed</p>		<p>Unacceptable</p>		<p>There are no stakeholders in the EU who currently rely on <i>T. sebifera</i> so preventing secondary spread of the species via waterways should be acceptable.</p>	
<p>Additional cost information ¹ When not already included above, or in the species Risk Assessment.</p> <ul style="list-style-type: none"> - implementation cost for Member States - the cost of inaction - the cost-effectiveness - the socio-economic aspects <p>Include quantitative &/or qualitative data, and case studies (incl. from countries outside the EU).</p>	<p>No information available.</p>								
<p>Level of confidence on the information provided ²</p> <p>Please select one of the confidence categories along with a statement to support the category chosen. See <i>Notes</i> section at the bottom of this document.</p> <p>NOTE – this is not related to the effectiveness of the measure</p>	<p>Inconclusive</p>		<p>Unresolved</p>		<p>Established but incomplete</p>		<p>Well established</p>	<p>X</p>	<p>There are multiple reports (Bruce et al., 1997; McCormick, 2005; Pile et al., 2017) that <i>T. sebifera</i> occurs near waterways and may be transported by water. Therefore, removing reproductive trees from near water should prevent secondary spread of the species by water.</p>

Surveillance measures to support early detection - Measures to run an effective surveillance system for achieving an early detection of a new occurrence (cf. Article 16). This section assumes that the species is not currently present in a Member State, or part of a Member State's territory. **This table is repeated for each of the early detection measures identified.**

<p>Measure description Provide a description of the measure, and identify its objective</p>	<p>Surveillance of the natural environment by natural resources professionals, gardeners, and citizens</p> <p>The species is sold online and occurs under limited cultivation in the EU but the species is not known to occur in natural areas so early detection will be critical to prevent invasions. An effective surveillance system for early detection should include education of amateur and professional botanists, natural resource professionals, people interested in gardening and horticulture, and citizen scientists so they can identify and report occurrences of the species in natural areas.</p> <p>Education about <i>T. sebifera</i> identification could be accomplished through pamphlets, email, web sites (e.g., www.bsbi.org), or social media. Systematic surveillance of susceptible habitats could focus specifically on <i>T. sebifera</i> but are not likely to be cost-effective given the relatively unlikely chance the species would be encountered. Instead, regular biological recording for Atlases and Floras is likely to capture occurrences of the species given that they already exist at a reasonable scale across the EU (e.g., Pescott et al., 2015).</p> <p><i>Triadica sebifera</i> is relatively easy to identify based on its unique leaf morphology. There are no congeneric species within the EU. The Flora of China provides the following identification (edited): "Bark is grey, brown, and rough. Exudes a milky sap. Twigs slender. Petioles 2-6 cm long, with 2 sessile disc-shaped glands at the apex. Leaf-blades broadly rhombic-ovate, 2-7.5 x 1.5-7 cm, abruptly acutely acuminate, broadly cuneate to rounded, subtruncate at the base, entire, lateral nerves 7-12 pairs, glaucous beneath. Stipules 1-2 mm long, obtuse." <i>Triadica sebifera</i> flowers from April to June, producing both male and female flowers. Fruits are 1 cm three-lobed capsules expected to mature in the autumn. Seeds are 8 mm long and chalky white (covered by a white wax) (Flora of China, 1972; Bruce et al., 1997; Scheld, Cowles, Engler, Kleiman & Schultz, 1984). Providing pictures of leaves, seedlings, mature trees, flowers, and fruit should therefore be sufficient for land managers, gardeners, and citizen scientists to identify the species.</p> <p><i>Triadic sebifera</i> can tolerate a relatively wide range of habitat conditions in the introduced range such as moderately closed-canopy forests, open fields, wetlands, and coastal prairie, as well as disturbed sites such as old fields and margins such as fencerows (Bruce et al., 1997; McCormick, 2005; Webster et al., 2006; Camarillo et al., 2015; Langeland & Enloe, 2015). Such habitats should be the primary focus for surveillance efforts.</p> <p>The objective of this measure is to achieve early detection by utilizing efforts of people who might encounter <i>T. sebifera</i> in natural areas.</p>
<p>Scale of application At what scale is the measure applied? What is the largest scale at which it has been successfully used? Please</p>	<p><i>Triadica sebifera</i> has been cultivated and kept in botanical gardens in the EU for many years, including in the UK, France, Germany, the Netherlands, Italy, Portugal, Spain (Hortus Paddingtonensis, 1797; Antonelli, 1830; Dillwyn & Collinson, 1843; Hortus Collinsonianus, 1843; Banfi & Visconti, 2013; Batchelor, 2017), thus surveillance efforts need to cover these Member States as a priority. Within Member States where <i>T. sebifera</i> is known to occur, natural areas should be surveyed for the species, with a</p>

provide examples, with areas (km ² or ha) if possible.	particular focus on areas near cultivated <i>T. sebifera</i> . Preferably, <i>T. sebifera</i> would be included in surveillance efforts throughout the temperate EU Member States.						
<p>Effectiveness of the measure Is it effective in relation to its objective? Has the measure previously worked, failed?</p> <p>Please select one of the categories of effectiveness (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<p>Effectiveness of measures</p>	<p><i>Effective</i></p>		<p><i>Neutral</i></p>	<p>X</p>	<p><i>Ineffective</i></p>	
<p><i>Rationale:</i> It is very difficult to determine the effectiveness of particular surveillance measures for early detection because such efforts are relatively new (within the last couple decades) and documentation of efforts are not widely disseminated. Also, as the species is not widely distributed and is likely to be new to most managers and citizens, people participating in surveillance are less likely to immediately recognize seedlings, although the species is relatively unique (as compared to, e.g., a grass). However, amateur and professional botanists are likely to recognize the plant is unique and regular biological recording already occurs in much of the EU and is known to be effective at identifying new species occurrences (Pescott et al., 2015).</p>							
<p>Effort required e.g. period of time over which measure need to be applied to have results</p>	<p>If surveillance measures for <i>T. sebifera</i> were to be combined with efforts for other species of concern then the effort required for any one species, including <i>T. sebifera</i>, would be relatively low. Species already listed by the EU that occur in the same or similar habitats include <i>Asclepias syriaca</i>, <i>Heracleum mantegazzianum</i>, <i>Microstegium vimineum</i>, and <i>Pueraria lobata</i>.</p>						
<p>Resources required¹ e.g. cost, staff, equipment etc.</p>	<p>Surveillance of the natural environment for <i>T. sebifera</i> should require few resources – simply educational materials such as web sites, smartphone apps (which often are already produced for other IAS), and educational handouts. Overall, the implementation cost is expected to be minimal because regular biological recording already occurs in many areas of the EU (e.g., Pescott et al., 2015).</p>						
<p>Side effects (incl. potential) – both positive and negative i.e. positive or negative side effects of the measure on public health, environment including non-targeted species, etc.</p> <p>For each of the side effect types please select one of the impact categories (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<p>Environmental effects</p>	<p><i>Positive</i></p>	<p>X</p>	<p><i>Neutral or mixed</i></p>		<p><i>Negative</i></p>	
<p>Social effects</p>		<p><i>Positive</i></p>	<p>X</p>	<p><i>Neutral or mixed</i></p>		<p><i>Negative</i></p>	
<p>Economic effects</p>		<p><i>Positive</i></p>		<p><i>Neutral or mixed</i></p>	<p>X</p>	<p><i>Negative</i></p>	
<p><i>Rationale:</i> Surveillance for <i>T. sebifera</i> could have significant positive environmental side effects if search efforts in natural areas result in identification of other invasive species or documentation of other environmental issues that need to be reported to natural resource managers.</p> <p>The positive social effects of surveillance are that citizen scientists will be more aware of environmental issues and they will have the opportunity to spend time in natural areas during search efforts.</p> <p>No economic side effects of <i>T. sebifera</i> surveillance efforts are expected.</p>							

<p>Acceptability to stakeholders e.g. impacted economic activities, animal welfare considerations, public perception, etc.</p> <p>Please select one of the categories of acceptability (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<table border="1" data-bbox="645 153 1888 220"> <tr> <td><i>Acceptability to stakeholders</i></td> <td><i>Acceptable</i></td> <td>X</td> <td><i>Neutral or mixed</i></td> <td></td> <td><i>Unacceptable</i></td> <td></td> </tr> </table> <p><i>Rationale:</i> Surveillance of natural areas for <i>T. sebifera</i> is expected to be acceptable to stakeholders because it will not impact economic activities or have consequences for animal welfare. Public perception should be neutral or possibly positive because participants will experience and become more aware of the natural environment.</p>	<i>Acceptability to stakeholders</i>	<i>Acceptable</i>	X	<i>Neutral or mixed</i>		<i>Unacceptable</i>		
<i>Acceptability to stakeholders</i>	<i>Acceptable</i>	X	<i>Neutral or mixed</i>		<i>Unacceptable</i>				
<p>Additional cost information¹ When not already included above, or in the species Risk Assessment.</p> <ul style="list-style-type: none"> - implementation cost for Member States - the cost of inaction - the cost-effectiveness - the socio-economic aspects <p>Include quantitative &/or qualitative data, and case studies (incl. from countries outside the EU).</p>	<p>As described under Rapid eradication below, the cost of inaction could be very high if surveillance efforts are not implemented for <i>T. sebifera</i> as invasions can spread rapidly once individuals reach maturity and the species can have significant impacts on biodiversity and ecosystem functions.</p> <p>If surveillance efforts for <i>T. sebifera</i> are combined with surveys of other species the measure should be cost-effective.</p>								
<p>Level of confidence on the information provided²</p> <p>Please select one of the confidence categories along with a statement to support the category chosen. See <i>Notes</i> section at the bottom of this document.</p> <p>NOTE – this is not related to the effectiveness of the measure</p>	<table border="1" data-bbox="645 847 1928 914"> <tr> <td><i>Inconclusive</i></td> <td></td> <td><i>Unresolved</i></td> <td></td> <td><i>Established but incomplete</i></td> <td>X</td> <td><i>Well established</i></td> <td></td> </tr> </table> <p><i>Rationale:</i> There is growing evidence on the application of surveillance, and citizen science as a tool for early detection.</p>	<i>Inconclusive</i>		<i>Unresolved</i>		<i>Established but incomplete</i>	X	<i>Well established</i>	
<i>Inconclusive</i>		<i>Unresolved</i>		<i>Established but incomplete</i>	X	<i>Well established</i>			

Rapid eradication for new introductions - Measures to achieve eradication at an early stage of invasion, after an early detection of a new occurrence (cf. Article 17). This section assumes that the species is not currently present in a Member State, or part of a Member State's territory. **This table is repeated for each of the eradication measures identified.**

<p>Measure description Provide a description of the measure, and identify its objective</p>	<p>Manual or mechanical removal</p> <p>If a new occurrence of <i>T. sebifera</i> is found in the EU in the natural environment, the most direct and effective method to rapidly eradicate newly established individuals or populations is by physical removal. To achieve eradication, it is critical to remove individuals before they are mature and produce seed. The species has been documented to reach reproductive maturity in as little as one year under ideal greenhouse conditions (Jubinsky & Anderson, 1996), although typically, plants in natural areas are mature in 3-8 years (Duke, 1983). Given the relatively rapid maturation of <i>T. sebifera</i>, newly identified invasions will require quick action to prevent seed production and achieve eradication.</p> <p><i>Triadica sebifera</i> creates a long-lived seedbank, which may complicate management efforts and further motivate rapid removal to prevent second spread. Cameron et al. (2000) showed that germination success peaked after two years in their experiment but that there was little loss in viability (only 3%) even after seven years.</p> <p>To remove an individual tree or small populations prior to maturity, mechanical or manual removal is preferred. Although multiple herbicides have been tested and utilized effectively on <i>T. sebifera</i> (see below), it is simpler, easier, and in many ways safer to remove seedling trees without the use of herbicides. Seedlings may be hand pulled or dug (i.e., “grubbed”) out but in either case, the entire plant must be removed because the species readily resprouts from root fragments. Machinery such as bulldozers or scrapers also can be used but, again, such techniques must remove all plant parts to prevent resprouting and ensure eradication.</p> <p>The objective of this measure is to rapidly eradicate new introductions of <i>T. sebifera</i> using manual and mechanical methods.</p>						
<p>Scale of application At what scale is the measure applied? What is the largest scale at which it has been successfully used? Please provide examples, with areas (km² or ha) if possible.</p>	<p>Removal of <i>T. sebifera</i> with mechanical or manual methods is labour intensive and can be time consuming. No information is available on the scale at which this measure has been applied successfully but it is likely to be restricted to areas of approximately 1 ha or less and with a limited number and size of stems, due to the difficulty of implementation and need for complete removal of all plant parts to achieve eradication. Invaded areas larger than 1 ha will require chemical control methods, especially if plants are mature and seed has been dispersed.</p>						
<p>Effectiveness of the measure Is it effective in relation to its objective? Has the measure previously worked, failed?</p> <p>Please select one of the categories of effectiveness (with an ‘X’), and provide a rationale, with supporting evidence and examples if possible.</p>	<p><i>Effectiveness of measures</i></p>	<p><i>Effective</i></p>	<p>X</p>	<p><i>Neutral</i></p>		<p><i>Ineffective</i></p>	
<p><i>Rationale:</i></p> <p>Removal of entire plants can be highly effective for eradication. However, because of the scale at which manual or mechanical removal can be implemented is quite limited, chemical control must be considered for larger infestations. Simply cutting a plant at the soil surface is ineffective because “cutting results in the immediate production of multiple small, independent shoots” (Jubinsky, 1993). Cut-stump application of herbicides will greatly reduce eradication success so when to use manual or mechanical methods and when to implement chemical control should be carefully considered.</p>							

<p>Effort required e.g. period of time over which measure need to be applied to have results</p>	<p>The intensity of effort required to eradicate a population with mechanical or manual methods is determined by the size of plants and the number of plants in a given area. Larger plants will require more work to pull or dig out and the larger the plant, the greater the likelihood some plant parts will be missed and the plant will resprout. Larger areas or more dense populations will increase the chances that an individual will be missed and potentially reach reproductive maturity. Because seed dispersal of <i>T. sebifera</i> can occur through both bird and water dispersal, and mature individuals can produce up to 100,000 seeds annually (Jubinsky & Anderson, 1996; Bruce et al., 1997; USDA, 2008), eradication will require exponentially more effort if seed production occurs.</p>																											
<p>Resources required¹ e.g. cost, staff, equipment etc.</p>	<p>The cost of <i>T. sebifera</i> eradication using manual and mechanical methods is relatively low, although perhaps higher than the cost of using chemical herbicides. The cost is predominantly for personnel to locate, identify, map, and remove individuals. Gloves and hand tools (e.g., shovels, spades) are all that is needed for seedlings and small trees. Larger trees will require chemical control (see below).</p>																											
<p>Side effects (incl. potential) – both positive and negative i.e. positive or negative side effects of the measure on public health, environment including non-targeted species, etc.</p> <p>For each of the side effect types please select one of the impact categories (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<table border="1"> <tr> <td>Environmental effects</td> <td>Positive</td> <td></td> <td>Neutral or mixed</td> <td></td> <td>Negative</td> <td>X</td> </tr> <tr> <td>Social effects</td> <td>Positive</td> <td></td> <td>Neutral or mixed</td> <td>X</td> <td>Negative</td> <td></td> </tr> <tr> <td>Economic effects</td> <td>Positive</td> <td></td> <td>Neutral or mixed</td> <td>X</td> <td>Negative</td> <td></td> </tr> </table>	Environmental effects	Positive		Neutral or mixed		Negative	X	Social effects	Positive		Neutral or mixed	X	Negative		Economic effects	Positive		Neutral or mixed	X	Negative							
Environmental effects	Positive		Neutral or mixed		Negative	X																						
Social effects	Positive		Neutral or mixed	X	Negative																							
Economic effects	Positive		Neutral or mixed	X	Negative																							
<p><i>Rationale:</i> There may be negative environmental side effects because of disturbance of the natural environment when larger individuals are dug out, but the effects can be minimized by replacing soil and taking care to not disturb native vegetation. The amount of disturbance would depend on the number and size of individuals. Few, if any, social, or economic effects are expected with manual or mechanical removal of <i>T. sebifera</i>.</p>																												
<p>Acceptability to stakeholders e.g. impacted economic activities, animal welfare considerations, public perception, etc.</p> <p>Please select one of the categories of acceptability (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<table border="1"> <tr> <td>Acceptability to stakeholders</td> <td>Acceptable</td> <td>X</td> <td>Neutral or mixed</td> <td></td> <td>Unacceptable</td> <td></td> </tr> </table>	Acceptability to stakeholders	Acceptable	X	Neutral or mixed		Unacceptable																					
Acceptability to stakeholders	Acceptable	X	Neutral or mixed		Unacceptable																							
<p><i>Rationale:</i> Manual or mechanical eradication of <i>T. sebifera</i> is likely to be acceptable to stakeholders because there is no commercial value of the species in the EU and little evidence that it is used in landscaping.</p>																												
<p>Additional cost information¹ When not already included above, or in the species Risk Assessment.</p>	<p>Given the rapid spread and significant impacts of <i>T. sebifera</i> invasions in the Southern US (see Prevention above) the cost of not eradicating newly established populations could be profound (McCormick, 2005). The population growth and long-distance dispersal potential of <i>T. sebifera</i> once a population establishes indicates that early detection and rapid eradication are absolutely</p>																											

<ul style="list-style-type: none"> - implementation cost for Member States - the cost of inaction - the cost-effectiveness - the socio-economic aspects <p>Include quantitative &/or qualitative data, and case studies (incl. from countries outside the EU).</p>	<p>critical for preventing widespread invasions. Once individuals or populations are found – and assuming plants are still small – manual or mechanical removal should be cost-effective with few socio-economic impacts.</p>								
<p>Level of confidence on the information provided²</p> <p>Please select one of the confidence categories along with a statement to support the category chosen. See <i>Notes</i> section at the bottom of this document.</p> <p>NOTE – this is not related to the effectiveness of the measure</p>	<i>Inconclusive</i>		<i>Unresolved</i>		<i>Established but incomplete</i>		<i>Well established</i>	X	
		<p><i>Rationale:</i></p> <p>It is well-known that larger, mature <i>T. sebifera</i> trees readily resprout when felled. However, there is a high level of confidence that at the earliest stage of invasion when plants are still small, manual or mechanical removal can be highly effective.</p>							

<p>Rapid eradication for new introductions - Measures to achieve eradication <u>at an early stage of invasion</u>, after an early detection of a new occurrence (cf. Article 17). This section assumes that the species is not currently present in a Member State, or part of a Member State’s territory. This table is repeated for each of the eradication measures identified.</p>	
<p>Measure description</p> <p>Provide a description of the measure, and identify its objective</p>	<p>Chemical control</p> <p>While rapid eradication of <i>T. sebifera</i> seedlings and small trees with manual or mechanical measures can be effective, larger trees that cannot be completely removed by hand or with hand tools or machinery will require chemical control. In addition, the difficulty of removing individual trees means that rapid eradication of larger infestations with many individuals will be impractical without chemicals.</p> <p>Chemical control for rapid eradication, can be achieved with foliar, basal bark, or cut stump application of herbicides. Foliar applications of herbicides for seedlings and small trees should occur during the growing season (July through October in most Member States). Products for foliar application should contain imazapyr (1% water-solution of products such as Arsenal™ or Habitat™), fosamine-containing herbicide (30% water solution of Krenite STM), or triclopyr ester-containing herbicide (2% water-solution of Garlon 4™) (Jubinsky & Anderson, 1996; Langeland & Enloe, 2015; NSW Government Factsheet, 2017)</p>

	<p>If it is acceptable for standing dead trees to remain in place, then basal bark application of herbicides is easier and more cost-effective than either foliar applications or cut-stump treatment. A 20% dilution in oil of triclopyr ester- containing herbicide such as Garlon 4TM or Pathfinder IITM is effective for trees up to 15cm.</p> <p>For cut stump application a 50% dilution of triclopyr ester or triclopyr amine containing herbicide such as Garlon 3ATM can be used immediately after cutting. Stumps may coppice and resprouts will need to be treated according to cut-stump recommendations. Herbicides that contain imazapyr can help to reduce resprouting (Jubinsky & Anderson, 1996; Langeland & Enloe, 2015; NSW Government Factsheet, 2017).</p> <p>Recent research showed that herbicides containing aminocyclopyrachlor or fluroxypyr were much more effective in controlling resprouting than products containing the traditionally used triclopyr amine and ester formulations (Enloe, Loewenstein, Streett & Lauer, 2015). Therefore, these herbicides should be used.</p> <p>It is illegal to use an herbicide in a manner inconsistent with the label's instructions; therefore, read the label carefully and follow the instructions. It is important to note that EU/national/local legislation on the use of plant protection products and biocides needs to be respected and authorities should check to ensure chemicals are licensed for use in their respective countries/regions.</p> <p>The objective of this measure is to rapidly eradicate new introductions of <i>T. sebifera</i> using chemical control methods.</p>							
<p>Scale of application At what scale is the measure applied? What is the largest scale at which it has been successfully used? Please provide examples, with areas (km² or ha) if possible.</p>	<p>No information is available on the scale at which the chemical control of <i>T. sebifera</i> or other woody invaders has been successfully implemented. But, given the method of applying either as basal bark or cut stump, widespread applications could be feasible given sufficient resources.</p>							
<p>Effectiveness of the measure Is it effective in relation to its objective? Has the measure previously worked, failed?</p> <p>Please select one of the categories of effectiveness (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<table border="1" data-bbox="645 1043 1888 1110"> <tr> <td data-bbox="645 1043 925 1110"><i>Effectiveness of measures</i></td> <td data-bbox="925 1043 1151 1110"><i>Effective</i></td> <td data-bbox="1151 1043 1211 1110" style="text-align: center;">X</td> <td data-bbox="1211 1043 1509 1110"><i>Neutral</i></td> <td data-bbox="1509 1043 1570 1110"></td> <td data-bbox="1570 1043 1832 1110"><i>Ineffective</i></td> <td data-bbox="1832 1043 1888 1110"></td> </tr> </table> <p><i>Rationale:</i> Enloe et al. (2015) conducted an extensive study of the effectiveness of chemical herbicide treatments to kill <i>T. sebifera</i>, including their effects on root and stump sprouting post treatment. They found that many treatments were more than 80% effective in preventing resprouts:</p> <p>“Aminopyralid reduced total sprouting by 95% compared with the untreated control and was different from all other herbicide treatments. Fluroxypyr reduced total sprouting by 79 and 88% at the low and high rates, respectively. Imazamox reduced total sprouting by 61 and 64% at the low and high rates, respectively, and triclopyr reduced total sprouting by 37%.”</p>	<i>Effectiveness of measures</i>	<i>Effective</i>	X	<i>Neutral</i>		<i>Ineffective</i>	
<i>Effectiveness of measures</i>	<i>Effective</i>	X	<i>Neutral</i>		<i>Ineffective</i>			

	<p>For basal bark treatment, multiple chemicals were effective against <i>T. sebifera</i>, including for inhibition of resprouting. The most effective chemical was aminocyclopyrachlor (Enloe et al., 2015).</p>																					
<p>Effort required e.g. period of time over which measure need to be applied to have results</p>	<p>The effort required to apply cut stump and basal bark chemical treatments is relatively low compared to manual or mechanical treatments. Basal bark application of chemicals is significantly easier than cut stump treatment because individuals do not need to be cut prior to treatment. When individuals are seedlings or small trees, the difference in application effort is relatively minor but for large trees, particularly those requiring chainsaws, the difference in effort is substantial.</p> <p>Chemical measures can be quite effective for rapid eradication of <i>T. sebifera</i> with only a single treatment but when resprouts occur they must be treated again. It has been suggested that infested sites be revisited for at least 3-5 years to ensure eradication (Jubinsky & Anderson, 1996; Bruce et al., 1997; Meyer, 2011).</p>																					
<p>Resources required¹ e.g. cost, staff, equipment etc.</p>	<p>Basal bark application requires fewer resources than cut stump application because trees are left in place. For basal bark, chemical, oil surfactant, sprayer, and personal protective equipment (e.g., gloves) are required. For cut stump application of chemicals, tools are also needed to cut the trees – hand pruners or loppers for smaller trees and a chainsaw for larger (i.e., >5cm) trees.</p> <p>More staff resources are required for cut stump than basal bark application and cut stump treatments may require specialized training, such as the use of chainsaws. In either case, the applicator would need safety training in the use of chemical herbicides.</p>																					
<p>Side effects (incl. potential) – both positive and negative i.e. positive or negative side effects of the measure on public health, environment including non-targeted species, etc.</p> <p>For each of the side effect types please select one of the impact categories (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<table border="1" data-bbox="645 979 1888 1082"> <tr> <td><i>Environmental effects</i></td> <td><i>Positive</i></td> <td></td> <td><i>Neutral or mixed</i></td> <td>X</td> <td><i>Negative</i></td> <td></td> </tr> <tr> <td><i>Social effects</i></td> <td><i>Positive</i></td> <td></td> <td><i>Neutral or mixed</i></td> <td>X</td> <td><i>Negative</i></td> <td></td> </tr> <tr> <td><i>Economic effects</i></td> <td><i>Positive</i></td> <td></td> <td><i>Neutral or mixed</i></td> <td>X</td> <td><i>Negative</i></td> <td></td> </tr> </table> <p><i>Rationale:</i></p> <p>Cut stump and basal bark chemical measures to rapidly eradicate <i>T. sebifera</i> are likely to have few side effects. Few, if any, negative environmental side effects are expected because the amount of chemical used is relatively small (Enloe et al., 2015) and is applied in a very targeted manner directly to the base of trees or cut stumps.</p> <p>Given that currently there are no significant social or economic uses of <i>T. sebifera</i> in the EU, any social or economic side effects are expected to be minimal or non-existent.</p>	<i>Environmental effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>	X	<i>Negative</i>		<i>Social effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>	X	<i>Negative</i>		<i>Economic effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>	X	<i>Negative</i>	
<i>Environmental effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>	X	<i>Negative</i>																	
<i>Social effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>	X	<i>Negative</i>																	
<i>Economic effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>	X	<i>Negative</i>																	

<p>Acceptability to stakeholders e.g. impacted economic activities, animal welfare considerations, public perception, etc.</p> <p>Please select one of the categories of acceptability (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<table border="1" data-bbox="645 153 1888 220"> <tr> <td>Acceptability to stakeholders</td> <td><i>Acceptable</i></td> <td></td> <td><i>Neutral or mixed</i></td> <td>X</td> <td><i>Unacceptable</i></td> <td></td> </tr> </table> <p><i>Rationale:</i> Both basal bark and cut stump application of herbicides for rapid eradication should be acceptable for stakeholders because the amount of herbicide is relatively low and there should be a low number of treated plants. However, the public may object to rapid eradication of <i>T. sebifera</i> if the treated trees are prominent such as near hiking trails or parks and they are removed or if trees are treated with basal bark applications and left standing.</p>	Acceptability to stakeholders	<i>Acceptable</i>		<i>Neutral or mixed</i>	X	<i>Unacceptable</i>		
Acceptability to stakeholders	<i>Acceptable</i>		<i>Neutral or mixed</i>	X	<i>Unacceptable</i>				
<p>Additional cost information¹ When not already included above, or in the species Risk Assessment.</p> <ul style="list-style-type: none"> - implementation cost for Member States - the cost of inaction - the cost-effectiveness - the socio-economic aspects <p>Include quantitative &/or qualitative data, and case studies (incl. from countries outside the EU).</p>	<p>As described above for rapid eradication using manual or mechanical control, the cost of inaction could be quite high if small populations spread and create larger invasions. Chemical eradication measures are often thought to be highly cost-effective but the costs of detection, application, and monitoring have not been quantified.</p> <p>One concern that might need to be addressed is that the physical removal of trees or leaving standing dead trees may not be acceptable to local people who pass by or use natural areas. These concerns may be addressed with educational campaigns.</p>								
<p>Level of confidence on the information provided²</p> <p>Please select one of the confidence categories along with a statement to support the category chosen. See <i>Notes</i> section at the bottom of this document.</p> <p>NOTE – this is not related to the effectiveness of the measure</p>	<table border="1" data-bbox="645 847 1924 914"> <tr> <td><i>Inconclusive</i></td> <td></td> <td><i>Unresolved</i></td> <td></td> <td><i>Established but incomplete</i></td> <td></td> <td><i>Well established</i></td> <td>X</td> </tr> </table> <p><i>Rationale:</i> It is well established that cut stump and basal bark chemical measures are effective for killing <i>T. sebifera</i> (e.g., Enloe et al., 2015), thus these measures are expected to be effective for rapid eradication of the species in the EU.</p>	<i>Inconclusive</i>		<i>Unresolved</i>		<i>Established but incomplete</i>		<i>Well established</i>	X
<i>Inconclusive</i>		<i>Unresolved</i>		<i>Established but incomplete</i>		<i>Well established</i>	X		

Management - Measures to achieve management of the species once it has become widely spread within a Member State, or part of a Member State's territory. (cf. Article 19), i.e. **not** at an early stage of invasion (see Rapid eradication table above). These measures can be aimed at eradication, population control or containment of a population of the species. **This table is repeated for each of the management measures identified.**

<p>Measure description Provide a description of the measure, and identify its objective</p>	<p>Chemical control</p> <p><i>Triadica sebifera</i> is not known to currently occur in natural areas within the EU. Thus, surveillance, early detection, and rapid eradication should be top priorities. However, if widespread invasions of <i>T. sebifera</i> are found in a Member State(s), the optimal management measure is chemical control. Other measures, including fire, alteration of hydrology, and mechanical removal (mulching) have been attempted where the species has invaded over large areas but they are not effective because of the ability of the <i>T. sebifera</i> to resprout from roots. Because little information is available on each of these measures, no specific tables have been produced for these measures. Here the focus is on the same chemical treatments of cut stumps and basal bark applications outlined above under Rapid Eradication.</p> <p>In brief, chemical control can be achieved by primarily foliar (seedlings or small trees only), basal bark, or cut stump application of herbicides. Foliar applications of herbicides for seedlings and small trees should occur during the growing season (July through October in most EU Member States). Products for foliar application should contain imazapyr (1% water-solution of products such as ArsenalTM or HabitatTM), fosamine-containing herbicide (30% water solution of Krenite STM), or triclopyr ester-containing herbicide (2% water-solution of Garlon 4TM) (Jubinsky & Anderson, 1996; Langeland & Enloe, 2015; NSW Government Factsheet, 2017). If it is acceptable for standing dead trees to remain in place, then basal bark application of herbicides is easier and more cost-effective than either foliar applications or cut-stump treatment. A 20% dilution in oil of triclopyr ester- containing herbicide such as Garlon 4TM or Pathfinder IITM is effective for trees up to 15cm. For cut stump application a 50% dilution of triclopyr ester or triclopyr amine containing herbicide such as Garlon 3ATM can be used immediately after cutting. Stumps may coppice and resprouts will need to be treated according to cut-stump recommendations. Herbicides that contain imazapyr can help to reduce resprouting (Jubinsky & Anderson, 1996, Langeland & Enloe, 2015, NSW Government Factsheet, 2017).</p> <p>Recent research showed that herbicides containing aminocyclopyrachlor or fluroxypyr were much more effective in controlling resprouting than products containing the traditionally used triclopyr amine and ester formulations (Enloe et al., 2015).</p> <p>It is illegal to use an herbicide in a manner inconsistent with the label's instructions; therefore, read the label carefully and follow the instructions. It is important to note that EU/national/local legislation on the use of plant protection products and biocides needs to be respected and authorities should check to ensure chemicals are licensed for use in their respective countries/regions.</p> <p>The objective of this measure is to manage established invasions of <i>T. sebifera</i> using chemical control methods.</p>
<p>Scale of application At what scale is the measure applied? What is the largest scale at which it has been successfully used? Please provide examples, with areas (km² or ha) if possible.</p>	<p>No information is available on the scale at which management of <i>T. sebifera</i> with chemical measures has been applied. However, as treatment is conducted on a per-individual basis there is little benefit for large-scale compared to local control. That is, there is not significant economy of scale as there would be with a measure that uses larger equipment for more extensive invasions (e.g. aerial application of herbicides). Similarly, there is little disadvantage of larger scale management of <i>T. sebifera</i> using basal bark and cut stump chemical methods other than the amount of area that needs to be mapped, treated, and monitored.</p>

<p>Effectiveness of the measure Is it effective in relation to its objective? Has the measure previously worked, failed?</p> <p>Please select one of the categories of effectiveness (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<table border="1" data-bbox="645 153 1888 220"> <tr> <td><i>Effectiveness of measures</i></td> <td><i>Effective</i></td> <td>X</td> <td><i>Neutral</i></td> <td></td> <td><i>Ineffective</i></td> <td></td> </tr> </table> <p><i>Rationale:</i> Multiple fact sheets (e.g., McCormick, 2005; NSW Government Factsheet, 2017; Meyer, 2011) and the extensive study by Enloe et al. (2015) document the effectiveness of basal bark and cut stump chemical measures for the management of <i>T. sebifera</i>.</p>	<i>Effectiveness of measures</i>	<i>Effective</i>	X	<i>Neutral</i>		<i>Ineffective</i>															
<i>Effectiveness of measures</i>	<i>Effective</i>	X	<i>Neutral</i>		<i>Ineffective</i>																	
<p>Effort required e.g. period of time over which measure need to be applied to have results</p>	<p>The effort required for management of <i>T. sebifera</i> using basal bark and cut stump chemical methods scales directly with the size of the infestation. Treatments per individual tree require relatively little effort but invasions covering multiple hectares would require significant person hours, equipment, and chemicals, particularly if trees cannot be treated with basal bark applications and left as standing dead. Results will be apparent within the first year but invaded areas will need to be retreated for resprouts for at least 3-5 years for full control of a population.</p>																					
<p>Resources required¹ e.g. cost, staff, equipment etc.</p>	<p>Resources required for management of established stands of <i>T. sebifera</i> invasions are similar to resources required for rapid eradication, except that it is likely that more people, time, and chemicals are needed depending on the area invaded, density of <i>T. sebifera</i> individuals, and size distribution of trees.</p>																					
<p>Side effects (incl. potential) – both positive and negative i.e. positive or negative side effects of the measure on public health, environment including non-targeted species, etc.</p> <p>For each of the side effect types please select one of the impact categories (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<table border="1" data-bbox="645 786 1888 887"> <tr> <td><i>Environmental effects</i></td> <td><i>Positive</i></td> <td></td> <td><i>Neutral or mixed</i></td> <td>X</td> <td><i>Negative</i></td> <td></td> </tr> <tr> <td><i>Social effects</i></td> <td><i>Positive</i></td> <td></td> <td><i>Neutral or mixed</i></td> <td>X</td> <td><i>Negative</i></td> <td></td> </tr> <tr> <td><i>Economic effects</i></td> <td><i>Positive</i></td> <td></td> <td><i>Neutral or mixed</i></td> <td>X</td> <td><i>Negative</i></td> <td></td> </tr> </table> <p><i>Rationale:</i> Broad scale management of <i>T. sebifera</i> could have both positive and negative environmental side effects. It is well-established that invasions of the species in the southern US cause substantial biodiversity loss and alter ecosystem functions (Bruce et al., 1995; Bruce et al., 1997; Jubinsky & Anderson, 1996; Cameron & Spencer, 1989; Neyland & Meyer, 1997; Pile et al., 2017). If such invasions also occurred in the EU, similar effects are expected. Removal of those invasions would thus have some positive environmental benefit. However, it is not known to what extent native biodiversity in the EU would utilize the new habitat created by large stands of <i>T. sebifera</i>. If invasive stands create habitat or provide a significant food source then removal of those stands could have negative environmental side effects. Such effects are not understood for <i>T. sebifera</i>.</p> <p>Given the current lack of economic activity or cultural or social uses associated with <i>T. sebifera</i> in the EU, no social or economic side effects are expected.</p>	<i>Environmental effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>	X	<i>Negative</i>		<i>Social effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>	X	<i>Negative</i>		<i>Economic effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>	X	<i>Negative</i>	
<i>Environmental effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>	X	<i>Negative</i>																	
<i>Social effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>	X	<i>Negative</i>																	
<i>Economic effects</i>	<i>Positive</i>		<i>Neutral or mixed</i>	X	<i>Negative</i>																	
<p>Acceptability to stakeholders</p>	<table border="1" data-bbox="645 1313 1888 1380"> <tr> <td><i>Acceptability to stakeholders</i></td> <td><i>Acceptable</i></td> <td>X</td> <td><i>Neutral or mixed</i></td> <td></td> <td><i>Unacceptable</i></td> <td></td> </tr> </table>	<i>Acceptability to stakeholders</i>	<i>Acceptable</i>	X	<i>Neutral or mixed</i>		<i>Unacceptable</i>															
<i>Acceptability to stakeholders</i>	<i>Acceptable</i>	X	<i>Neutral or mixed</i>		<i>Unacceptable</i>																	

<p>e.g. impacted economic activities, animal welfare considerations, public perception, etc.</p> <p>Please select one of the categories of acceptability (with an 'X'), and provide a rationale, with supporting evidence and examples if possible.</p>	<p><i>Rationale:</i> No significant economic activities are associated with <i>T. sebifera</i> in the EU currently so chemical management measures are expected to be acceptable to stakeholders.</p>							
<p>Additional cost information ¹ When not already included above, or in the species Risk Assessment.</p> <ul style="list-style-type: none"> - implementation cost for Member States - the cost of inaction - the cost-effectiveness - the socio-economic aspects <p>Include quantitative &/or qualitative data, and case studies (incl. from countries outside the EU).</p>	<p>As with the cost of inaction for rapid eradication, the cost of not conducting management of <i>T. sebifera</i> could be substantial given the known ecological impacts of invasions in the Southern US.</p>							
<p>Level of confidence on the information provided ²</p> <p>Please select one of the confidence categories along with a statement to support the category chosen. See <i>Notes</i> section at the bottom of this document.</p> <p>NOTE – this is not related to the effectiveness of the measure</p>	<i>Inconclusive</i>	<input type="checkbox"/>	<i>Unresolved</i>	<input type="checkbox"/>	<i>Established but incomplete</i>	<input type="checkbox"/>	<i>Well established</i>	<input checked="" type="checkbox"/>
<p><i>Rationale:</i> There is a reasonable amount of information on the effectiveness, costs, and resources required for basal bark and cut stump chemical management techniques for <i>T. sebifera</i>, thus there is a high level of confidence in the information provided here.</p>								

<p>Bibliography ³</p> <p>See guidance section</p>
<p>Antonelli, G. (1830). Nuovo Dizionario universale tecnologico o di arti e mestieri e della economia industriale e commerciale compilato dai signori Lenormand etc. Prima trad. <i>Italiana</i>, Volume 1.</p>

- Banfi, E., & Visconti, A. (2013). L'Orto di Brera alla fine della dominazione asburgica e durante l'età napoleonica, *Atti Soc. it. Sci. nat. Museo civ. Stor. nat. Milano*, 154 (II): 173-264
- Batchelor, R. (2017). John Bradby Blake, the Chinese tallow tree and the infrastructure of botanical experimentation. *Curtis's Botanical Magazine* 2017 vol. 34 (4): pp. 402-426
- Bruce, K. A., Cameron, G. N., & Harcombe, P. A. (1995). Initiation of a new woodland type on the Texas coastal prairie by the Chinese tallow tree (*Sapium sebiferum* (L.) Roxb.). *Bulletin of the Torrey Botanical Club*, 215-225.
- Bruce, K. A., Cameron, G. N., Harcombe, P. A., & Jubinsky, G. (1997). Introduction, impact on native habitats, and management of a woody invader, the Chinese tallow tree, *Sapium sebiferum* (L.) Roxb. *Natural Areas Journal*, 255-260.
- CABI (2018). *Sapium sebiferum*. Distribution. <http://www.cabi.org/isc/datasheet/48351>
- Camarillo, S. A., Stovall, J. P., & Sunda, C. J. (2015). The impact of Chinese tallow (*Triadica sebifera*) on stand dynamics in bottomland hardwood forests. *Forest Ecology and Management*, 344, 10-19.
- Cameron, G. N., & Spencer, S. R. (1989). Rapid leaf decay and nutrient release in a Chinese tallow forest. *Oecologia*, 80(2), 222-228.
- Cameron, G. N., Glumac, E. G., & Eschelman, B. D. (2000). Germination and dormancy in seeds of *Sapium sebiferum* (Chinese tallow tree). *Journal of Coastal Research*, 16(2): 391-395.
- Diaz, S., Smith, J. R., Zaleski, S. F., & Murray, S. N. (2012). Effectiveness of the California state ban on the sale of *Caulerpa* species in aquarium retail stores in southern California. *Environmental management*, 50(1), 89-96.
- Dillwyn, L. W., & Collinson, P. (1843). *Hortus Collinsonianus: An account of the plants cultivated by the late Peter Collinson. Arranged alphabetically according to their modern names, from the catalogue of his garden, and other manuscripts*, 64 pp. - Publisher Printed by W. C. Murray and D. Rees. Book digitized by Google from the library of Oxford University and uploaded to the Internet Archive by user tpb. - Source: <http://books.google.com/books?id=gEQAAAAAQAAJ&oe=UTF-8>
- Duke, J. A. (1983). *Sapium sebiferum* (L.) Roxb., [Online]. In: Handbook of energy crops. In: NewCrops: New crop resource online. West Lafayette, IN: Purdue University, Department of Horticulture and Landscape Architecture, Center for New Crops & Plant Products (Producer). Available: https://hort.purdue.edu/newcrop/duke_energy/Sapium_sebiferum.html accessed August 2018.
- EddMapps (2018). Chinese Tallow Distribution. <https://www.eddmaps.org/distribution/usstate.cfm?sub=3079>
- Enloe, S. F., Loewenstein, N. J., Streett, D., & Lauer, D. K. (2015). Herbicide treatment and application method influence root sprouting in Chinese tallowtree (*Triadica sebifera*). *Invasive Plant Science and Management*, 8(2), 160-168.
- Flora of China (1972). Available: <http://www.efloras.org/index.aspx>
- Gan, J., Miller, J. H., Wang, H., & Taylor, J. W. (2009). Invasion of tallow tree into southern US forests: influencing factors and implications for mitigation. *Canadian journal of forest research*, 39(7), 1346-1356.
- GBIF (2018). *Triadica sebifera* Distribution. Available: <https://www.gbif.org/species/3054399>
- Gao, R., Su, Z., Yin, Y., & Li, S. (2016). Germplasm, chemical constituents, biological activities, utilization, and control of Chinese tallow (*Triadica sebifera* (L.) Small). *Biol Invasions*, 18:809-829
- Hortus collinsonianus (1843). An Account of the Plants cultivated by the late Peter Collinson, Esq., F.R.S. Available: https://books.google.fr/books?id=gEQAAAAAQAAJ&pg=PR9&source=gbs_selected_pages&cad=2#v=onepage&q&f=false
- Hortus Paddingtonensis (1797). A catalogue of plants cultivated in the garden of J. Symmons, Esq., Paddington-House. Available at: <https://archive.org/details/b22386890>
- ITIS Standard Report. Retrieved July 20, 2018 from the Integrated Taxonomic Information System on-line database, <http://www.itis.gov>.
- Jubinsky, G. (1993). A review of the literature: *Sapium sebiferum* Roxb. Report No. TSS 93-03. Florida Dept. Natur. Resour., Bur. Aq. Plant Manage., Tallahassee.
- Jubinsky, G., & Anderson, L. C. (1996). The invasive potential of Chinese tallow-tree (*Sapium sebiferum* Roxb.) in the Southeast. *Castanea*, 226-231.

- Langeland, K. A., & Enloe, S. (2015) Natural area weeds: Chinese tallow (*Sapium sebiferum* L.), [Online]. In: EDIS (Electronic Data Information Source). Publication #SS-AGR-45. Gainesville, FL: University of Florida, IFAS (Institute of Food and Agricultural Science) Extension (Producer). Available: <http://edis.ifas.ufl.edu/ag148>
- Lin, W. C., Chen, A. C., Tseng, C. J., & Hwang, S. G. (1958). An investigation and study of Chinese tallow tree in Taiwan (*Sapium sebiferum*, Roxb.). *Bulletin of Taiwan Forestry Research Institute*, 57, 32-37.
- McCormick, C. (2005). Chinese Tallow Management Plan for Florida: Recommendations from The Chinese Tallow Task Force. The Florida Exotic Pest Plant Council.
- Meyer, R. (2011). *Triadica sebifera*. In: Fire Effects Information System, [Online]. U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station, Fire Sciences Laboratory (Producer). Available: <http://www.fs.fed.us/database/feis/> [2017, August 30]
- Neyland, R., & Meyer, H. A. (1997). Species diversity of Louisiana chenier woody vegetation remnants. *Journal of the Torrey Botanical Society*, 254-261.
- NSW Government, Australia (2017). NSW WeedWise: Chinese tallow tree (*Triadica sebifera*). Department of Primary Industries. Factsheet: <http://weeds.dpi.nsw.gov.au/Weeds/Details/38>
- Pescott, O. L., Walker, K. J., Pocock, M. J., Jitlal, M., Outhwaite, C. L., Cheffings, C. M., ... & Roy, D. B. (2015). Ecological monitoring with citizen science: the design and implementation of schemes for recording plants in Britain and Ireland. *Biological Journal of the Linnean Society*, 115(3), 505-521.
- Pile, L. S., Wang, G. G., Knapp, B. O., Walker, J. L., & Stambaugh, M. C. (2017). Chinese tallow (*Triadica sebifera*) invasion in maritime forests: The role of anthropogenic disturbance and its management implication. *Forest Ecology and Management*, 398, 10-24.
- Renne, I. J., Barrow, W. C., Randall, J., Lori, A., & Bridges, W. C. (2002). Generalized avian dispersal syndrome contributes to Chinese tallow tree (*Sapium sebiferum*, Euphorbiaceae) invasiveness. *Diversity and Distributions*, 8(5), 285-295.
- Renne, I. J., Gauthreaux Jr, S. A., & Gresham, C. A. (2000). Seed dispersal of the Chinese tallow tree (*Sapium sebiferum* (L.) Roxb.) by birds in coastal South Carolina. *The American Midland Naturalist*, 144(1), 202-215.
- Renne, I. J., Spira, T. P., & Bridges Jr, W. C. (2001). Effects of habitat, burial, age and passage through birds on germination and establishment of Chinese tallow tree in coastal South Carolina. *Journal of the Torrey Botanical Society*, 109-119.
- Roy, H. E., Adriaens, T., Aldridge, D. C., Bacher, S., Bishop, J. D. D., Blackburn, T. M., Branquart, E., Brodie, J., Carboneras, C., Cook, E. J., Copp, G. H., Dean, H. J., Eilenberg, J., Essl, F., Gallardo, B., Garcia, M., García-Berthou, E., Genovesi, P., Hulme, P. E., Kenis, M., Kerckhof, F., Kettunen, M., Minchin, D., Nentwig, W., Nieto, A., Pergl, J., Pescott, O., Peyton, J., Preda, C., Rabitsch, W., Roques, A., Rorke, S., Scalera, R., Schindler, S., Schönrogge, K., Sewell, J., Solarz, W., Stewart, A., Tricarico, E., Vanderhoeven, S., van der Velde, G., Vilà, M., Wood, C. A., & Zenetos, A. (2015). *Invasive Alien Species - Prioritising prevention efforts through horizon scanning* ENV.B.2/ETU/2014/0016. European Commission.
- Scheld, H. W., & Cowles, J.R. (1981). Woody biomass potential of the Chinese tallow tree. *Economic Botany*, 35(4), 391-397.
- Scheld, H. W., Cowles, J. R., Engler, C. R., Kleiman, R., & Schultz Jr, E. B. (1984). Seeds of the Chinese tallow tree as a source of chemicals and fuels. In *Fuels and chemicals from oil seeds: Technology and policy options*. AAAS Selected symposium (Vol. 91, pp. 81-101).
- Siemann, E., & Rogers, W.E. (2003a). Herbivory, disease, recruitment limitation, and success of alien and native tree species. *Ecology*. 84(6): 1489-1505.
- Siemann, E., & Rogers, W. E. (2003b). Increased competitive ability of an invasive tree may be limited by an invasive beetle. *Ecological Applications*. 13(6): 1503-1507.
- USDA (2008). Plant guide: Chinese tallow tree--*Triadica sebifera* (L.) Small, [Online]. In: PLANTS Database--fact sheets & plant guides. Baton Rouge, LA: U.S. Department of Agriculture, Natural Resources Conservation Service, National Plant Data Center (Producer). Available: http://plants.usda.gov/plantguide/pdf/pg_trse6.pdf
- Webster, C. R., Jenkins, M. A., & Jose, S. (2006). Woody invaders and the challenges they pose to forest ecosystems in the eastern United States. *Journal of Forestry* 104:366-374.
- Zenni, R. D., & Nuñez, M. A. (2013). The elephant in the room: the role of failed invasions in understanding invasion biology. *Oikos*, 122(6), 801-815.

Notes

1. Costs information. The assessment of the potential costs shall describe those costs quantitatively and/or qualitatively depending on what information is available. This can include case studies from across the Union or third countries.

2. Level of confidence²: based on the quantity, quality and level of agreement in the evidence.



- **Well established:** comprehensive meta-analysis³ or other synthesis or multiple independent studies that agree.
- **Established but incomplete:** general agreement although only a limited number of studies exist but no comprehensive synthesis and, or the studies that exist imprecisely address the question.
- **Unresolved:** multiple independent studies exist but conclusions do not agree.
- **Inconclusive:** limited evidence, recognising major knowledge gaps

3. Citations and bibliography. The APA formatting style for citing references in the text and in the bibliography is used.

e.g. Peer review papers will be written as follows:

In text citation: (Author & Author, Year)

In bibliography: Author, A. A., & Author, B. B. (Publication Year). Article title. *Periodical Title*, Volume(Issue), pp.-pp.

(see <http://www.waikato.ac.nz/library/study/referencing/styles/apa>)

² Assessment of confidence methodology is taken from IPBES. 2016. Guide on the production and integration of assessments from and across all scales (IPBES-4-INF-9), which is adapted from Moss and Schneider (2000).

³ A statistical method for combining results from different studies which aims to identify patterns among study results, sources of disagreement among those results, or other relationships that may come to light in the context of multiple studies.